


[Print this Page for Your Records](#)
[Close Window](#)
Control/Tracking Number: 2025-A-375-ESHG

Activity: ESHG Abstract

Current Date/Time: 3/19/2025 11:32:32 PM

Identifying genetic variants associated with recurrent pregnancy loss in the Uzbek population

Author Block: Anastasiya Punko¹, Abror Abdurakhimov², Yuliya Kapralova¹, Darya Zakirova¹, Abdushukur Rakhmatullayev¹, Gulnoza Esonova¹, Xurshid Meylikov¹, Alisher Abdullaev¹, Dilbar Dalimova¹, Shahlo Turdikulova¹.

¹Center for Advanced Technologies, Tashkent, Uzbekistan, ²Institute of Biophysics and Biochemistry at the National University of Uzbekistan, Tashkent, Uzbekistan.

Abstract:

Background: Recurrent pregnancy loss (RPL) is defined as the loss of two or more pregnancies before the 20th week of gestation and affects approximately 15% of pregnancies. Despite advances in infertility diagnostics, 35–60% of RPL cases remain unexplained, suggesting genetic, epigenetic, and environmental contributions. In this study, we analysed the genetic factors associated with RPL among women in Uzbekistan.

Material and Methods: This study included 161 patients diagnosed with RPL and 511 healthy individuals. Genetic screening was performed using the Global Screening Array v.3.0. Hardy–Weinberg equilibrium test and logistic regression analyses were conducted using the SNPAssoc software package to identify statistically significant associations between genetic variants and disease risk under an additive genetic model.

Results: Several genetic variants showed significant associations with RPL. Protective effects were observed for rs4871372 (*LOC105375725*, $p = 3.7 \times 10^{-5}$, OR = 0.58) and rs2704035 (*SULF1*, $p = 3.9 \times 10^{-4}$, OR = 0.63). Increased RPL risk was linked to rs4916848 (*ADGRV1*, $p = 1.8 \times 10^{-6}$, OR = 1.82) and rs478993 (*TENM4*, $p = 4.5 \times 10^{-4}$, OR = 1.58).

Conclusion: Our study identified novel genetic associations with RPL in Uzbek women. The lncRNA *LOC105375725* (rs4871372) may function as an epigenetic regulator, while rs4916848 likely acts as an enhancer. Additionally, *TENM4* has been implicated in mesoderm induction during early embryogenesis. Therefore, these variants can be used as diagnostic and prognostic markers. Further research is needed to understand their functional role of these variants in RPL pathogenesis.

Author Disclosure Information:

A. Punko: None. **A. Abdurakhimov:** None. **Y. Kapralova:** None. **D. Zakirova:** None. **A. Rakhmatullayev:** None. **G. Esonova:** None. **X. Meylikov:** None. **A. Abdullaev:** None. **D. Dalimova:** None. **S. Turdikulova:** None.

Topic (Complete): 18. Large scale genomics (GWAS) and other -omics association studies

Keyword (Complete): recurrent pregnancy loss ; Uzbek population

Presentation Preference (Complete): Hybrid Poster

Additional information (Complete):

Select one of the options below: Regular Abstract

Please specify all grants & references related to your abstract: : None

Fellowships & Awards (Complete):

Do you wish to apply for any of the Fellowships or Awards?: No

I apply for the Conference Fellowship for European Countries: No

I apply for the Fellowship for Non-European Countries: No

I apply for the Fellowship of Excellence: No

I apply for the Early Career Award: No

Attached Files: No Files Attached

Status: Complete

ESHG Conference – Wiener Medizinische Akademie GmbH

Alser Strasse 4, A-1090 Vienna, Austria – E: conference@eshg.org – T: +43 1 405 13 83 14 – F: +43 1 407 82 74

 Feedback

Powered by [cOASIS](#), The Online Abstract Submission and Invitation System SM
© 1996 – 2025 [CTI Meeting Technology](#). All rights reserved. [Privacy Policy](#).