

Calibration of Stochastic Volatility Models using Particle Markov Chain Monte Carlo Methods

Jonas Hallgren¹

¹Department of Mathematics
KTH Royal Institute of Technology
Stockholm, Sweden

BFS 2012
June 21, 2012
Sydney, Australia



TABLE OF CONTENTS

INTRODUCTION

Financial Times Series

Model Proposal

PARAMETER ESTIMATION

Bayesian Inference

Parameter Simulation

Sequential Monte Carlo Methods

Filters and Smoothers

Monte Carlo Integration

Sequential Importance Sampling

Particle MCMC

Particle Marginal Metropolis Hastings

Unbiased Parallel Metropolis Hastings




Simulations and Results

Estimations

Prediction



LOGRETURNS



$$Y_k = \log \left(\frac{S_k}{S_{k-1}} \right)$$

MODEL PROPOSAL

$$Y_k = \beta e^{\frac{1}{2}X_k} u_k, \quad (1)$$

$$X_k = \alpha X_{k-1} + \sigma w_k, \quad (2)$$

$$(u_k, w_k) \sim \mathcal{N}(0, \Sigma), \quad (3)$$

$$\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix} \quad (4)$$

BAYESIAN INFERENCE

Bayesian inference, view the parameter as a random variable

Observation:

$$Y \mid \theta^* = \theta \sim p(y \mid \theta^* = \theta), \quad \theta^* \sim p(\theta) \quad (5)$$

Parameter posterior distribution:

$$p(\theta \mid y) = \frac{p(y \mid \theta)p(\theta)}{\int_{\Theta} p(y \mid \theta')p(d\theta')} \propto p(y \mid \theta)p(\theta) \quad (6)$$

Example:

$$p(\beta \mid \alpha, \sigma, \rho, x_{0:n}, y_{0:n}) \propto p(\beta, \alpha, \sigma, \rho, x_{0:n}, y_{0:n}) \quad (7)$$



POSTERIOR FOR β

$$p(\beta|\alpha, \sigma, \rho, x_{0:n}, y_{0:n}) \propto p(\beta, \alpha, \sigma, \rho, x_{0:n}, y_{0:n}) \quad (8)$$

$$= p(x_{0:n}, y_{0:n}|\beta, \alpha, \rho, \sigma)p(\beta)p(\alpha, \rho, \sigma) \quad (9)$$

$$\propto p(x_{0:n}, y_{0:n}|\beta, \alpha, \rho, \sigma)p(\beta) \quad (10)$$

$$= p(x_n, y_n|\beta, \alpha, \rho, \sigma, x_{0:n-1}, y_{0:n-1}) \quad (11)$$

$$\times p(x_{0:n-1}, y_{0:n-1}|\beta, \alpha, \rho, \sigma)p(\beta) \quad (12)$$

$$= p(\beta)p(y_0, x_0|\beta, \alpha, \rho, \sigma) \\ \times \prod_{k=1}^n p(x_k, y_k|\beta, \alpha, \rho, \sigma, x_{k-1})$$

POSTERIOR FOR β

$$p(x, y) = \frac{1}{|\beta| \sigma 2\pi \sqrt{1 - \rho^2}} \quad (13)$$

$$\times \exp \left(- \frac{\left[\left(\frac{y}{\beta e^{\frac{1}{2}x}} \right)^2 + \left(\frac{x - \alpha x_{k-1}}{\sigma} \right)^2 - 2\rho \frac{y(x - \alpha x_{k-1})}{\sigma \beta e^{\frac{1}{2}x}} \right]}{2(1 - \rho^2)} - \frac{1}{2}x \right)$$

PRIOR SELECTION

$$p(\beta) = \frac{1}{\beta^2} \quad (14)$$

$$p(\alpha) = (\alpha + 1)^{\delta-1} (1 - \alpha)^{\gamma-1} \quad (15)$$

$$p(\rho) = \frac{1}{2} \quad (16)$$

$$p(\sigma) = \frac{1}{\sigma^2 \sigma^{2(t/2-1)}} e^{-\frac{1}{2\sigma^2} \tilde{S}_0} \quad (17)$$

Example:

$$p(\beta \mid \alpha, \sigma, \rho, x_{0:n}, y_{0:n}) \propto p(x_{0:n}, y_{0:n} \mid \alpha, \beta, \rho, \sigma) p(\beta) \quad (18)$$



IDEA

- ▶ Simulate the parameters from the posterior distributions!

THE GIBBS SAMPLER ¹

1. For the first iteration choose $\xi^{(0)} = \{x_{0:n}^{(0)}, \theta^{(0)}\}$ arbitrarily
2. For $k = 1, 2, \dots, N$, draw random samples

$$2.1 \quad x_{0:n}^{(k)} \sim p_X(\cdot \mid \theta^{(k-1)}, y_{0:n})$$

$$2.2 \quad \theta_1^{(k)} \sim p_X(\cdot \mid x_{0:n}^{(k)}, \theta^{(k-1)}, y_{0:n})$$

$$\vdots$$

$$\theta_D^{(k)} \sim p_X(\cdot \mid x_{0:n}^{(k)}, \theta_1^{(k)}, \theta_2^{(k)}, \dots, \theta_D^{(k-1)}, y_{0:n})$$

Now as N tend to infinity, the sequence $\{\xi^{(k)}\}_{k=0}^N$ will have p_X as its stationary distribution.

New problem: How do we sample θ and x ?

¹Geman (1984)

METROPOLIS-HASTINGS SAMPLER ²

Choose θ_0 arbitrarily, then for $k = 1, \dots, N$

1. Sample $\theta^* \sim q(\cdot \mid \theta^{(k)})$
2. With probability

$$1 \wedge \frac{p(\theta^*)q(\theta^{(k)} \mid \theta^*)}{p(\theta^{(k)})q(\theta^* \mid \theta^{(k)})} \quad (19)$$

set $\theta^{(k+1)} = \theta^*$, otherwise set $\theta^{(k+1)} = \theta^{(k)}$

²Metropolis et. al. (1953), Hastings (1970)

SEQUENTIAL MONTE CARLO (PARTICLE FILTER)

$$\phi_k \triangleq p(x_k \mid y_{0:k}) \quad (20)$$

Propose

$$\phi_{k+1}(\tilde{\xi}) = \frac{\int l_k(\xi, \tilde{\xi}) \phi_k(\xi) d\xi}{\int \phi_k(\xi) \int l_k(\xi, \tilde{\xi}) d\tilde{\xi} d\xi} \quad (21)$$



OUR MODEL

In our setting:

$$\phi_{k+1} = p(x_{k+1}, y_{0:k+1})/p(y_{0:k+1}) \quad (22)$$

$$\propto \int p(y_{k+1} \mid x_{k+1}, x_k, y_{0:k}) \quad (23)$$

$$\begin{aligned} &\times p(x_{k+1} \mid x_k, y_{0:k}) p(x_k, y_{0:k}) dx_k \\ &= \int p(y_{k+1} \mid x_{k:k+1}) p(x_{k+1} \mid x_k) p(x_k \mid y_{0:k}) p(y_{0:k}) dx_k \end{aligned} \quad (24)$$

$$= \int p(y_{k+1} \mid x_{k:k+1}) p(x_{k+1} \mid x_k) \phi_k p(y_{0:k}) dx_k \quad (25)$$

$$= \int G(y_{k+1} \mid x_{k:k+1}) Q(x_{k+1} \mid x_k) \phi_k p(y_{0:k}) dx_k \quad (26)$$



SUMMARIZED

Filter:

$$\phi_{k+1} = \frac{\int G(y_{k+1} | x_{k:k+1}) Q(x_{k+1} | x_k) \phi_k dx_k}{\int \int G(y_{k+1} | x_{k:k+1}) Q(x_{k+1} | x_k) \phi_k dx_k dx_{k+1}} \quad (27)$$

Smoother:

$$\begin{aligned} \phi_{0:k+1|k+1} &= p(x_{0:k+1} | y_{0:k+1}) \\ &= \frac{\int G(y_{k+1} | x_{k:k+1}) Q(x_{k+1} | x_k) \phi_{0:k|k} dx_{0:k}}{\int \int G(y_{k+1} | x_{k:k+1}) Q(x_{k+1} | x_k) \phi_{0:k|k} dx_{0:k} dx_{0:k+1}} \end{aligned} \quad (28)$$

MONTE CARLO INTEGRATION

We want to evaluate:

$$\mu(f) = \int f(x) \frac{d\mu}{d\nu}(x) \nu(dx) \quad (29)$$

We use the estimate:

$$N^{-1} \sum_{j=1}^N f(\xi^j) \frac{d\mu}{d\nu}(\xi^j) \xrightarrow[N \rightarrow \infty]{a.s.} \mu(f) \quad (30)$$



SEQUENTIAL IMPORTANCE SAMPLING

1. Sampling: for $k = 0, 1, \dots$

Draw $\tilde{\xi}_{k+1}^1, \dots, \tilde{\xi}_{k+1}^N \mid \tilde{\xi}_{0:k}^1, \dots, \tilde{\xi}_{0:k}^N$

- 1.1 Compute the importance weights

$$\omega_{k+1}^j = \omega_k^j g_{k+1}(\tilde{\xi}_{k+1}^j) \quad (31)$$


2. Resampling: Draw N particles from the N -sized population where the probability of selecting particle j is

$$\frac{\omega_{k+1}^j}{\sum_s \omega_{k+1}^s} \quad (32)$$

3. Update the trajectory: Copy the resampled particles trajectories and replace the ones we discarded.



EXAMPLE



RECAP

- ▶ Object: Model the price
- ▶ Need parameters
 - ▶ Need X trajectories

Which we now have!

$$Y_k = \beta e^{\frac{1}{2}X_k} u_k, \quad (33)$$

$$X_k = \alpha X_{k-1} + \sigma w_k, \quad (34)$$

$$(u_k, w_k) \sim \mathcal{N}(0, \Sigma), \quad (35)$$

$$\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix} \quad (36)$$



THE GIBBS SAMPLER

1. For the first iteration choose $\xi^{(0)} = \{x_{0:n}^{(0)}, \theta^{(0)}\}$ arbitrarily
2. For $k = 1, 2, \dots, N$, draw random samples
 - 2.1 $x_{0:n}^{(k)} \sim p_X(\cdot \mid \theta^{(k-1)}, y_{0:n})$
 - 2.2 $\theta_1^{(k)} \sim p_X(\cdot \mid x_{0:n}^{(k)}, \theta^{(k-1)}, y_{0:n})$
 - \vdots
 - $\theta_D^{(k)} \sim p_X(\cdot \mid x_{0:n}^{(k)}, \theta_1^{(k)}, \theta_2^{(k)}, \dots, \theta_D^{(k-1)}, y_{0:n})$

Now as N tend to infinity, the sequence $\{\xi^{(k)}\}_{k=0}^N$ will have p_X as its stationary distribution

INTRODUCTION

Financial Times Series
Model Proposal

PARAMETER ESTIMATION

Bayesian Inference
Parameter Simulation

Sequential Monte Carlo Methods

Filters and Smoothers
Monte Carlo Integration
Sequential Importance Sampling

Particle MCMC

Particle Marginal Metropolis Hastings
Unbiased Parallel Metropolis Hastings

Simulations and Results

Estimations
Prediction



METROPOLIS-HASTINGS SAMPLER ²

Choose θ_0 arbitrarily, then for $k = 1, \dots, N$

1. Sample $\theta^* \sim q(\cdot \mid \theta^{(k)})$
2. With probability

$$1 \wedge \frac{p(\theta^*)q(\theta^{(k)} \mid \theta^*)}{p(\theta^{(k)})q(\theta^* \mid \theta^{(k)})} \quad (19)$$

set $\theta^{(k+1)} = \theta^*$, otherwise set $\theta^{(k+1)} = \theta^{(k)}$

²Metropolis et. al. (1953), Hastings (1970)

PARTICLE MARGINAL METROPOLIS HASTINGS³

1. Initialization, $k = 0$
 - 1.1 Set θ_0 arbitrarily
 - 1.2 Run an SMC algorithm targeting $p_{\theta^{(0)}}(x_{1:T}|y_{1:T})$, sample our first trajectory of particles $\tilde{\xi}_{1:T}^{(0)} \sim \hat{p}_{\theta^{(0)}}(\cdot|y_{1:T})$ and denote the marginal likelihood by \hat{p}_{θ_0}
2. For iteration $k \geq 1$
 - 2.1 Sample $\theta^* \sim q(\cdot | \theta_{k-1})$
 - 2.2 Run an SMC algorithm targeting $p_{\theta^*}(x_{1:T} | y_{1:T})$, sample the trajectory of particles as in 1.2
 - 2.3 With probability

$$1 \wedge \frac{\hat{p}_{\theta^*}(y_{1:T})p(\theta^*)q(\theta_{k-1} | \theta^*)}{\hat{p}_{\theta_{k-1}}(y_{1:T})p(\theta_{k-1})q(\theta^* | \theta_{k-1})} \quad (37)$$

put $\theta_k = \theta^*$, $\xi_{1:T}^{(k)} = \xi_{1:T}^*$, and $p_{\theta_k}(y_{1:T}) = p_{\theta^*}$

³Andrieu et. al. (2010)

UNBIASED PARALLEL METROPOLIS HASTINGS

Choose θ_0^m arbitrarily, then for $k = 1, \dots, N$

1. For each of the C cores (where $\theta^{(k)} = \theta_k^m$):
 - 1.1 Sample $\theta^* \sim q(\cdot \mid \theta^{(k)})$
 - 1.2 With probability

$$1 \wedge \frac{p(\theta^*)q(\theta^{(k)} \mid \theta^*)}{p(\theta^{(k)})q(\theta^* \mid \theta^{(k)})} \quad (38)$$

set $\theta^{(k+1)} = \theta^*$, otherwise set $\theta^{(k+1)} = \theta^{(k)}$

2. Iterate through the C cores, take the first accepted sample and put $\theta_{k+\gamma}^m$ equal to it. Put $\theta_{k:k+\gamma-1}^m = \theta_k^m$. Throw away all samples after that. If no sample is accepted, $\theta_{k:C}^m = \theta_k^m$



UNBIASED PARALLEL METROPOLIS HASTINGS

- ▶ Roughly $(\text{Acceptance rate})^{-1}$ times faster as numbers of cores grow large
- ▶ Easy to implement



MODEL PROPOSAL


$$Y_k = \beta e^{\frac{1}{2}X_k} u_k, \quad (1)$$

$$X_k = \alpha X_{k-1} + \sigma w_k, \quad (2)$$

$$(u_k, w_k) \sim \mathcal{N}(0, \Sigma), \quad (3)$$

$$\Sigma = \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix} \quad (4)$$

SIMULATION OF S&P500



PREDICTION

Dataset	Model	RMSE (10^{-3})
GBP/USD	<i>SVOL</i>	11.706
GBP/USD	$SVOL_{\rho=0}$	11.714
BIDU	<i>SVOL</i>	20.188
BIDU	$SVOL_{\rho=0}$	20.232
S&P500	<i>SVOL</i>	252.94
S&P500	$SVOL_{\rho=0}$	252.93
XBC/USD	<i>SVOL</i>	5.5762
XBC/USD	$SVOL_{\rho=0}$	5.5920

Any remark, question or suggestion is welcomed!

`jonas@math.kth.se`

`http://www.math.kth.se/~jhallg/`

