

# Last Time

# Last Time

$$\begin{aligned} & \rightarrow \max_a R(s,a) \quad \leftarrow \text{Myopic} \\ & \max_a \underbrace{R(s,a) + \gamma E[U(s')]}_{Q(s,a)} \\ & \rightarrow \max_a Q^*(s,a) \end{aligned}$$

- How do we reason about the **future consequences** of actions in an MDP?

# Last Time

- How do we reason about the **future consequences** of actions in an MDP?
- What are the basic **algorithms for solving MDPs**?

Policy Iteration  
Evaluate policy  
Improve policy

Value Iteration  
Improve  $V$  estimate

# Guiding Questions

# Guiding Questions

- Does value iteration always converge?
- Is the value function unique?

# Value Iteration: The Bellman Operator

# Value Iteration: The Bellman Operator

Algorithm: Value Iteration

while  $\|V - V'\|_\infty < \epsilon$

$V \leftarrow V'$

$V' \leftarrow B[V]$

return  $V'$

# Value Iteration: The Bellman Operator

Algorithm: Value Iteration

while  $\|V - V'\|_\infty < \epsilon$

$V \leftarrow V'$

$V' \leftarrow B[V]$

return  $V'$

$$B[V](s) = \max_{a \in A} (R(s, a) + \gamma E[V(s')])$$



# Value Iteration Convergence

# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

# Metrics

# Metrics

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

# Metrics

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$

# Metrics


Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$

# Metrics

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$



# Contraction Mappings



# Contraction Mappings

Definition: A *contraction mapping* on metric space  $(M, d)$  is a function  $f : M \rightarrow M$  satisfying

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for some  $\alpha$ ,  $0 \leq \alpha \leq 1$  and all  $x$  and  $y$  in  $M$ .


# Contraction Mappings

Definition: A *contraction mapping* on metric space  $(M, d)$  is a function  $f : M \rightarrow M$  satisfying

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for some  $\alpha$ ,  $0 \leq \alpha \leq 1$  and all  $x$  and  $y$  in  $M$ .

Definition:  $x^*$  is said to be a *fixed point* of  $f$  if  $f(x^*) = x^*$ .



# Contraction Mappings

Definition: A *contraction mapping* on metric space  $(M, d)$  is a function  $f : M \rightarrow M$  satisfying

$$d(f(x), f(y)) \leq \alpha d(x, y)$$

for some  $\alpha$ ,  $0 \leq \alpha \leq 1$  and all  $x$  and  $y$  in  $M$ .

Definition:  $x^*$  is said to be a *fixed point* of  $f$  if  $f(x^*) = x^*$ .

Script: contraction\_mapping.jl

# Banach's Theorem

# Banach's Theorem

Theorem (Banach): If  $f$  is a contraction mapping on metric space  $(M, d)$ , then

1.  $f$  has a single, unique fixed point  $x^*$ .
2. If  $\{x_k\}$  is a sequence defined by  $x_{k+1} = f(x_k)$ , then  $\lim_{k \rightarrow \infty} x_k = x^*$ .

# Max Norm

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$



# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:

# Max Norm

$$\|x\|_n = \left(\sum_{i=1}^n |x_i|^n\right)^{1/n}$$
$$\|x - y\|_1$$

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

- 1.  $d(x, y) = 0$  if and only if  $x = y$
- 2.  $d(x, y) = d(y, x)$
- 3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:

Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:      Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:      Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

$$2. |x - y| = |-(x - y)| = |y - x|$$
$$\therefore \max |x - y| = \max |y - x|$$

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:      Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

$$2. |x - y| = |-(x - y)| = |y - x|$$
$$\therefore \max |x - y| = \max |y - x|$$

$$3. \max |x - z| = \max |x - y + y - z|$$

Assumption:  $|S| < \infty$

$$S = \{1, 2, 3\}$$

$\mathbb{R}^{|S|}$  = space  
of 3D vectors

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:      Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

$$2. |x - y| = |-(x - y)| = |y - x|$$
$$\therefore \max |x - y| = \max |y - x|$$

$$3. \max |x - z| = \max |x - y + y - z|$$
$$\leq \max(|x - y| + |y - z|)$$

# Max Norm

Lemma 1:  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$  is a metric space.

Definition: Let  $M$  be a set. A *metric* on  $M$  is a function  $d : M \times M \rightarrow [0, \infty)$  which satisfies the following three conditions for all  $x, y, z \in M$ :

1.  $d(x, y) = 0$  if and only if  $x = y$
2.  $d(x, y) = d(y, x)$
3.  $d(x, y) \leq d(x, z) + d(z, y)$

Proof:      Note:  $\|x - y\|_\infty = \max_i |x_i - y_i|$

$$1. \max |x - y| = 0 \text{ iff } x_i = y_i \quad \forall i$$

$$2. |x - y| = |-(x - y)| = |y - x|$$
$$\therefore \max |x - y| = \max |y - x|$$

$$3. \max |x - z| = \max |x - y + y - z|$$
$$\leq \max(|x - y| + |y - z|)$$
$$\leq \max |x - y| + \max |y - z|$$

# Bellman Operator Contraction



# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\|B[V_1] - B[V_2]\|_\infty = \max_{s \in S} |B[V_1](s) - B[V_2](s)|$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}\|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\ &= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right|\end{aligned}$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}\|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\ &= \max_{s \in S} \left| \max_{\substack{a \in A \\ \text{blue wavy}}} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{\substack{a \in A \\ \text{blue wavy}}} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\ &\leq \max_{s \in S} \left| \max_{\substack{a \in A \\ \text{blue wavy}}} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right|\end{aligned}$$

# Bellman Operator Contraction

$\begin{bmatrix} 3 \\ -4 \end{bmatrix}$

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned} \|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\ &= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\ &\leq \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\ &\qquad\qquad\qquad |\max(x)| \leq \max |x| \end{aligned}$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}
 \|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\
 &= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\
 &\leq \max_{s \in S} \left| \max_{a \in A} \left( \cancel{R(s, a)} + \gamma \sum_{s' \in S} \underline{T(s'|s, a)} V_1(s') - \cancel{R(s, a)} - \gamma \sum_{s' \in S} \underline{T(s'|s, a)} V_2(s') \right) \right| \\
 &\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} \underline{T(s'|s, a)} (V_1(s') - V_2(s')) \right| \qquad \underline{|\max(x)| \leq \max |x|}
 \end{aligned}$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}\|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\&= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} T(s'|s, a) (V_1(s') - V_2(s')) \right| \quad \left| \max(x) \right| \leq \max |x| \\&\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) \underline{|V_1(s') - V_2(s')|}\end{aligned}$$



# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}\|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\&= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} T(s'|s, a) (V_1(s') - V_2(s')) \right| \quad \left| \max(x) \right| \leq \max |x| \\&\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) |V_1(s') - V_2(s')| \\&\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) \|V_1 - V_2\|_\infty\end{aligned}$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}\|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\&= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\&\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} T(s'|s, a) (V_1(s') - V_2(s')) \right| \quad \left| \max(x) \right| \leq \max |x| \\&\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) |V_1(s') - V_2(s')| \\&\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) \underbrace{\|V_1 - V_2\|_\infty} \\&= \gamma \|V_1 - V_2\|_\infty \max_{s \in S, a \in A} \sum_{s' \in S} T(s'|s, a)\end{aligned}$$

# Bellman Operator Contraction

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Proof

$$\begin{aligned}
 \|B[V_1] - B[V_2]\|_\infty &= \max_{s \in S} |B[V_1](s) - B[V_2](s)| \\
 &= \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') \right) - \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\
 &\leq \max_{s \in S} \left| \max_{a \in A} \left( R(s, a) + \gamma \sum_{s' \in S} T(s'|s, a) V_1(s') - R(s, a) - \gamma \sum_{s' \in S} T(s'|s, a) V_2(s') \right) \right| \\
 &\leq \max_{s \in S, a \in A} \left| \gamma \sum_{s' \in S} T(s'|s, a) (V_1(s') - V_2(s')) \right| \quad |\max(x)| \leq \max |x| \\
 &\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) |V_1(s') - V_2(s')| \\
 &\leq \max_{s \in S, a \in A} \gamma \sum_{s' \in S} T(s'|s, a) \|V_1 - V_2\|_\infty \\
 &= \gamma \|V_1 - V_2\|_\infty \max_{s \in S, a \in A} \sum_{s' \in S} T(s'|s, a) \\
 &= \gamma \|V_1 - V_2\|_\infty
 \end{aligned}$$

# Value Iteration Convergence

# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Theorem (Banach): If  $f$  is a contraction mapping on metric space  $(M, d)$ , then

1.  $f$  has a single, unique fixed point  $x^*$ .
2. If  $\{x_k\}$  is a sequence defined by  $x_{k+1} = f(x_k)$ , then  $\lim_{k \rightarrow \infty} x_k = x^*$ .

# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

Proof:

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Theorem (Banach): If  $f$  is a contraction mapping on metric space  $(M, d)$ , then

1.  $f$  has a single, unique fixed point  $x^*$ .
2. If  $\{x_k\}$  is a sequence defined by  $x_{k+1} = f(x_k)$ , then  $\lim_{k \rightarrow \infty} x_k = x^*$ .



# Value Iteration Convergence

Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

Proof:


Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Theorem (Banach): If  $f$  is a contraction mapping on metric space  $(M, d)$ , then

1.  $f$  has a single, unique fixed point  $x^*$ .
2. If  $\{x_k\}$  is a sequence defined by  $x_{k+1} = f(x_k)$ , then  $\lim_{k \rightarrow \infty} x_k = x^*$ .

By Lemma 2 and Banach's theorem (part 2), repeated application of the Bellman operator always has a fixed point limit,  $\hat{V}$ .

# Value Iteration Convergence



Theorem 1: Let  $\{V_1, \dots, V_\infty\}$  be a sequence of value functions for a discrete MDP generated by the recurrence  $V_{k+1} = B[V_k]$ . If  $\gamma < 1$ , then  $\lim_{k \rightarrow \infty} V_k = V^*$ .

Proof:

Lemma 2:  $B$  is a  $\gamma$  contraction mapping on  $(\mathbb{R}^{|S|}, \|\cdot\|_\infty)$ .

Theorem (Banach): If  $f$  is a contraction mapping on metric space  $(M, d)$ , then

1.  $f$  has a single, unique fixed point  $x^*$ .
2. If  $\{x_k\}$  is a sequence defined by  $x_{k+1} = f(x_k)$ , then  $\lim_{k \rightarrow \infty} x_k = x^*$ .

By Lemma 2 and Banach's theorem (part 2), repeated application of the Bellman operator always has a fixed point limit,  $\hat{V}$ .

By Banach's theorem (part 1),  $\hat{V} = B[\hat{V}]$ . Since  $\hat{V}$  satisfies Bellman's equation, it is optimal and  $\hat{V} = \underline{V^*}$ .

# Does Policy Iteration Converge?

# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

1. The policy will either improve or stay the same at each iteration

# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

1. The policy will either improve or stay the same at each iteration
2. The policy will stay the same if and only if  $V^\pi = V^*$

# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

1. The policy will either improve or stay the same at each iteration
2. The policy will stay the same if and only if  $V^\pi = V^*$
3. There are a finite number of possible policies



# Does Policy Iteration Converge?

Theorem: Policy iteration converges to an optimal policy for a finite MDP in finite time.

Proof (sketch):

1. The policy will either improve or stay the same at each iteration
2. The policy will stay the same if and only if  $V^\pi = V^*$
3. There are a finite number of possible policies
4. By (1), (2), and (3), the policy will improve until it finds the optimal policy, and it will always find the optimal policy.

# Guiding Questions

# Guiding Questions

- Does value iteration always converge?
- Is the value function unique?

$$S = \{1, 0.8, \dots, 0\}$$

## Conservation MDP

$$S = \{0, 1\} \quad \text{extinct}$$

$$A = \{p, n\}$$

$$R(s, a) = R(s) + R(a)$$

$$R(1) = 1$$

$$R(0) = -1$$

$$R(p) = -1$$

$$R(n) = 1$$

$$T^p = \begin{bmatrix} 0 & 1 \\ 1 & 0.1 \end{bmatrix}$$

$$T^n = \begin{bmatrix} 1 & 0 \\ 0.5 & 0.5 \end{bmatrix}$$

$$\gamma = 0.9$$

## Break

Find

$$V^*, \pi^*$$

$$V^*(0) = \max_a (R(s, a) + \gamma E[V^*(s')])$$

$$\max_a (R(s, a) + 0.9 V^*(0))$$

$$V^*(0) = 0 + 0.9 V^*(0)$$

$$(1 - 0.9) V^*(0) = 0$$

$$V^*(0) = 0$$

$$\pi^*(0) = n$$

Two options:  $\pi_p(1) = p$   $\pi_n(1) = n$

$$V^{\pi_p}(1) = R(1, p) + \gamma (0.1 V^{\pi_p}(0) + 0.9 V^{\pi_p}(1))$$

$$0 + \gamma (0.9 V^{\pi_p}(1))$$

$$(1 - 0.81) V^{\pi_p}(1) = 0$$

$$V^{\pi_p}(1) = 0$$

$$V^{\pi_n}(1) = R(1, n) + \gamma (0.5 V^{\pi_n}(0) + 0.5 V^{\pi_n}(1))$$

$$2 + \gamma (0.5 V^{\pi_n}(1))$$

$$(1 - 0.45) V^{\pi_n}(1) = 2$$

$$V^{\pi_n}(1) = 3.64 = V^*(1)$$

$$\pi^*(1) = n$$

Bellman's  
Optimality  
Equation

Policy  
Eval,  
Bellman's  
Exp. Eqn.