

Guiding Questions:

Guiding Questions:

1. How do we **encode relationships** between random variables?

Guiding Questions:

1. How do we **encode relationships** between random variables?
2. How do we **infer** something about one random variable given the value of another related one?

Plausibility and Probability

Full story: <https://projecteuclid.org/journals/statistical-science/volume-1/issue-3/The-Axioms-of-Subjective-Probability/10.1214/ss/1177013611.full>

slide not on exam

What is a Random Variable?

R.V. X

Vocabulary/Notation

Vocabulary/Notation

Term	Definition	Coinflip Example	Uniform Example
------	------------	------------------	-----------------

Vocabulary/Notation

Bernoulli(0.5)

Term

Definition

Coinflip Example

Uniform Example

Vocabulary/Notation

		Bernoulli(0.5)	$\mathcal{U}(0, 1)$
Term	Definition	Coinflip Example	Uniform Example

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X)			

Vocabulary/Notation

Term	Definition	Bernoulli(0.5)	$\mathcal{U}(0, 1)$
Coinflip Example	Uniform Example		
support(X)	All the values that X can take		

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$	All the values that X can take		

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take		

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution			

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$

Distribution

- Discrete: PMF
- Continuous: PDF

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution <ul style="list-style-type: none">• Discrete: PMF• Continuous: PDF	Maps each value in the support to a real number indicating its probability		

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$
Distribution <ul style="list-style-type: none">• Discrete: PMF• Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$	


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution • Discrete: PMF • Continuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>X</th><th>P(X)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	X	P(X)	0	0.5	1	0.5	
X	P(X)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table border="1"> <thead> <tr> <th>X</th> <th>P(X)</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </tbody> </table>	X	P(X)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$
X	P(X)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none">Discrete: PMFContinuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ 
x	P(x)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none">Discrete: PMFContinuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) =$ 
x	P(x)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none">Discrete: PMFContinuous: PDF	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table><tr><th>x</th><th>P(x)</th></tr><tr><td>0</td><td>0.5</td></tr><tr><td>1</td><td>0.5</td></tr></table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ 
x	P(x)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$ 						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <table data-bbox="1332 948 1574 1128"> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table>	x	P(x)	0	0.5	1	0.5	$p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation									


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation	Single representative value of the random variable, "mean"								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	Single representative value of the random variable, "mean"								


Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	Single representative value of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$							

Vocabulary/Notation


Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	Single representative value of the random variable, "mean"	$E[X] = \sum_{x \in X} x P(x)$ $= 0.5$							

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x)dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	Single representative value of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	$E[X] = \int_{x \in X} xp(x)dx$						

4/25

Vocabulary/Notation

Term	Definition	Bernoulli(0.5) Coinflip Example	$\mathcal{U}(0, 1)$ Uniform Example						
support(X) $x \in X$ $X \in [0, 1]$	All the values that X can take	$\{h, t\}$ or $\{0, 1\}$	$[0, 1]$						
Distribution <ul style="list-style-type: none"> Discrete: PMF Continuous: PDF 	Maps each value in the support to a real number indicating its probability	$P(X = 1) = 0.5$ $P(X = 0) = 0.5$ $P(X)$ is a table <div> <table> <tr> <th>x</th> <th>P(x)</th> </tr> <tr> <td>0</td> <td>0.5</td> </tr> <tr> <td>1</td> <td>0.5</td> </tr> </table> </div>	x	P(x)	0	0.5	1	0.5	 $p(x) = \begin{cases} 1 & \text{if } x \in [0, 1] \\ 0 & \text{o.w.} \end{cases}$ $P(X = 1) = 0$ $P(X \in [a, b]) = \int_a^b p(x) dx$
x	P(x)								
0	0.5								
1	0.5								
Expectation $E[X]$	Single representative value of the random variable, "mean"	$E[X] = \sum_{x \in X} xP(x)$ $= 0.5$	$E[X] = \int_{x \in X} xp(x) dx$ $= 0.5$						

4.28

Distributions of related R.V.s

Distributions of related R.V.s

Joint Distribution

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Distributions of related R.V.s

Joint Distribution

Conditional Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X \mid Y, Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	$P(X Y=1, Z=1)$
0	0.84
1	0.16

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	$P(X Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X \mid Y, Z)$$

(Distribution - valued function)

X	$P(X \mid Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

X	Y	Z	$P(X, Y, Z)$
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

Conditional Distribution

$$P(X | Y, Z)$$

(Distribution - valued function)

X	$P(X Y=1, Z=1)$
0	0.84
1	0.16

Marginal Distribution

$$P(X) \quad P(Y) \quad P(Z)$$

X	$P(X)$	Y	$P(Y)$
0	0.85	0	0.45
1	0.15	1	0.55

Z	$P(Z)$
0	0.20
1	0.80

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

(Burrito-level)

(Filet Minion Level: Axioms of Probability)

AXIOM 1. STRUCTURE OF UNKNOWN REAL NUMBERS AND PLAUSIBLE VALUE. We assume a set T of unknown numbers is a partially ordered commutative algebra over \mathbb{R} with identity, 1.

We assume in addition a given sub-Boolean algebra E of $E(T)$ with $0, 1 \in E$ and denote by E_0 the set of non-zero members of E . We assume that the partial ordering in $E(T)$ as a Boolean algebra coincides with the ordering that $E(T)$ inherits from the algebra T . Finally, we assume a function $PV : T \times E_0 \rightarrow \mathbb{R}$, called **PLAUSIBLE VALUE**, whose value on the pair (x, e) is denoted $PV(x|e)$.

not on exam

AXIOM 2. STRONG RESCALING FOR PLAUSIBLE VALUE. If a, b belong to \mathbb{R} , if x belongs to T , and if e belongs to E_0 , then

$$PV(ax + b|e) = aPV(x|e) + b. \quad (2)$$

AXIOM 3. ORDER CONSISTENCY FOR PLAUSIBLE VALUE. If $x, y \in T$ and if $e \in E_0$, implies that $x \leq y$, then $PV(x|e) \leq PV(y|e)$.

Notice that if $e \in E(T)$, then $0 \leq e \leq 1$, in T , as it is true in the lattice ordering of $E(T)$.

AXIOM 4. THE COX AXIOM FOR PLAUSIBLE VALUE: If e, c are fixed in E , with $ec \in E_0$, if x_1, x_2 are in T , if $PV(x_1|ec) = PV(x_2|ec)$, then $PV(x_1|c) = PV(x_2|c)$. That is, we assume that as a function of x , the plausible value $PV(x|c)$ depends only on $PV(x|ec)$.

AXIOM 5. RESTRICTED ADDITIVITY OF PLAUSIBLE VALUE. For each fixed $y \in T$ and $e \in E_0$, the plausible value $PV(x + y|e)$ as a function of $x \in T$ depends only on $PV(x|e)$, which is to say that if $x_1, x_2 \in T$ and $PV(x_1|e) = PV(x_2|e)$, then $PV(x_1 + y|e) = PV(x_2 + y|e)$.

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

1)

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

1) a) $0 \leq P(X \mid Y) \leq 1$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

Joint \rightarrow Marginal

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) P(Y) P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Joint + Marginal \rightarrow Conditional

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules (Burrito-level)

- 1) a) $0 \leq P(X \mid Y) \leq 1$
b) $\sum_{x \in X} P(x \mid Y) = 1$

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Joint + Marginal \rightarrow Conditional

Marginal + Conditional \rightarrow Joint

$$P(X, Y) = P(X \mid Y) P(Y)$$

Distributions of related R.V.s

Joint Distribution

$$P(X, Y, Z)$$

Conditional Distribution

$$P(X \mid Y, Z)$$

Marginal Distribution

$$P(X) \ P(Y) \ P(Z)$$

3 Rules

- 1) a) $0 \leq P(X \mid Y) \leq 1$
 b) $\sum_{x \in X} P(x \mid Y) = 1$

X	Y	Z	P(X, Y, Z)
0	0	0	0.08
0	0	1	0.31
0	1	0	0.09
0	1	1	0.37
1	0	0	0.01
1	0	1	0.05
1	1	0	0.02
1	1	1	0.07

- 2) "Law of total probability"

$$P(X) = \sum_{y \in Y} P(X, y)$$

- 3) Definition of Conditional Probability

$$P(X \mid Y) = \frac{P(X, Y)}{P(Y)}$$

Joint \rightarrow Marginal

Joint + Marginal \rightarrow Conditional

Marginal + Conditional \rightarrow Joint

$$P(X, Y) = P(X \mid Y) P(Y)$$

Break

1) a) $0 \leq P(X | Y) \leq 1$

b) $\sum_{x \in X} P(x | Y) = 1$

2) $P(X) = \sum_{y \in Y} P(X, y)$

3) $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X|Y) P(Y)$$

- $P \in \{0, 1\}$: Powder Day
 - $C \in \{0, 1\}$: Pass Clear
 - 1 in 5 days is a powder day
 - The pass is clear 8 in 10 days
 - If it is a powder day, there is a 50% chance the pass is blocked
-
- What is the probability that there is a powder day and the pass is clear?
 - What is the probability that the pass is blocked on a non-powder day

Bayes Rule

- Know: $P(B \mid A), P(A), P(B)$
- Want: $P(A \mid B)$

Independence

Independence

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

Independence

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

Independence

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Independence

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definition: X and Y are *conditionally independent* given Z iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

Independence

Definition: X and Y are *independent* iff $P(X, Y) = P(X) P(Y)$

$$X \perp Y$$

$$P(X|Y) = P(X)$$

Definition: X and Y are *conditionally independent* given Z iff

$$P(X, Y | Z) = P(X | Z) P(Y | Z)$$

$$X \perp Y | Z$$

Rules for Continuous RVs

Discrete

1) a) $0 \leq P(X | Y) \leq 1$

b) $\sum_{x \in X} P(x | Y) = 1$

2) $P(X) = \sum_{y \in Y} P(X, y)$

3) $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

Continuous

1)

Rules for Continuous RVs

Discrete

1) a) $0 \leq P(X | Y) \leq 1$
b) $\sum_{x \in X} P(x | Y) = 1$

2) $P(X) = \sum_{y \in Y} P(X, y)$

3) $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

Continuous

1) $0 \leq p(X | Y)$

Rules for Continuous RVs

Discrete

$$1) \text{ a) } 0 \leq P(X | Y) \leq 1$$
$$\text{b) } \sum_{x \in X} P(x | Y) = 1$$

$$2) P(X) = \sum_{y \in Y} P(X, y)$$

$$3) P(X | Y) = \frac{P(X, Y)}{P(Y)}$$

$$P(X, Y) = P(X | Y) P(Y)$$

Continuous

$$1) \quad 0 \leq p(X | Y)$$
$$\int_X p(x|Y) dx = 1$$

Rules for Continuous RVs

Discrete

1) a) $0 \leq P(X | Y) \leq 1$

b) $\sum_{x \in X} P(x | Y) = 1$

2) $P(X) = \sum_{y \in Y} P(X, y)$

3) $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

Continuous

1) $0 \leq p(X | Y)$

$$\int_X p(x|Y) dx = 1$$

2)

$$p(X) = \int_Y p(X, y) dy$$

Rules for Continuous RVs

Discrete

1) a) $0 \leq P(X | Y) \leq 1$

b) $\sum_{x \in X} P(x | Y) = 1$

2) $P(X) = \sum_{y \in Y} P(X, y)$

3) $P(X | Y) = \frac{P(X, Y)}{P(Y)}$

$$P(X, Y) = P(X | Y) P(Y)$$

Continuous

1) $0 \leq p(X | Y)$

$$\int_X p(x|Y) dx = 1$$

2)

$$p(X) = \int_Y p(X, y) dy$$

3) $p(X | Y) = \frac{p(X, Y)}{p(Y)}$

$$p(X, Y) = p(X | Y) p(Y)$$

Multivariate Gaussian Distribution

Joint Distribution

Conditional Distribution

Marginal Distribution

Guiding Questions:

Guiding Questions:

1. How do we **encode relationships** between random variables?

Guiding Questions:

1. How do we **encode relationships** between random variables?
2. How do we **infer** something about one random variable given the value of another related one?