Combining IQ Scores

Analysis and a corrected statistical treatment

31 October 2026

The covariance identities reproduced in Ronald K. Hoeflin's article are correct. Their application to qualifying IQ scores requires additional assumptions about the reference population, the composite's distribution, and the meaning of the score. This report distinguishes those assumptions from the algebra, corrects the numerical table, and supplies an independently written replacement article. All displayed numerical examples have been recalculated. A technical appendix illustrates why a standardized composite and an estimate of latent ability need not coincide.

1 Analysis of the original article

Hoeflin reproduces George A. Ferguson's treatment of variances and covariances and applies it to combinations of IQ scores. The basic identity

$$\operatorname{Var}(X+Y) = \operatorname{Var}(X) + \operatorname{Var}(Y) + 2\operatorname{Cov}(X,Y) \tag{1}$$

is correct. It determines the variance of a sum without requiring normality. The principal weaknesses arise when a standardized sum is assigned an extreme population percentile and used as a qualifying score.

- **Population calibration.** Correlations among the article's 27 selected high scorers do not establish the correlations in the population used to standardize the tests. Selection can change the relationships, and the appropriate correction depends on how selection occurred (Sackett and Yang, 2000).
- **Distributional assumptions.** Mean zero and variance one do not establish a normal distribution. A one-in-a-million upper-tail claim requires additional justification for the tail model.
- **Numerical correction.** The 100-test entry should be about 3.69, rather than 3.71, standard deviations. The more accurate normal quantile also changes some final digits elsewhere; for example, the four-test result rounds to 3.98 rather than 3.97.
- **Score interpretation.** A standardized composite, a latent ability estimate, and a validated IQ score are distinct quantities. Their equivalence requires evidence about the battery and its intended use (American Educational Research Association et al., 2014, Standard 1.14).
- **Threshold precision.** Rounded whole-number scores are useful illustrations, but are not exact qualifying cutoffs. The scale's standard deviation and the classification rule must be specified.

The replacement below preserves the valid calculation while making its assumptions explicit. The article is newly written; it is not a transcription or a reproduction of the original scans.

2 Corrected article: a statistical method and its limits

Several cognitive test scores can be combined into a single score. The calculation must account for the tests' correlations: correlated scores cannot be treated as independent when calculating the composite's variance. The resulting composite also needs an explicit interpretation. Rescaling it to an IQ-style scale supplies units; evidence about the battery establishes what those units mean.

Before calculating a composite, specify the tests, their weights, the administration procedure, and the population against which scores will be compared. Composite construction and interpretation require their own justification; evidence supporting the individual tests does not automatically establish the meaning of their combination (American Educational Research Association et al., 2014, Standard 1.14).

2.1 Put the component scores on comparable scales

For test i, convert the observed score x_i to a standardized score:

$$z_i = \frac{x_i - \mu_i}{\sigma_i}, \tag{2}$$

where μ_i and σ_i are its mean and standard deviation in the intended reference population. The norms must support comparison with that population. Using different numbers labelled "IQ" does not establish their comparability.

For example, a score of 160 on a scale with mean 100 and standard deviation 15 corresponds to $z = 4$. The same numerical score on a scale with standard deviation 16 corresponds to $z = 3.75$. These scores should be standardized before they are combined.

2.2 Standardize the weighted composite

Let R be the population correlation matrix of the standardized test scores, and let w contain fixed weights. The weighted sum has variance

$$\mathrm{Var}(w^\mathsf{T} Z) = w^\mathsf{T} R w. \tag{3}$$

Consequently, provided this variance is positive, a standardized composite is

$$\boxed{Z_C = \frac{w^\mathsf{T} z}{\sqrt{w^\mathsf{T} R w}}.} \tag{4}$$

With the correct population parameters, this composite has mean zero and variance one. This statement requires finite variances, but does not require normality. When the parameters are estimated, their uncertainty affects the calibration of the composite.

For an equally weighted battery of $n \geq 2$ tests, write $\bar{z}$ for the average standardized score and $\bar{r}$ for the average correlation across the distinct pairs of tests:

$$\bar{r} = \frac{2}{n(n-1)} \sum_{i<j} r_{ij}. \tag{5}$$

Then

$$\begin{aligned}\mathrm{Var}(\bar{Z}) &= \frac{n + 2\sum_{i<j} r_{ij}}{n^2} \\ &= \frac{1 + (n-1)\bar{r}}{n} = \bar{r} + \frac{1-\bar{r}}{n}.\end{aligned} \tag{6}$$

Therefore,

$$\boxed{Z_C = \frac{\bar{z}}{\sqrt{\bar{r} + (1-\bar{r})/n}}.} \tag{7}$$

For a single test, $Z_C = z_1$; no inter-test correlation is needed.

Identical pairwise correlations are unnecessary for equation (6); their average is sufficient for this particular equal-weight composite. The full matrix remains relevant to other weighting schemes and to understanding the battery.

An IQ-style reporting scale with mean 100 and standard deviation s can be defined by

$$I_C = 100 + sZ_C. \tag{8}$$

When all component scores already use mean 100 and the same standard deviation s, the formula becomes

$$I_C = 100 + \frac{\bar{I} - 100}{\sqrt{\bar{r} + (1-\bar{r})/n}}, \tag{9}$$

where $\bar{I}$ is their arithmetic average.

For example, three scores averaging 160, each on an SD-16 scale, produce $\bar{z} = 3.75$. If their population average correlation is 0.60, then

$$Z_C = \frac{3.75}{\sqrt{0.60 + 0.40/3}} \approx 4.3791, \qquad I_C \approx 170.06. \tag{10}$$

This is a score on the specified composite. Interpreting it as a measure of general intelligence requires evidence about what the composite measures.

2.3 Separate standardization from percentile estimation

To assign a percentile, we also need the composite's distribution. Mean zero and variance one do not make a distribution normal. Joint multivariate normality of the component scores is a sufficient condition for a fixed linear composite to be normal. Normal distributions for the individual tests alone are insufficient.

If a standard-normal distribution is justified for the composite, its percentile is $100\Phi(Z_C)$, where Φ is the standard-normal cumulative distribution function. A one-in-a-million upper-tail probability corresponds to

$$q = \Phi^{-1}(0.999999) \approx 4.7534243088. \tag{11}$$

Thus the model-based 99.9999th-percentile threshold is

$$\boxed{\bar{z}_{\min} = q\sqrt{\bar{r} + \frac{1-\bar{r}}{n}}.} \tag{12}$$

The quantile can be reproduced in R with `qnorm(1e-6, lower.tail = FALSE)`; the official documentation defines the normal quantile function and its upper-tail option (R Core Team).

For $\bar{r} = 0.60$, the thresholds are shown in table 1.

Table 1: Model-based 99.9999th-percentile thresholds, assuming a normal composite and average pairwise correlation 0.60.

Number of tests	**Required average component z**	**Required average component score, mean 100 and SD 16**
1	4.7534	176.05
2	4.2516	168.03
3	4.0706	165.13
4	3.9770	163.63
5	3.9198	162.72
10	3.8027	160.84
100	3.6942	159.11
1,000	3.6832	158.93
Limit as n increases	3.6820	158.91

These are thresholds for the *average component score*, under the stated model. The target composite score remains approximately 176.05 on the SD-16 scale in every row. Values are rounded for display; an operational classification rule must specify rounding and use the appropriate attainable scores.

For an SD-15 reporting scale, use $100 + 15\bar{z}_{\min}$ instead. In particular, the single-test threshold would be approximately 171.30. An IQ number without its scale convention is ambiguous.

2.4 Interpret the limiting threshold and correlation sensitivity

The limiting average threshold follows from

$$\lim_{n\to\infty} \bar{z}_{\min} = q\sqrt{\bar{r}}, \tag{13}$$

if the average correlation remains fixed as tests are added. With $\bar{r} = 0.60$, the limiting average component score on the SD-16 scale is approximately 158.91. This is a feature of the assumed sequence of batteries and its threshold; it is not a ceiling on possible composite scores. Adding tests can also change the average correlation, so the fixed-correlation limit should not be treated as a universal rule.

The choice of correlation matters substantially. For three tests, an average correlation of 0.90 raises the required average SD-16 component score to approximately 173.48, compared with 165.13 at 0.60. A smaller assumed correlation gives a larger composite score for the same positive average performance. The correlation therefore needs to be estimated for the relevant population and administration procedure. A lower correlation does not, by itself, establish better measurement; it can also reflect different constructs or greater measurement noise.

2.5 Calibrate the population and the scoring procedure

Correlations in a selected high-scoring sample need not equal correlations in the population used to standardize the tests. Selection can change score variances and relationships. The appropriate treatment depends on the selection mechanism and the information available about the unrestricted population; there

is no universal adjustment. Sackett and Yang (2000) explain why range-restriction corrections depend on those details. A small selected sample cannot directly establish a one-in-a-million population frequency.

The scoring procedure also matters. A fixed battery, the best few results from many attempts, and a rule that keeps adding tests until someone qualifies have different score distributions. Each procedure requires calibration for the rule actually used. Repeated administrations can introduce practice effects, as documented by Hausknecht et al. (2007). Repeated or overlapping tests should not be counted as fresh independent observations without accounting for their dependence and administration history.

Finally, variability across people is different from uncertainty about one person. The correlation matrix determines how the composite varies in the reference population. It does not, by itself, provide an individual confidence interval, separate general ability from shared methods, or establish measurement precision at extreme scores. Those questions require an appropriate measurement model and supporting data.

A defensible report should identify the battery, norms, weights, estimated correlations, and uncertainty. Direct composite norms should be used when they adequately support the intended interpretation. If an extreme percentile comes from a distributional model beyond the available normative evidence, report it as a model-based extrapolation and show how sensitive it is to the assumptions. The testing standards require clear descriptions of reference populations and norming procedures in Standards 5.8–5.9, and justification for score interpretations in Standards 1.1–1.4 (American Educational Research Association et al., 2014).

A Composite scores and latent ability: a mathematical illustration

To see why these quantities differ, suppose, expressly as a hypothetical model, that

$$Z_i = \sqrt{\rho}\, G + \sqrt{1-\rho}\, \varepsilon_i, \qquad 0 < \rho < 1, \tag{14}$$

where G and all ε_i are independent standard-normal variables. This model treats all shared variance as one common factor and all residuals as mutually independent. Real batteries need not satisfy those assumptions.

In this particular model, the equal-weight composite's squared correlation with G is

$$\mathcal{R} = \mathrm{Corr}(G, Z_C)^2 = \frac{n\rho}{1 + (n-1)\rho}. \tag{15}$$

The conditional distribution of the latent factor has

$$\mathrm{E}[G \mid Z_1, \ldots, Z_n] = \sqrt{\mathcal{R}}\, Z_C, \tag{16}$$

$$\mathrm{Var}(G \mid Z_1, \ldots, Z_n) = \frac{1-\rho}{1 + (n-1)\rho}. \tag{17}$$

These identities follow from Gaussian conditioning. In particular, $\mathrm{Cov}(G, \bar{Z}) = \sqrt{\rho}$ and $\mathrm{Var}(\bar{Z}) = \rho + (1-\rho)/n$; the residual contrasts contain no additional information about G under this model.

For three tests with $\rho = 0.60$, a composite at $Z_C = 4.7534$ has conditional mean G of approximately 4.30 and conditional standard deviation approximately 0.43. On a hypothetical latent scale with mean 100 and SD 16, these become a mean of approximately 168.79 and a standard deviation of 6.82. The composite's reporting-scale value is approximately 176.05. The two values answer different questions, even in this unusually simple model.

The illustration is a mathematical counterexample to treating a standardized composite as automatically identical to a latent-ability estimate. It is not a proposed correction factor for actual IQ scores, and inter-test correlation should not generally be equated with reliability or common-factor variance.

B Numerical reproduction

The accompanying script `scripts/verify_calculations.py` recalculates the normal quantile, checks the table against `data/thresholds.csv` and the numerical entries in this source, and verifies the worked examples. It uses the upper-tail probability directly through `scipy.stats.norm.isf(1e-6)`. The mathematical results do not depend on the choice of software.

C Final software verification and its limits

The final checks were executed on `computer10` on 31 October 2026. The following results refer to the source and evidence preserved at repository commit `2636a42`. The Deal verification report and the SPARK integration report link the exact commands, tool versions, source hashes, checksums, and raw results. Repository-wide Python counts below cover both analyses and their shared tooling; they are not counts for this article alone.

C.1 Deal contracts and Python checks

The contract-coverage gate found 20 Python files, 142 named functions, and 416 Deal decorators, with zero coverage gaps. All 19 positive files, including tests and the gate itself, passed static lint with zero diagnostics, empty output, and exit status 0, both normally and under `python -O`. The deliberately incorrect fixture was checked separately: it produced exactly the expected `DEL012` postcondition diagnostic and exit status 1. Its failure is a required negative control, not an unresolved production finding.

All 14 earlier production diagnostics were resolved. Bounded quotient helpers retain the original division operations, numerical tolerances remain unchanged, and the fixes do not disable contracts or suppress diagnostics. The runtime, property, and generated-test suite passed 110 tests with no failures or skips. All 27 recorded command checks had their expected outcomes, including numerical scripts, Ada-output validators, replay, optimized-Python execution, and evidence audits.

Deal Solver proved the two pure proof-acceptance gate postconditions; the deliberately wrong symbolic contract was rejected. This focused symbolic result does not prove the complete Python programs. A clean linter result is also not a universal proof: input-dependent contracts remain runtime checks, and finite tests cannot establish correctness for every possible input. The recorded environment used Python 3.13.5, Deal 4.24.6, and Deal Solver 0.1.2.

C.2 SPARK results for the Hoeflin/Ferguson core

The fresh five-prover portfolio completed all 47 native checks for this analysis: 8 flow checks and 39 proof checks, with zero unproved and zero justified checks. CVC5, Z3, Alt-Ergo, classic COLIBRI, and Yices were requested together. This is a portfolio result; it does not mean that each prover independently discharged every obligation. Incomplete individual runs remain identified in the earlier evidence. The deliberately incorrect inclusion-exclusion control was rejected with an unproved functional contract.

The proof project selects only `hoeflin_core.ads` and `hoeflin_core.adb`. Under their stated input preconditions, the contracts establish exact integer variance bookkeeping, bounded floating-point transforms, and nested event counts with their specified increments. The core uses no project assumptions, imported proof axioms, or suppressed checks. The native Hoeflin proof report records the totals. The toolchain manifest records the compiler banner as GNAT 13.2.0 and the GNATprove banner as `0.0w`; these distinct identifiers should not be conflated.

C.3 Numerical, simulation, and model boundaries

The Ada drivers and `normal_support` are outside SPARK. Their elementary functions, C `erfc`, and inverse-tail bisection are checked numerically against SciPy, not certified by the core proof. Likewise, a proof of a floating-point bound does not establish an approximation-error bound. The proof-friendly evaluation orders for thresholds and composites are algebraically equivalent to the report's formulas, but are not claimed to give bit-identical floating-point results.

The retained simulation uses 1,000,000 trials for each of eight test-count cases, with seed `20261030` reset across cases. These common random numbers make the cases dependent. Independent Python replay reproduced the tail counts and checked the means and variances within the recorded tolerances, normally and under optimized Python. This establishes reproducibility and numerical agreement, not independence of the pseudo-random draws. Counts beyond the one-in-a-million cutoff remain descriptive; this sample size does not validate that extreme tail.

Neither Deal nor SPARK proves the Gaussian common-factor model, the population covariance calibration, an identification of correlation with reliability, or the empirical validity of a psychometric interpretation. Those assumptions and the distinction between a standardized composite and latent ability remain as discussed above.

From a full Git clone, install the pinned development dependencies and run `make check` at the repository root. The Python auditor `verification/verify_evidence.py` checks the retained hashes, native totals, and negative-control result. Historical Python identities are resolved through verified Git blobs; changed Ada sources cannot use that fallback. An evidence audit verifies record integrity and source identity; it does not rerun the provers or independently validate their SMT proofs.

C.4 Final consistency audit

A final audit of revision `9a244d8` on 31 October 2026 found no discrepancies in the corrected numerical results or verification claims. Fresh Ada calculation and simulation CSVs matched the retained files byte for byte; validation and replay passed normally and under `python -O`. All 110 tests and the audits of all three SPARK campaigns passed. The provers were not rerun. The audit record is retained under

`verification/evidence/final-consistency-computer10-20261031/`.

References

American Educational Research Association, American Psychological Association, and National Council on Measurement in Education. *Standards for Educational and Psychological Testing*. American Educational Research Association, Washington, DC, 2014. ISBN 978-0-935302-35-6. URL https://www.testingstandards.net/uploads/7/6/6/4/76643089/9780935302356.pdf.

John P. Hausknecht, Jane A. Halpert, Nicole T. Di Paolo, and Meghan O. Moriarty Gerrard. Retesting in selection: A meta-analysis of coaching and practice effects for tests of cognitive ability. *Journal of Applied Psychology*, 92(2):373–385, 2007. doi: 10.1037/0021-9010.92.2.373. URL https://ecommons.cornell.edu/entities/publication/97becf46-3c0a-4de4-a48f-79b715ba8446.

Ronald K. Hoeflin. The statistical technique for combining IQ scores. Noesis, issue 141; reprinted from The Circle. URL https://megasociety.org/noesis/141/ferguson.html. Includes an excerpt from George A. Ferguson, Statistical Analysis in Psychology and Education, third edition, pp. 103–105. Accessed 30 October 2026.

R Core Team. *The Normal Distribution*. R Foundation for Statistical Computing. URL https://stat.ethz.ch/R-manual/R-devel/library/stats/html/Normal.html. R stats package reference documentation. Accessed 30 October 2026.

Paul R. Sackett and Hyuck Yang. Correction for range restriction: An expanded typology. *Journal of Applied Psychology*, 85(1):112–118, 2000. doi: 10.1037/0021-9010.85.1.112. URL https://pubmed.ncbi.nlm.nih.gov/10740961/.