

NP-Complete graph problems - KombSøg 2017

Anders H Pedersen

May 2017

The graph problems and reductions below were introduced by Thomas Dueholm Hansen, at the lecture on NP-Complete graph problems may 3 2017.

1 Graph problems

1.1 INDEPENDENT SET

Given: Graph $G = (V, E)$, target K .

Question: Does there exist $I \subseteq V$ such that $|I| \geq K$ and for all $u, v \in I$ we have $(u, v) \notin E$?

1.2 CLIQUE

Given: Graph $G = (V, E)$, target K .

Question: Does there exist $C \subseteq V$ such that $|C| \geq K$ and for all $u, v \in C$ we have $(u, v) \in E$?

1.3 VERTEX COVER

Given: Graph $G = (V, E)$, budget B .

Question: Does there exist $C \subseteq V$ such that $|C| \leq B$ and for all $(u, v) \in E$ we have $(u \in C \vee v \in C)$?

1.4 MAX CUT

Given: Graph $G = (V, E)$, target K .

Question: Does there exist a cut $(S, V \setminus S)$ of size at least K ?

1.5 MAX BISECTION

Given: Graph $G = (V, E)$, target K .

Question: Does there exist a cut $(S, V \setminus S)$ of size at least K , such that $|S| = |V \setminus S|$?

1.6 BISECTION WIDTH

Given: Graph $G = (V, E)$, target K .

Question: Does there exist a cut $(S, V \setminus S)$ of size at most K , such that $|S| = |V \setminus S|$?

1.7 HAMILTONIAN PATH

Given: Graph $G = (V, E)$.

Question: Does G have a path that visits every vertex exactly once?

1.8 TSP

Given: Distance matrix D , target t .

Question: Is there a tour of length at most t that visits every node in the graph defined by D exactly once?

2 Reductions

As always, in a reduction $L_1 \leq L_2$, we describe a polynomial time computable function r , such that $\forall x : x \in L_1 \iff r(x) \in L_2$. We then argue that it is indeed polynomial, and that both directions of the bi-implication holds.

2.1 3SAT \leq INDEPENDENT SET

We prove **Theorem 9.4: INDEPENDENT SET is NP-Complete**, by reducing from 3SAT, which we know is NP-Complete.

For this reduction we need a gadget, the triangle. The logic behind this is that, if a graph contains a triangle, then at most one of the nodes can be in the independent set. We restrict the class of graphs we consider, to graphs whose nodes can be partitioned in m disjoint triangles. This ensures that an independent set can contain at most m nodes.

For each of the m clauses in our input CNF formula, we create a triangle where the nodes are labeled with the literals of the clause. Next we add an edge between two nodes in different triangles if and only if the nodes correspond to opposite literals. For example an edge between x_1 and $\neg x_1$. Adding these edges between opposite literals, ensures that we cannot pick both (it would not be an independent set). The reduction is completed by setting the independent set goal $k = m$.

Example:

Given a 3CNF formula: $f = (x_1 \vee x_2 \vee x_3) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_3) \wedge (\neg x_1 \vee x_2 \vee x_3)$
We construct the graph shown in figure 1.




Figure 1: Graph $r(f)$ corresponding to the formula f

Analysis

For each clause, we just create a triangle. Then we add at most $3 * (m - 1)$ edges between the triangles - so the reduction is clearly polynomial.

\Rightarrow :

Given a satisfying assignment for the CNF formula f from above, we must show that the graph, $r(f)$, has an independent set of size $K=m$. The assignment: $x_1 = 0, x_2 = 1, x_3 = 0$ satisfies the formula, and for our independent set of size K , we pick a node in each triangle which make the corresponding clause true. For the truth assignment we chose above, this means x_2 from triangle 1, $\neg x_1$ in triangle 2, $\neg x_1$ in triangle 3. (Note that this is not the only independent set of size K).

\Leftarrow :

Read more in papadimitriou page 188-190.

2.2 INDEPENDENT SET \leq CLIQUE

2.3 INDEPENDENT SET \leq VERTEX COVER

2.4 NAESAT \leq MAX CUT

2.5 3SAT \leq HAMILTONIAN PATH

2.6 HAMILTONIAN PATH \leq TSP