

# Kinetic energy

Ayoub Gouasmi

March 13th

Let's start with the 1D compressible Euler equations. We have:

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x}(\rho u) = 0, \quad (1)$$

$$\frac{\partial \rho u}{\partial t} + \frac{\partial}{\partial x}(\rho u^2 + p) = 0, \quad (2)$$

$$\frac{\partial \rho e^t}{\partial t} + \frac{\partial}{\partial x}(u(\rho e^t + p)) = 0. \quad (3)$$

where  $e^t := e + k$ ,  $k := \frac{u^2}{2}$ . We assume a calorically perfect gas, so pressure and internal energy are related through:

$$p := (\gamma - 1)\rho e.$$

Let's express the time derivative of  $\rho k$  (the quantity of interest) in terms of the time derivatives of mass, momentum and total energy (the quantities whose equations we have). Using the chain-rule, we obtain:

$$\frac{\partial(\rho k)}{\partial t} = \frac{1}{2} \frac{\partial}{\partial t} \left( \frac{(\rho u)^2}{\rho} \right) = u \frac{\partial}{\partial t}(\rho u) - \frac{1}{2} u^2 \frac{\partial \rho}{\partial t}.$$

We can substitute the time derivatives on the right-hand side with spatial derivatives through equations (1) and (2). This gives:

$$\begin{aligned} \frac{\partial(\rho k)}{\partial t} &= -u \frac{\partial}{\partial x}(\rho u^2 + p) + \frac{1}{2} u^2 \frac{\partial}{\partial x}(\rho u) \\ &= -u \frac{\partial}{\partial x}(\rho u^2) + \frac{1}{2} u^2 \frac{\partial}{\partial x}(\rho u) - u \frac{\partial p}{\partial x} \\ &= -u^2 \frac{\partial}{\partial x}(\rho u) - \rho u^2 \frac{\partial u}{\partial x} + \frac{1}{2} u^2 \frac{\partial}{\partial x}(\rho u) - u \frac{\partial p}{\partial x} \\ &= -\frac{1}{2} u^2 \frac{\partial}{\partial x}(\rho u) - \rho u^2 \frac{\partial u}{\partial x} - u \frac{\partial p}{\partial x} \\ &= -\frac{\partial}{\partial x} \left( \frac{1}{2} u^2 (\rho u) \right) - u \frac{\partial p}{\partial x} \\ &= -\frac{\partial}{\partial x} \left( \frac{1}{2} u^2 (\rho u) + p u \right) + p \frac{\partial u}{\partial x} \\ &= -\frac{\partial}{\partial x} (u(\rho k + p)) + p \frac{\partial u}{\partial x} \end{aligned}$$

This gives the final result:

$$\frac{\partial(\rho k)}{\partial t} + \frac{\partial}{\partial x} (u(\rho k + p)) = p \frac{\partial u}{\partial x}. \quad (4)$$

This equation does not express conservation unless the right-hand side term of equation (4) is zero. This is the case for incompressible flow since for such flows:

$$\frac{\partial u}{\partial x} = 0.$$

Let's do it in 2D now. We have:

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial x}(\rho u) + \frac{\partial}{\partial y}(\rho v) = 0, \quad (5)$$

$$\frac{\partial \rho u}{\partial t} + \frac{\partial}{\partial x}(\rho u^2 + p) + \frac{\partial}{\partial y}(\rho uv) = 0, \quad (6)$$

$$\frac{\partial \rho v}{\partial t} + \frac{\partial}{\partial x}(\rho uv) + \frac{\partial}{\partial y}(\rho v^2 + p) = 0, \quad (7)$$

$$\frac{\partial \rho e^t}{\partial t} + \frac{\partial}{\partial x}(u(\rho e^t + p)) + \frac{\partial}{\partial y}(v(\rho e^t + p)) = 0. \quad (8)$$

where  $e^t := e + k$ ,  $k := \frac{u^2 + v^2}{2}$ . Just as before, express the time derivative of  $\rho k$  (the quantity of interest) in terms of the time derivatives of mass, momentum and total energy (the quantities whose equations we have). Using the chain-rule, we obtain:

$$\frac{\partial(\rho k)}{\partial t} = \frac{1}{2} \frac{\partial}{\partial t} \left( \frac{(\rho u)^2}{\rho} + \frac{(\rho v)^2}{\rho} \right) = u \frac{\partial}{\partial t}(\rho u) + v \frac{\partial}{\partial t}(\rho v) - \frac{1}{2}(u^2 + v^2) \frac{\partial \rho}{\partial t}.$$

We can substitute the time derivatives on the right-hand side with spatial derivatives through equations (5), (6) and (7).

**Question 1:** This gives:

$$\begin{aligned} \frac{\partial(\rho k)}{\partial t} &= \text{Figure} \\ &= \text{it} \\ &= \text{out} \\ &= \text{with} \\ &= \text{chain} \\ &= \text{rules} \end{aligned}$$

This gives the final result:

$$\frac{\partial(\rho k)}{\partial t} + \frac{\partial}{\partial x}(u(\rho k + p)) + \frac{\partial}{\partial y}(v(\rho k + p)) = p \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right). \quad (9)$$

For incompressible flows the right-hand side term is zero (divergence free).

**Question 2:** Find a problem that illustrates that kinetic energy is not conserved for compressible flows. Let's stick to the 1D system (1) - (3) and set periodic boundary conditions. Run it with a decent numerical scheme on a decently fine grid (so that your numerical solution can be trusted) and show a time history of:

$$\int_{\Omega} (\rho k) dV$$

**Question 3:** Consider the two-dimensional compressible Navier-Stokes equation (just add a viscous diffusion term). Prove that viscous diffusion dissipates kinetic energy.

## References