

SYDE 556/750

Simulating Neurobiological Systems
Lecture 2: Neurons


Andreas Stöckel

January 9 & 14, 2020




UNIVERSITY OF
WATERLOO



FACULTY OF
ENGINEERING



Textbook Neuron and Cell Membrane




Injecting a Current Into a Detailed Neuron Model




Computer simulation of an Hodgkin-Huxley type neuron with Traub kinematics (Roger D. Traub and Richard Miles, *Neuronal Networks of the Hippocampus*, Cambridge University Press, 1991)

The Leaky Integrate-and-Fire Equivalent Circuit




Injecting a Current Ramp into a Detailed Neuron Model





Computer simulation of an Hodgkin-Huxley type neuron with Traub kinematics (Roger D. Traub and Richard Miles, *Neuronal Networks of the Hippocampus*, Cambridge University Press, 1991)


Injecting a Current Ramp into a LIF Neuron Model




Exploring the LIF Rate Approximation



Limitations of the LIF Neuron Model



Artificial Rate Neurons: LIF




$$G[J] = \frac{1}{\tau_{\text{ref}} - \tau_{\text{RC}} \log \left(1 - \frac{1}{J}\right)}$$

Usefulness to neurobiological systems modellers:

- ⊕ Biologically motivated
- ⊕ Captures saturation effects
- Relatively slow to evaluate numerically (for machine-learning people)
- ⊖ Spike onset is smooth in noisy systems

Artificial Rate Neurons: ReLU




$$G[J] = \max\{0, J\}$$

Usefulness to neurobiological systems modellers:

- ⊕ Fast to evaluate
- Rough approximation of the LIF response curve
- ⊖ Does not capture saturation effects
- ⊖ Spike onset is smooth in noisy systems

Artificial Rate Neurons: Smooth ReLU (Softplus)




$$G[J] = \log(1 + \exp(J))$$

Usefulness to neurobiological systems modellers:

- ⊕ Models smooth spike onset
- Rough approximation of the LIF response curve
- ⊖ Does not capture saturation effects

Artificial Rate Neurons: Logistic Function




$$G[J] = \frac{1}{1 + e^{-J}}$$

Usefulness to neurobiological systems modellers:

- Models smooth spike onset and saturation (?)

Artificial Rate Neurons: Hyperbolic Tangent



$$G[J] = \tanh(J) = \frac{e^J - e^{-J}}{e^J + e^{-J}}$$

Usefulness to neurobiological systems modellers:

- Models smooth spike onset and saturation (?)
- Negative rates

Image sources

Title slide

Image of rat primary cortical neurons in culture.

Author: ZEISS Microscopy, <http://www.zeiss.com/celldiscoverer>.

From Wikimedia.