

PMATH 370 Winter 2024:

Lecture Notes

1	Iteration and Orbits	3
1.1	Orbits	3
1.2	Real analysis review	5
1.3	Orbits, revisited	8
2	Graphical Analysis	12
2.1	Cobweb plots	12
3	Fixed Points	15
3.1	Attracting/repelling fixed point theorems	15
3.2	Neutral fixed points	18
4	Bifurcations	22
5	Cantor set	25
6	Symbolic dynamics	28
6.1	Intro to topology	29
6.2	Revisiting the itinerary	31
7	Chaos	34
7.1	Prerequisites to chaos	34
7.2	Defining chaos	36
8	Sarkovskii's Theorem	38
9	Fractals	41
	Back Matter	44
	List of Named Results	44
	Index of Defined Terms	45

Lecture notes taken, unless otherwise specified, by myself during the Winter 2024 offering of PMATH 370, taught by Blake Madill.

Lectures

			Lecture 9	Jan 26	25
			Lecture 10	Jan 29	28
Lecture 1	Jan 8	3	Lecture 11	Jan 31	30
Lecture 2	Jan 10	5	Lecture 12	Feb 2	32
Lecture 3	Jan 12	7	Lecture 13	Feb 5	34
Lecture 4	Jan 15	10	Lecture 14	Feb 7	36
Lecture 5	Jan 17	13	Lecture 15	Feb 9	38
Lecture 6	Jan 19	18	Lecture 16	Feb 12	39
Lecture 7	Jan 22	20	Lecture 17	Feb 14	41
Lecture 8	Jan 24	23			

Chapter 1

Iteration and Orbits

1.1 Orbits

Definition 1.1.1 (iteration)

Let $f : A \rightarrow \mathbb{R}$ such that $A \subseteq \mathbb{R}$ and $f(A) \subseteq A$. For $a \in A$ we may iterate the function at a :

$$x_1 = a, x_2 = f(a), x_3 = \underbrace{f(f(a))}_{f^2(a)}, \dots, x_i = f^{i-1}(a), \dots$$

The sequence $(x_n)_{n=1}^\infty$ is the orbit of a under f (abbreviated (x_n) without limits).

Lecture 1
Jan 8

Example 1.1.2. Let $f(x) = x^4 + 2x^2 - 2$, $a = -1$. What is the orbit of a under f ?

Solution. $a = -1$, $f(a) = 1$, $f(f(a)) = f(1) = 1$, so we have $-1, 1, 1, 1, \dots$. We call this eventually constant. \square

Example 1.1.3. Let $f(x) = -x^2 - x + 1$, $a = 0$. What is the orbit of a under f ?

Solution. Calculate: $0, 1, -1, 1, -1, 1, \dots$. We call this eventually periodic (with period 2). \square

Example 1.1.4. Let $f(x) = x^3 - 3x + 1$, $a = 1$. What is the orbit of a under f ?

Solution. Calculate the first few terms: $1, -1, 3, 19, \dots$ (too big). This is a divergence to infinity. \square

Example 1.1.5. Let $f(x) = x^2 + 2x$, $a = -0.5$. What is the orbit of a under f ?

Solution. Calculate: $-0.5, -0.75, -0.9375, -0.9961 \dots$ and we make an educated guess that this converges to -1 since $f(-1) = -1$, a fixed point. \square

Example 1.1.6. Let $f(x) = x^3 - 3x$, $a = 0.75$. What is the orbit of a under f ?

Solution. Calculate: $0.75, -1.828, -0.625, 1.631, -0.552, \dots$. There is no clear pattern, so we call this chaotic. In fact, the orbit is dense in a neighbourhood of 0. \square

We can start to formalize the examples.

Definition 1.1.7 (fixed point)

Let $f : A \rightarrow \mathbb{R}$ such that $f(A) \subseteq A$. A point $a \in A$ is fixed if $f(a) = a$.

Then, the orbit of a under f is (a, a, a, \dots) which is constant.

Example 1.1.8. Find all fixed points of $f(x) = x^2 + x - 4$.


Solution. We find points where $f(x) = x$, i.e., $x^2 + x - 4 = x$.

$$x^2 + x - 4 = x \iff x^2 = 4 \iff x = \pm 2$$

\square

Example 1.1.9. How many fixed points does $f(x) = 2 \sin x$ have?

Solution. Consider where the curve $y = 2 \sin x$ meets $y = x$:



We can see there are three fixed points. \square

Example 1.1.10. Prove that $f(x) = x^4 - 3x + 1$ has a fixed point.

Proof. We must show there is a solution to $x^4 - 3x + 1 \iff x^4 - 4x + 1 = 0$. Let $g(x) = x^4 - 4x + 1$. Since $g(x)$ is continuous, $g(0) = 1 > 0$, and $g(1) = -2 < 0$, by the Intermediate Value Theorem, there must exist a root of g on the interval $(0, 1)$. That is, a fixed point of f . \square

Definition 1.1.11 (periodicity)

Let $f : A \rightarrow \mathbb{R}, f(A) \subseteq A$.

1. A point $a \in A$ is periodic for f if its orbit is periodic. An orbit is periodic if for some $n \in \mathbb{N}$, $f^n(a) = a$. The smallest n is the period of (the orbit of) a .
2. An orbit (of a point) is eventually periodic if there exists $n < m$ such that $f^n(a) = f^m(a)$. The smallest difference $m - n$ is the period of the orbit.

Definition 1.1.12 (doubling function)

$D : [0, 1) \rightarrow [0, 1) : x \mapsto 2x - \lfloor 2x \rfloor$ returns the fractional part of $2x$.

Lecture 2
Jan 10

Example 1.1.13. $D(0.4) = 0.8$, $D(0.6) = 0.2$, $D(0.8) = 0.6$, $D(0.5) = 0$.

This is a nice function that gives lots of periodic orbits for funsies.

Example 1.1.14. Find the orbit of $a = \frac{1}{5}$ under D .

Solution. Double until we pass 1: $\frac{1}{5}, \frac{2}{5}, \frac{4}{5}, \frac{8}{5} \rightarrow \frac{3}{5}, \frac{6}{5} \rightarrow \frac{1}{5}$. The period is $|\{\frac{1}{5}, \frac{2}{5}, \frac{4}{5}, \frac{3}{5}\}| = 4$. □

Example 1.1.15. Find the orbit of $a = \frac{1}{20}$ under D .

Solution. Double: $\frac{1}{20}, \frac{1}{10}, \frac{1}{5}$ and we can stop because ex. 1.1.14 showed $\frac{1}{5}$ is periodic.

So this is eventually periodic with period 4. □

Problem 1.1.16

Given f and a , does $f^n(a)$ tend towards some limit L ?

To solve this problem, we need to rigorously define “tend” and “limit”.

1.2 Real analysis review

Notation. If $(x_n)_{n=1}^\infty$ is a sequence of real numbers, we write $(x_n) \subseteq \mathbb{R}$.

Definition 1.2.1 (convergence of a sequence)

Let $(x_n) \subseteq \mathbb{R}$, $x \in \mathbb{R}$.

We say (x_n) converges to x if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $|x_n - x| < \varepsilon$ for all $n \geq N$.

Then, we write $x_n \rightarrow x$ or $\lim x_n = x$.

Example 1.2.2. Show that $\frac{1}{n} \rightarrow 0$.

Proof. Let $\varepsilon > 0$. Consider $N = \frac{2}{\varepsilon} > \frac{1}{\varepsilon}$. For $n \geq N$, we have

$$\left| \frac{1}{n} - 0 \right| = \frac{1}{n} < \varepsilon$$

Therefore, $\frac{1}{n} \rightarrow 0$. □

Example 1.2.3. Prove that $\frac{2n}{n+3} \rightarrow 2$.

Proof. Let $\varepsilon > 0$. Since we know $\frac{1}{n} \rightarrow 0$, let $N \in \mathbb{N}$ such that $\frac{1}{N} < \frac{\varepsilon}{6}$.

For $n \geq N$,

$$\left| \frac{2n}{n+3} - 2 \right| = \left| \frac{2n}{n+3} - \frac{2n+6}{n+3} \right| = \left| \frac{-6}{n+3} \right| = \frac{6}{n+3} < \frac{6}{n} \leq \frac{6}{N} < 6 \cdot \frac{\varepsilon}{6} = \varepsilon$$

Therefore, $\frac{2n}{n+3} \rightarrow 2$. □

Definition 1.2.4 (bounded sequence)

A sequence (x_n) is bounded (by M) if there exists $M > 0$ such that $\forall n \in \mathbb{N}$, $|x_n| \leq M$.

Proposition 1.2.5 (convergence implies boundedness)

If (x_n) is convergent, then (x_n) is bounded.

Proof. Suppose $x_n \rightarrow x$. Then, there exists $N \in \mathbb{N}$ such that if $n \geq N$, then $|x_n - x| < 1$.

For $n \geq N$, $|x_n| - |x| \leq |x_n - x| < 1$. That is, $|x_n| < 1 + |x|$.

Let $M = \max\{|x_1|, \dots, |x_{N-1}|, 1 + |x|\}$. Then, for both all $n < N$ and $n \geq N$, we have $|x_n| \leq M$. □

Remark 1.2.6. The converse is not true. Notice that $x_n = (-1)^n$ is bounded by 1 but obviously not convergent.

Proposition 1.2.7 (limit laws)

Let $x_n \rightarrow x$ and $y_n \rightarrow y$. Then:

- (1) $x_n + y_n \rightarrow x + y$
- (2) $x_n y_n \rightarrow xy$

Proof. (1) Let $\varepsilon > 0$. Then, since $x_n \rightarrow x$ and $y_n \rightarrow y$, there exist $N_1, N_2 \in \mathbb{N}$ such that $n \geq N_1 \implies |x_n - x| < \frac{\varepsilon}{2}$ and $n \geq N_2 \implies |y_n - y| < \frac{\varepsilon}{2}$.

For $N = \max\{N_1, N_2\}$ and $n \geq N$,

$$\begin{aligned} |(x_n + y_n) - (x + y)| &= |(x_n - x) + (y_n - y)| \\ &\leq |x_n - x| + |y_n - y| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

That is, $x_n + y_n \rightarrow x + y$.

(2) Let $\varepsilon > 0$. Notice that:

$$|x_n y_n - xy| = |x_n y_n - x_n y + x_n y - xy| \leq |x_n| \cdot |y_n - y| + |y| \cdot |x_n - x| \quad (*)$$

Since x_n is bounded, there exists $M > 0$ such that $|x_n| \leq M$ for all n .

Let $N_1, N_2 \in \mathbb{N}$ such that

$$\begin{aligned} n \geq N_1 &\implies |x_n - x| \leq \frac{\varepsilon}{2(|y| + 1)} \text{ and} \\ n \geq N_2 &\implies |y_n - y| < \frac{\varepsilon}{2M}. \end{aligned}$$

Then, for $n \geq N := \max\{N_1, N_2\}$, $|x_n y_n - xy| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$ by (*). □

Definition 1.2.8 (Cauchy sequence)

We say $(x_n) \in \mathbb{R}$ is Cauchy if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all n and m ,

$$n, m \geq N \implies |x_n - x_m| < \varepsilon$$

Proposition 1.2.9

Every convergent sequence is Cauchy.

Proof. Intuitively: if the terms get arbitrarily close to some limit, they must get arbitrarily close to each other.

Formally: Let $x_n \rightarrow x$ be a convergent sequence and $\varepsilon > 0$. Since x_n converges, there exists $N \in \mathbb{N}$ such that $n \geq N \implies |x_n - x| < \frac{\varepsilon}{2}$.

Lecture 3
Jan 12

Then, when $n, m \geq N$, we have:

$$\begin{aligned}
 |x_n - x_m| &= |x_n - x_m + x - x| \\
 &= |(x_n - x) + (x - x_m)| \\
 &\leq |x_n - x| + |x_m - x| \\
 &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\
 &= \varepsilon
 \end{aligned}$$

as desired. □

We take the following theorem from real analysis without proof.

Theorem 1.2.10 (completeness of \mathbb{R})

A sequence is Cauchy if and only if it is convergent.

The big idea here: To prove (x_n) is Cauchy, you do not have to guess the limit first. That is, if you must prove convergence but do not care about the limit's value, prove that it is Cauchy instead.

Definition 1.2.11 (continuity of a function)

Let $f : A \rightarrow \mathbb{R}$, $A \subseteq \mathbb{R}$, $a \in A$. We say f is continuous at a if for all $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(a)| < \varepsilon$ whenever $x \in A$ and $|x - a| < \delta$.

If f is continuous at all $a \in A$, we say it is continuous.

We also take this theorem from MATH 137 without proof.

Theorem 1.2.12

A function $f : A \rightarrow \mathbb{R}$ is continuous at $a \in A$ if and only if for all sequences $(x_n) \subseteq A$ with $x_n \rightarrow a$, we have $f(x_n) \rightarrow f(a)$.

1.3 Orbits, revisited

Proposition 1.3.1

If $f : [a, b] \rightarrow [a, b]$ is continuous, then $f(x)$ has a fixed point.

Proof. We know by the domain and codomain that $f(a) \geq a$ and $f(b) \leq b$. This means $f(a) - a \geq 0$ and $f(b) - b \leq 0$. By the IVT on the continuous function $g(x) = f(x) - x$, we know there exists an $x \in [a, b]$ such that $g(x) = f(x) - x = 0 \iff f(x) = x$, i.e., x is a fixed point. □

Definition 1.3.2 (contraction)

Let $f : A \rightarrow \mathbb{R}$, $A \subseteq \mathbb{R}$. We say f is a contraction if there exists $C \in [0, 1)$ such that for all $x, y \in A$,

$$|f(x) - f(y)| \leq C|x - y|$$

This is just a Lipschitz function with Lipschitz constant less than 1.

Proposition 1.3.3

Contractions are continuous.

Proof. Let $\varepsilon > 0$. Suppose f is a contraction such that $|f(x) - f(y)| \leq C|x - y|$.

Consider $y \in A$. Let $\delta = \frac{\varepsilon}{C+1}$ and assume that $x \in A$ and $|x - y| < \delta$. But we have:

$$|f(x) - f(y)| \leq C|x - y| \leq C\delta < \varepsilon$$

as desired. □

Definition 1.3.4 (closure of an interval)

We say $A \subseteq \mathbb{R}$ is closed if whenever $(x_n) \subseteq A$ with $x_n \rightarrow x$, then $x \in A$.

Example 1.3.5. $[a, b]$ is closed but $(0, 1]$ is not because $\frac{1}{n} \rightarrow 0 \notin (0, 1]$.

Theorem 1.3.6 (Banach contraction mapping theorem)

Suppose $A \subseteq \mathbb{R}$ is closed and $f : A \rightarrow A$ is a contraction. Then, there exists a unique fixed point $a \in A$ for f .

Moreover, for all $x \in A$, $f^n(x) \rightarrow a$.

Example 1.3.7. Analyze the orbit of $f : [0, 1] \rightarrow [0, 1]$, $f(x) = \frac{1}{3-x}$.

Solution. We can observe that $\frac{1}{3} \leq \frac{1}{3-x} \leq \frac{1}{2}$.

Also, $f'(x) = \frac{1}{(3-x)^2}$. Notice that $\frac{1}{9} \leq |f'(x)| \leq \frac{1}{4}$. So by the mean value theorem, for all $x, y \in [0, 1]$, there exists $c \in (0, 1)$ such that:

$$\begin{aligned} f(x) - f(y) &= f'(c)(x - y) \\ |f(x) - f(y)| &= |f'(c)| \cdot |x - y| \\ &\leq \frac{1}{4}|x - y| \end{aligned}$$

Then, identifying $C = \frac{1}{4}$, f is a contraction. Now,

$$\frac{1}{3-x} = x \iff 1 = 3x - x^2 \iff x^2 - 3x + 1 = 0 \iff x = \frac{3 \pm \sqrt{9-4}}{2} \iff x = \frac{3 - \sqrt{5}}{2}$$

where we pick the negative root because we need $x \in [0, 1]$.

Therefore, by the [Banach contraction mapping theorem](#), for all $x \in [0, 1]$, $f^n(x) \rightarrow \frac{3-\sqrt{5}}{2}$. \square

Definition 1.3.8

A sequence $(a_n) \subseteq \mathbb{R}$ is strongly-Cauchy if there exists $(\varepsilon_n) \subseteq [0, \infty)$ such that $\sum_{n=1}^{\infty} \varepsilon_n < \infty$ and for all n , $|a_n - a_{n+1}| < \varepsilon_n$.

Informally, “far enough along the sequence, the *neighbours* must get close”. This is distinct from Cauchy, which is “far enough along the sequence, the *terms* must get close”.

Remark 1.3.9 (assignment hint!). Let $\sum_{n=1}^{\infty} a_n = L$. This means that $\sum_{k=1}^n a_k \xrightarrow{n \rightarrow \infty} L$.

That is, for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $n \geq N$ implies $|\sum_{k=1}^n a_k - L| < \varepsilon$.

But $|\sum_{k=1}^n a_k - L| = |\sum_{k=1}^{\infty} a_k - \sum_{k=1}^n a_k| = |\sum_{k=n+1}^{\infty} a_k| < \varepsilon$.

We can now prove the [Banach contraction mapping theorem](#).

Proof. Let $A \subseteq \mathbb{R}$ be closed and suppose there exists $f : A \rightarrow A$ and $C \in [0, 1)$ such that $|f(x) - f(y)| \leq C|x - y|$ for all x and y in A .

Fix $x_0 \in A$ and construct the orbit $x_1 = f(x_0)$, $x_2 = f(x_1)$, \dots , $x_n = f(x_{n-1}) = f^n(x_0)$.

For $n \in \mathbb{N}$, since f is a contraction,

$$\begin{aligned} |x_{n+1} - x_n| &= |f(x_n) - f(x_{n-1})| \\ &\leq C|x_n - x_{n-1}| \\ &= C|f(x_{n-1}) - f(x_{n-2})| \\ &\leq C^2|x_{n-1} - x_{n-2}| \\ &\vdots \\ &\leq C^n|x_1 - x_0| \end{aligned}$$

Since $\sum_{n=1}^{\infty} C^n|x_1 - x_0| = |x_1 - x_0| \sum_{n=1}^{\infty} C^n$ is a convergent geometric series, we have that the sequence (x_n) is strongly-Cauchy.

Hence, by Assignment 1, $x_n \rightarrow a$ for some limit point $a \in A$ since A is closed.

Since f is continuous (prop. 1.3.3), we have that $f(x_n) \rightarrow f(a)$. By definition, $f(x_n) = x_{n+1}$, so $x_n \rightarrow f(a)$. But we already know $x_n \rightarrow a$, so $a = f(a)$. That is, a is a fixed point of f .

It remains to show uniqueness.

Lecture 4
Jan 15

Suppose $a, b \in A$ such that $f(a) = a$ and $f(b) = b$.

$$\begin{aligned} |f(a) - f(b)| &\leq C|a - b| \\ |a - b| &\leq C|a - b| \end{aligned}$$

Since $C < 1$, we must have $|a - b| = 0$, that is, $a = b$. □

Chapter 2

Graphical Analysis

2.1 Cobweb plots

Recall ex. 1.1.9. To visualize the orbit of a under f , we can:

1. Superimpose $y = f(x)$ over the line $y = x$.
 2. Connect a vertical line $(a, a) - (a, f(a))$
 3. Connect a horizontal line $(a, f(a)) - (f(a), f(a))$
 4. Connect a vertical line $(f(a), f(a)) - (f(a), f(f(a)))$
 5. Connect a horizontal line $(f(a), f(f(a))) - (f(f(a)), f(f(a)))$
- etc.


This is sometimes called a cobweb plot. We will be using <https://marksmath.org/visualization/cobwebs/> to make cobweb plots.

Within these lecture notes, I use a \LaTeX macro to draw plots [defined here](#).

Example 2.1.1. Conduct a complete orbit analysis of $f(x) = x^2 - x + 1$

Solution. Playing around, we find that there is one fixed point $x = 1$.

When $x \in [0, 1]$, $f^n(x) \rightarrow 1$. Otherwise, $f^n(x) \rightarrow \infty$.




□

↓ Lectures 5 and 6 adapted from *Rosie* ↓


Lecture 5
Jan 17

Example 2.1.2. Conduct a complete orbit analysis of $f(x) = -\frac{3}{2}x^2 + \frac{5}{2}x + 1$.

Solution. At $x = 0$, we can see there is a cycle going from $0 \rightarrow 1 \rightarrow 2 \rightarrow 0$:



At points near 0, like $x = -0.3$ or $x = 0.3$, the graph becomes chaotic:



It appears that the cobweb densely covers the graph. □

As an aside, note that we cannot actually hit every point in the interval because the orbit is countable (i.e., has the same size as the naturals) but the interval is uncountable. We will later show that the points are dense (as the rationals are).


Chapter 3

Fixed Points


3.1 Attracting/repelling fixed point theorems

Remark 3.1.1. If $f(x)$ is continuous and $f^n(a) \rightarrow L$, then $f^{n+1}(a) \rightarrow f(L)$. Therefore, $f(L) = L$ is a fixed point.

Example 3.1.2. The function $f(x) = x^3$ has three fixed points: $0, \pm 1$. For $x \in (-1, 1)$, we see that $f^n(x) \rightarrow 0$:




It looks like point 0 is attracting the orbit. For $x \in (-\infty, -1) \cup (1, \infty)$, we see $f^n(x) \rightarrow \infty$:




so the points ± 1 are repelling the orbit.


Example 3.1.3. The function $f(x) = x^3 - 3x$ also has three fixed points: $0, \pm 2$. To the right (left) of ± 2 , orbits go to infinity:



The point 0 is repelling (in a different sense) since we get chaos:



At $x_0 = \pm 1$, the orbit is eventually constant, jumping to the fixed point ∓ 2 :

**Definition 3.1.4**

Let a be a fixed point of $f(x)$.

1. If $|f'(a)| > 1$, we call a a repelling fixed point
2. If $|f'(a)| < 1$, we call a a attracting fixed point
3. If $|f'(a)| = 1$, we call a a neutral fixed point

Neutral fixed points can be a lot of different things.

Theorem 3.1.5 (attracting fixed point theorem)

Suppose a is an attracting fixed point of $f(x)$. Then, there exists an open interval I containing a such that

1. for all $x \in I$, $n \in \mathbb{N}$, $f^n(x) \in I$
2. for all $x \in I$, $f^n(x) \rightarrow a$

Recall the ε - δ definition of a limit.

Definition 3.1.6 (limit of a function at a point)

Let $f : A \rightarrow \mathbb{R}$, $A \subseteq \mathbb{R}$.

We say a point $a \in A$ is non-isolated if for each $\varepsilon > 0$ there exists $b \in A$, $b \neq a$ with $b \in (a - \varepsilon, a + \varepsilon)$.

Suppose a is non-isolated. We say $\lim_{x \rightarrow a} f(x) = L$ if for all $\varepsilon > 0$, there exists a $\delta > 0$ such that $|f(x) - L| < \varepsilon$ whenever $a \in A$ and $0 < |x - a| < \delta$.

It is important to define non-isolation. If a is isolated, we can choose a δ where $|x - a| < \delta$ is false. Then, every point is vacuously a limit point.

We now give the proof of the [attracting fixed point theorem](#):

Proof. Assume $|f'(a)| < 1$. Then, there exists $c \in \mathbb{R}$ such that $|f'(a)| < c < 1$. By definition of the

derivative, this means we can write

$$\lim_{x \rightarrow a} \frac{|f(x) - f(a)|}{|x - a|} < c$$

and by the definition of the limit, we know there exists $\delta > 0$ such that

$$\frac{|f(x) - f(a)|}{|x - a|} \leq c, \quad \forall x \in (a - \delta, a + \delta)$$

Hence, for $x \in I := (a - \delta, a + \delta)$, we have $|f(x) - f(a)| \leq c|x - a|$ and f is a contraction.

In particular, for $x \in I$, we have

$$\begin{aligned} |f(x) - a| &= |f(x) - f(a)| && (a \text{ is a fixed point}) \\ &\leq c|x - a| \leq |x - a| && (c \in (0, 1)) \\ &< \delta \end{aligned}$$

That is, $f(x) \in (a - \delta, a + \delta) = I$. Continuing for the rest of the orbit, for all $n \in \mathbb{N}$,

$$|f^n(x) - a| \leq c^n|x - a| \leq |x - a| < \delta$$

so we also have $f^n(x) \in I$.

Finally, notice that $0 \leq |f^n(x) - a| \leq c^n|x - a|$ and $c^n|x - a| \rightarrow 0$ since $c \in (0, 1)$. By the squeeze theorem, $|f^n(x) - a| \rightarrow 0$. \square

Theorem 3.1.7 (repelling fixed point theorem)

Suppose a is a repelling fixed point for $f(x)$. Then, there exists an open interval I containing a such that for all $x \in I$, $x \neq a$, there exists $n \in \mathbb{N}$ such that $f^n(x) \notin I$.

Proof. Say $|f'(a)| > c > 1$. Then, as above, there exists a δ such that

$$\lim_{x \rightarrow a} \frac{|f(x) - f(a)|}{|x - a|} > c \implies |f(x) - f(a)| \geq c|x - a|$$


for all $x \in I := (a - \delta, a + \delta)$.

Since a is a fixed point, $|f(x) - f(a)| = |f(x) - a|$. Suppose for a contradiction that for all n , $f^n(x) \in I$. But since $c > 1$, $|f^n(x) - a| \geq c^n|x - a| \rightarrow \infty$. That is, δ must be arbitrarily large, which it is not. \square


3.2 Neutral fixed points

Neutral fixed points can exhibit a lot of different behaviours.


Example 3.2.1. For $f(x) = -x$, 0 is a fixed point with $|f'(0)| = 1$. The orbit bounces:




Example 3.2.2. For $f(x) = x - x^2$, $|f'(1)| = 1$ is a neutral fixed point. It is attracting from the right and repelling from the left:




Example 3.2.3. For $f(x) = x - x^3$, $|f'(0)| = 1$ is a neutral fixed point. It is weakly attracting, attracting but too slowly.



Example 3.2.4. For $f(x) = x + x^3$, $|f'(0)| = 1$ is a neutral fixed point. It is weakly repelling, repelling but too slowly:



Example 3.2.5. Consider $f(x) = x^2 - 1$. The orbit at $a = 0$ is periodic $(0, -1, 0, -1, \dots)$ with period 2. Near 0, the orbit tends to the $(0, -1)$ -cycle:



We will call 0 an attracting periodic point because 0 is an attracting point of $f^2(x)$.

↑ Lectures 5 and 6 adapted from *Rosie* ↑

Definition 3.2.6

Let a be a periodic point for $f(x)$ with period n .

We say a is an attracting/repelling/neutral periodic point if a is an attracting/repelling/neutral fixed point of f^n

Lecture 7
Jan 22

Finding a closed form expression for something like $f^{10}(x)$ is a nightmare, so we need a better way.

Proposition 3.2.7

Let $f(x)$ be a differentiable function. Then, $(f^n)'(x) = f'(x) \cdot f'(f(x)) \cdots f'(f^{n-1}(x))$.

Proof. Proceed by induction on n .

If $n = 1$, we have $f'(x) = f'(x)$ and we are done.

Suppose $(f^n)'(x) = \prod_{k=0}^{n-1} f'(f^k(x))$ for some $n \geq 1$. Consider f^{n+1} :

$$\frac{d}{dx} f^{n+1}(x) = \frac{d}{dx} f(f^n(x)) = f'(f^n(x)) \cdot (f^n)'(x)$$

by the chain rule. Then,

$$\begin{aligned} (f^{n+1})'(x) &= f'(f^n(x)) \cdot (f^n)'(x) \\ &= f'(f^n(x)) \cdot \prod_{k=0}^{n-1} f'(f^k(x)) \\ &= \prod_{k=0}^n f'(f^k(x)) \end{aligned}$$

completing the proof. □

Example 3.2.8. Analyze the periodic point $f(x) = -\frac{3}{2}x^2 + \frac{5}{2}x + 1$, $a = 0$

Solution. The orbit is $(0, 1, 2, 0, 1, 2, \dots)$ with period 3.

We have $f'(x) = -3x + \frac{5}{2}$. Then, $(f^3)'(0) = f'(0)f'(1)f'(2) = (-\frac{7}{2})(-\frac{1}{2})(\frac{5}{2}) = \frac{35}{8} > 1$.

Therefore, the point is repelling. □

Chapter 4

Bifurcations

In general, bifurcation theory is the study of how a family of curves can change when a defining parameter is changed.

Consider the quadratic family:

$$Q_C(x) = x^2 + C$$

defined by the parameter $C \in \mathbb{R}$.

Problem 4.0.1

How does the behaviour (fixed points, orbits, etc.) of Q_C change based on C ?

First, we can find the fixed points (if they exist) by solving

$$Q_C(x) = x \iff x^2 - x + C = 0 \iff x = \frac{1 \pm \sqrt{1 - 4C}}{2}$$

and note that $Q_C(x)$ has 2 fixed points when $C < \frac{1}{4}$, 1 fixed point when $C = \frac{1}{4}$, and no fixed points when $C > \frac{1}{4}$.

Suppose $C > \frac{1}{4}$. Then, we must have $Q_C^n(x) \rightarrow \infty$ for all x .

Instead, if $C = \frac{1}{4}$, $Q_C(x)$ has the unique fixed point $p = \frac{1}{2}$. Since $Q'_C(x) = 2x$ and $Q'_C(p) = 1$, this is a neutral fixed point. In fact, it attracts to one side and repels from the other.

Finally, if $C < \frac{1}{4}$, $Q_C(x)$ has two fixed points $p_+ = \frac{1 + \sqrt{1 - 4C}}{2}$ and $p_- = \frac{1 - \sqrt{1 - 4C}}{2}$. Then, $Q'_C(p_+) =$

$1 + \sqrt{1 - 4C} > 1$ is repelling. Next,

$$\begin{aligned}
 & -1 < Q'_C(p_-) < 1 \\
 \Leftrightarrow & -1 < 1 - \sqrt{1 - 4C} < 1 \\
 \Leftrightarrow & -2 < -\sqrt{1 - 4C} < 0 \\
 \Leftrightarrow & 0 < \sqrt{1 - 4C} < 2 \\
 \Leftrightarrow & -\frac{3}{4} < C < \frac{1}{4}
 \end{aligned}$$

and in fact if $C < -\frac{3}{4}$, $Q'_C(p_-) < -1$ and if $C = -\frac{3}{4}$, $Q'_C(p_-) = -1$.

Theorem 4.0.2

For the family

$$Q_C(x) = x^2 + C,$$

depending on C :

1. All orbits tend to ∞ if $C > \frac{1}{4}$.
2. When $C = \frac{1}{4}$, $Q_C(x)$ has a unique fixed point $\frac{1}{2}$ and it is neutral.
3. If $C < \frac{1}{4}$, $Q_C(x)$ has two fixed points p_+ and p_- . The point p_+ is repelling. Moreover,
 - (a) if $-\frac{3}{4} < C < \frac{1}{4}$, p_- is attracting;
 - (b) if $C = -\frac{3}{4}$, p_- is neutral; and
 - (c) if $C < -\frac{3}{4}$, p_- is repelling.

Definition 4.0.3 (bifurcation)

We say a family of functions $F_\lambda(x)$ undergoes a bifurcation at λ_0 if there is a change in fixed point structure at λ_0 .

Lecture 8
Jan 24

Example 4.0.4. The quadratic family $Q_C(x) = x^2 + C$ undergoes a bifurcation at $\lambda_0 = \frac{1}{4}$.

Definition 4.0.5 (tangent bifurcation)

A family $F_\lambda(x)$ undergoes a tangent bifurcation at λ_0 if there is an open interval I and an $\varepsilon > 0$ such that:

1. for $\lambda_0 - \varepsilon < \lambda < \lambda_0$, $F_\lambda(x)$ has no fixed points on I ;
2. for $\lambda = \lambda_0$, $F_\lambda(x)$ has one fixed point and it is neutral; and
3. for $\lambda_0 < \lambda < \lambda_0 + \varepsilon$, $F_\lambda(x)$ has two fixed points in I , one of which is attracting and the other repelling.

(or with all inequalities flipped)

Visually, you have situations like

TODO: graphs

for $\lambda < \lambda_0$, $\lambda = \lambda_0$, and $\lambda > \lambda_0$.

Example 4.0.6. Consider the exponential family $E_\lambda(x) = e^x + \lambda$ at $\lambda_0 = -1$.

This is a tangent bifurcation.

Example 4.0.7. $F_\lambda(x) = \lambda x(1 - x)$, $\lambda_0 = 1$


Here, we have two fixed points on one side of λ_0 and one fixed point on the other. So this is a bifurcation but not a tangent bifurcation.

Chapter 5

Cantor set

Recall the quadratic family $Q_C(x) = x^2 + C$ for $C < -2$. Then, $p_+ = \frac{1+\sqrt{1-4C}}{2} > 2$ and $-p_+ < -2$. Consider the interval/region $I = [-p_+, p_+]$ and $I \times I$.

Draw the picture of $y = x$, $y = Q_C(x)$, and the box $I \times I$:



Let $J_1 \subseteq I$ be the interval such that $Q_C(x) \notin I$ for all $x \in J_1$.


For $x \in J_1$, $Q_C^n(x) \rightarrow \infty$. Moreover, if there exists n such that $Q_C^n(x) \in J_1$, then $Q_C^n(x) \rightarrow \infty$.

Consider the set of points $\Lambda = \{x \in I : \forall n, Q_C^n(x) \in I\}$ with “interesting” orbits staying inside I .

Now, let $J_2 = \{x \in I : Q_C(x) \in J_1\} = \{x \in I : Q_C^2(x) \notin I\}$ and define higher J_n likewise.

Then, $\Lambda = I \setminus (J_1 \cup J_2 \cup \dots)$ is a Cantor set, that is, a fractal. (roll credits!)

Drawing Λ on the x -axis, we get something that looks like



↓ Lecture 9 adapted from Imaad ↓

Lecture 9
Jan 26

Definition 5.0.1 (Cantor middle thirds set)

Let $C_0 = [0, 1]$. Remove the open middle third interval each time.

That is, $C_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$, $C_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$, and so on.

The set $K = \bigcap_{n=1}^{\infty} C_n$ is the Cantor (middle thirds) set.

Proposition 5.0.2

Suppose a bunch of sets $A_n \subseteq \mathbb{R}$ are closed. Then, $\bigcap A_n$ is also closed.

Proof. Let $(a_k) \subseteq \bigcap A_n$ where $(a_k) \rightarrow a$.

Note that for all n , $(a_k) \subseteq A_n \implies a \in A_n \implies a \in \bigcap A_n$ □

Proposition 5.0.3

Let $A, B \subseteq \mathbb{R}$ be closed. Then, $A \cup B$ is closed.

Proof. Let $(a_n) \subseteq A \cup B$ where $a_n \rightarrow a$.

WLOG, $\{n : a_n \in A\}$ is infinite. This allows us to construct $(b_n) \subseteq A$ such that $b_n \rightarrow a$.

Since A is closed, $a \in A \subseteq A \cup B$. □

Theorem 5.0.4 (Cantor sets are closed)

Any Cantor set, in particular K , is closed.

Theorem 5.0.5

K contains no non-empty open intervals.

Proof. Consider $I \subseteq K$. Then $\forall n, I \subseteq C_n$.

Then $\ell(I) \leq \frac{1}{3^n} \implies \ell(I) = 0 \implies I = \emptyset$, contradiction. □

Now, let's consider the base-3 expansion of $x \in [0, 1]$. $x = 0.s_1s_2s_3, \dots, s_i \in \{0, 1, 2\}$

Consider $\underbrace{[0, 1/3]}_{s_1=0}$ and $\underbrace{[2/3, 1]}_{s_1=2}$ and $\underbrace{[0, 1/9]}_{s_1=0, s_2=0}$ $[2/9, 1/3]$ $[2/3, 7/9]$ $[8/9, 1]$.

Remark 5.0.6. $x \in K$ if and only if x can be written in base 3 using only 0s and 2s

Example 5.0.7. $\frac{1}{3} \in K$. $\frac{1}{3} = 0.1_3 = 0.02222\dots_3$

Theorem 5.0.8

K is uncountable and $|K| = |\mathbb{R}|$.


↑ *Lecture 9 adapted from Imaad* ↑

Symbolic dynamics

Lecture 10
Jan 29

$$\begin{aligned} J_1 &= \{x \in I : Q_C(x) \notin I\} \\ J_2 &= \{x \in I : Q_C(x) \in J_1\} \\ J_3 &= \{x \in I : Q_C(x) \in J_2\} \\ &\vdots \end{aligned}$$

Notation. Define closed intervals $I_0 \cup I_1 := I \setminus J_1$ on the left/right of J_1 :



For $x \in \Lambda$, the itinerary of x is the sequence $S(x) = (x_0 x_1 x_2 x_3 \cdots)$ with $x_i \in \{0, 1\}$ where $x_i = 0 \iff Q_G^i(x) \in I_0$ and $x_i = 1 \iff Q_G^i(x) \in I_1$.

Notation. Let $\Sigma = \{(x_0x_1x_2\cdots) : x_i \in \{0,1\}\}$ be the sequence space. Write elements of Σ as binary strings. Then, $S : \Lambda \rightarrow \Sigma$ is a function.

28

6.1 Intro to topology

Definition 6.1.1 (metric space)

Let X be a set. A function $d : X \times X \rightarrow [0, \infty)$ is a metric if

1. $d(x, y) = 0 \iff x = y$ (positive definiteness),
2. $d(x, y) = d(y, x)$ (symmetry), and
3. $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality).

The pair (X, d) is a metric space.

Once we have a metric space with a notion d of distance, we can adapt all our definitions from real analysis to an abstract space.

Example 6.1.2. The following are all metrics:

- $X = \mathbb{R}$, $d(x, y) = |x - y|$
- $X = \mathbb{R}^n$, $d(\mathbf{x}, \mathbf{y}) = \sqrt{(x_1 - y_1)^2 + \cdots + (x_n - y_n)^2}$
- For any set X , the discrete metric $d(x, y) = [x \neq y]$ (but is not particularly useful).
- For a subset $A \subseteq \mathbb{R}$, $d(x, y) = |x - y|$ is a metric.

Extremely helpfully, we can define a metric on the sequence space.

Definition 6.1.3 (Cantor space)

Let $X = \Sigma$. Define $d(x, y) = \sum_{i=0}^{\infty} 2^{-i} |x_i - y_i|$.

This is well-defined (converges) since $|x_i - y_i| \leq 1$ and $\sum 2^{-i}$ converges.

Example 6.1.4. Let $x = (1111\cdots)$ and $y = (1010\cdots)$. Calculate $d(x, y)$.

Solution. By definition,

$$\begin{aligned}
 d(x, y) &= \sum_{i=0}^{\infty} \frac{x_i - y_i}{2^i} \\
 &= \sum_{i=0}^{\infty} \frac{1}{2^{2i+1}} && \text{(even indices cancel)} \\
 &= \frac{1}{2} \sum_{i=0}^{\infty} \frac{1}{4^i} \\
 &= \frac{1}{2} \left(\frac{1}{1 - \frac{1}{4}} \right) = \frac{1}{2} \left(\frac{4}{3} \right) = \frac{4}{6} = \frac{2}{3}
 \end{aligned}$$

□

We don't want to do this manual calculation every time.

Proposition 6.1.5

Let $x, y \in \Sigma$.

1. If $x_i = y_i$ for $i \leq n$, then $d(x, y) \leq \frac{1}{2^n}$.
2. If $d(x, y) < \frac{1}{2^n}$, then $x_i = y_i$ for $i \leq n$.

Proof. Suppose $x_i = y_i$ for $i \leq n$. Then, $d(x, y) \leq \sum_{k=n+1}^{\infty} \frac{1}{2^k}$ since the first n terms will be 0 and $|x_i - y_i| \leq 1$. That is, $d(x, y) \leq \frac{1/2^{n+1}}{1 - \frac{1}{2}} = \frac{1}{2^n}$.

Conversely, suppose $d(x, y) < \frac{1}{2^n}$ and for a contradiction that there exists $k \leq n$ where $x_k \neq y_k$. Then, there will be a $\frac{1}{2^k}$ term in the sum, so $d(x, y) \geq \frac{1}{2^k} \geq \frac{1}{2^n}$. Contradiction. \square

Example 6.1.6. Let $x = (0000\cdots)$ and $y = (1000\cdots)$. Then, the distance is $\frac{1}{2^0} = 1$. However, $x_0 \neq y_0$.

Definition 6.1.7 (shift map)

The map $\sigma : \Sigma \rightarrow \Sigma : (x_0x_1x_2\cdots) \mapsto (x_1x_2x_3\cdots)$ that shifts a bitstring one bit to the left.

Remark 6.1.8. $\sigma^k(x_0x_1x_2\cdots) = x_kx_{k+1}x_{k+2}\cdots$

Definition 6.1.9 (continuity in metric spaces)

Suppose (X, d) and (Y, d') are (possibly distinct) metric spaces.

A function $f : X \rightarrow Y$ is continuous at $y \in X$ if for all $\varepsilon > 0$, there exists a $\delta > 0$ such that for all $x \in X$,

$$d(x, y) < \delta \implies d'(f(x), f(y)) < \varepsilon$$

We say f is continuous if it is continuous at every $y \in X$

Proposition 6.1.10

The shift map $\sigma : \Sigma \rightarrow \Sigma$ is continuous.

Proof. Fix $y = (y_0y_1y_2\cdots) \in \Sigma$ and let $\varepsilon > 0$. Take $n \in \mathbb{N}$ such that $\frac{1}{2^n} < \varepsilon$.

Consider $\delta = \frac{1}{2^{n+1}}$. Let $x = (x_0x_1x_2\cdots) \in \Sigma$ such that $d(x, y) < \delta$.

Therefore, by prop. 6.1.5, $x_i = y_i$ for $i = 0, 1, \dots, n+1$. Then, $\sigma(x) = (x_1x_2x_3\cdots)$ and $\sigma(y) = (y_1y_2y_3\cdots)$ agree for the first n terms.

Again by prop. 6.1.5, $d(\sigma(x), \sigma(y)) \leq \frac{1}{2^n} < \varepsilon$. \square

Lecture 11
Jan 31

Definition 6.1.11 (convergence in metric spaces)

Let (X, d) be a metric space, $(x_n) \subseteq X$, and $x \in X$.

We say (x_n) converges to x ($x_n \rightarrow x$) if for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$n \geq N \implies d(x_n, x) < \varepsilon.$$

Proposition 6.1.12 (sequential characterization of continuity in metric spaces)

Let (X, d) and (Y, d') be metric spaces and $f : X \rightarrow Y$. Then, f is continuous if and only if $f(x_n) \rightarrow f(x)$ whenever $x_n \rightarrow x$.

Definition 6.1.13 (homeomorphism)

Let (X, d) and (Y, d') be metric spaces. A function $f : X \rightarrow Y$ is a homeomorphism if

1. f is injective,
2. f is surjective,
3. f is continuous, and
4. f^{-1} is continuous.

Suppose $f : X \rightarrow Y$ is a homeomorphism. Then, if $(x_n) \subseteq X$ with $x_n \rightarrow x$, then $f(x_n) \rightarrow f(x)$.

Likewise, suppose $(y_n) \subseteq Y$ with $y_n \rightarrow y$. Say $y_n = f(x_n)$ and $y = f(x)$. Then, $f(x_n) \rightarrow f(x)$, so $f^{-1}(f(x_n)) \rightarrow f^{-1}(f(x))$ and $x_n \rightarrow x$.

That is, f is a *relabelling* of X to Y . We think of X and Y as the “same metric space”.

6.2 Revisiting the itinerary

Remark 6.2.1. Suppose we have $x \in \Lambda$ with $S(x) = (x_0 x_1 \dots)$. Then, by definition, $x \in I_{x_0}$, $Q_c(x) \in I_{x_1}$, $Q_c^2(x) \in I_{x_2}$, etc. Therefore, $S(Q_c(x)) = (x_1 x_2 \dots) = \sigma(S(x))$.

Iterating, $S(Q_c^n(x)) = \sigma^n(x)$.

Theorem 6.2.2

$S : \Lambda \rightarrow \Sigma$ is a homeomorphism.

We will prove this with some more tools. Recall from MATH 137:

Theorem 6.2.3 (monotone convergence theorem)

If $(a_n) \subseteq \mathbb{R}$ is increasing/decreasing and bounded, then (a_n) converges.

Instead of using this directly, we use a lemma:

Lemma 6.2.4 (nested intervals lemma)

If $I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$ are closed intervals, then $\bigcap_{i=1}^{\infty} I_i \neq \emptyset$.

Proof. Let $I_k = [a_k, b_k]$.

That is, $[a_1, b_1] \supseteq [a_2, b_2] \supseteq [a_3, b_3] \dots$.

Then, (a_n) is increasing and $(a_n) \subseteq [a_1, b_1]$. Likewise, (b_n) is decreasing and $(b_n) \subseteq [a_1, b_1]$. By the [monotone convergence theorem](#), $a_n \rightarrow a$ and $b_n \rightarrow b$ for some limit points a and b .

Therefore (handwavey), $\emptyset \neq [a, b] \subseteq \bigcap_{n=1}^{\infty} I_n$. □

We will now prove thm. 6.2.2. It is true for $c < -2$, but we will show it for $c < -\frac{5+2\sqrt{5}}{4}$.

Lecture 12
Feb 2

Proof. (injective) Suppose $x, y \in \Lambda$ with $S(x) = S(y)$ but $x \neq y$. Then, for all n , $Q_c^n(x)$ and $Q_c^n(y)$ live in the same I_0 or I_1 . Recall from Assignment 2 that for all $x \in I \setminus J_1 = I_0 \cup I_1$, we have $|Q'_c(x)| \geq \mu > 1$. By the mean value theorem,

$$|Q_c(x) - Q_c(y)| \geq \mu|x - y|.$$

Since Q_c is injective on I_0 and I_1 , we have that $Q_c(x) \neq Q_c(y)$. Thus,

$$\begin{aligned} |Q_c^2(x) - Q_c^2(y)| &\geq \mu^2|x - y| \\ &\vdots \\ |Q_c^n(x) - Q_c^n(y)| &\geq \mu^n|x - y| \end{aligned}$$

Since $\mu > 1$, we have $\mu^n|x - y| \rightarrow \infty$. However, $|Q_c^n(x) - Q_c^n(y)| \leq \max\{\ell(I_0), \ell(I_1)\}$, so it cannot blow up to infinity. Contradiction, so we have injectivity.

(surjective) Let $y = (y_0 y_1 \dots) \in \Sigma$. For $n \in \mathbb{N}$, define

$$I_{y_0 y_1 \dots y_n} := \{x \in I : x \in I_{y_0}, Q_c(x) \in I_{y_1}, \dots, Q_c^n(x) \in I_{y_n}\}.$$

It is enough to show there exists

$$x \in \bigcap_{n=1}^{\infty} I_{y_0 y_1 \dots y_n}$$

which would imply $S(x) = y$. Clearly, by definition, $I_{y_0} \supseteq I_{y_0 y_1} \supseteq I_{y_0 y_1 y_2} \supseteq \dots$


We claim that each $I_{y_0 y_1 \dots y_n}$ is a closed interval. Proceed by induction.

First, $I_{y_0} \in \{I_0, I_1\}$ so it is closed. Assume $I_{y_0 y_1 \dots y_n}$ is closed for some $n \geq 0$. Note:

$$\begin{aligned} x &\in I_{y_0 y_1 \dots y_{n+1}} \\ \iff x &\in I_{y_0}, Q_c(x) \in I_{y_1}, Q_c(Q_c(x)) \in I_{y_2}, Q_c(Q_c^2(x)) \in I_{y_3}, \dots, Q_c(Q_c^n(x)) \in I_{y_{n+1}} \\ \iff x &\in I_{y_0} \cap Q_c^{-1}(I_{y_1 y_2 \dots y_{n+1}}) \end{aligned} \tag{*}$$

By the inductive hypothesis, $I_{y_1 y_2 \dots y_{n+1}}$ is a closed interval (the subscript has length n).

We have



That is, $Q_c^{-1}(I_{y_1 y_2 \dots y_{n+1}})$ is a union of **two disjoint closed intervals**, one in I_0 and one in I_1 .

In particular, returning to (\star) , $I_{y_0 y_1 \dots y_{n+1}} = I_{y_0} \cap Q_c^{-1}(I_{y_1 y_2 \dots y_{n+1}})$ is one of these closed intervals.

By the **nested intervals lemma**, there must exist $x \in \bigcap_{n=1}^{\infty} I_{y_0 y_1 \dots y_n}$. Hence, $S(x) = y$ and we have surjectivity.

(continuous) Fix $y \in \Lambda$ and say $S(y) = (y_0 y_1 y_2 \dots)$. Let $\varepsilon > 0$ and choose n such that $\frac{1}{2^n} < \varepsilon$.

Consider the 2^{n+1} disjoint, closed intervals $I_{t_0 t_1 \dots t_n}$.

Pick $\delta > 0$ such that $(y - \delta, y + \delta)$ only overlaps with $I_{y_0 y_1 \dots y_n}$. We know δ exists since we have a finite set of disjoint closed intervals.

For $x \in \Lambda$ with $|x - y| < \delta$, $x \in I_{y_0 y_1 \dots y_n}$ and so $d(S(x), S(y)) \leq \frac{1}{2^n} < \varepsilon$.

(continuous inverse) Similar. □

Chapter 7

Chaos

Lecture 13
Feb 5

7.1 Prerequisites to chaos

Definition 7.1.1 (density)

Let (X, d) be a metric space. We say $A \subseteq X$ is dense in X if for all $x \in X$ and $\varepsilon > 0$, there exists $a \in A$ such that $d(a, x) < \varepsilon$.

Informally, there is always something “that close” to any point.

Example 7.1.2. \mathbb{Q} is dense in \mathbb{R} . Given a real number, there is always a decimal approximation with arbitrary accuracy.

\mathbb{Z} is not dense in \mathbb{R} . Given $x = \frac{1}{2} \in \mathbb{R}$, there are no integers within $\varepsilon = \frac{1}{4}$.

Example 7.1.3. Let $A = \{x \in \Sigma : \exists N, \forall i > N, x_i = 0\}$, i.e., the sequences which are eventually constant 0s. This is dense in Σ .

Proof. Let $x = (x_0x_1x_2\cdots) \in \Sigma$ and let $\varepsilon > 0$. As usual, take $n \in \mathbb{N}$ such that $\frac{1}{2^n} < \varepsilon$.

Consider $y = (x_0x_1x_2\cdots x_n0000\cdots) \in A$. Then, by prop. 6.1.5, $d(x, y) \leq \frac{1}{2^n} < \varepsilon$. \square

Exercise 7.1.4. Let $A = \{x \in \Sigma : x \text{ is periodic}\}$. Show that this is dense in Σ .

Remark 7.1.5. A in exercise 7.1.4 is exactly the set of periodic points for the shift map $\sigma : \Sigma \rightarrow \Sigma$.

Proposition 7.1.6

There exists $z \in \Sigma$ such that $\{\sigma^k(z) : k \in \mathbb{N} \cup \{0\}\}$ is dense in Σ .

Proof. Take $z = (0\ 1\ 00\ 01\ 10\ 11\ 000\ 001\ \dots)$ to contain all possible blocks of increasing sizes.

Let $x \in \Sigma$ and $\varepsilon > 0$. Again, let $\frac{1}{2^n} < \varepsilon$.

For some k , $\sigma^k(z)$ and x agree on the first n terms. This must exist because z has *every possible* sequence of n terms. That is, by prop. 6.1.5, $d(\sigma^k(z), x) \leq \frac{1}{2^n} < \varepsilon$. \square

Warning: definition 7.1.7 is not the normal definition from applied math textbooks, but it is what we will use in the course.

Definition 7.1.7 (dynamical system)

A metric space (X, d) together with a continuous function $f : X \rightarrow X$.

This is an abstract space in which we can do orbit analysis and all our fun stuff.

Example 7.1.8. $\sigma : \Sigma \rightarrow \Sigma$ is a dynamical system (see thm. 6.2.2).

Definition 7.1.9 (transitivity)

We say a dynamical system $f : X \rightarrow X$ is transitive if for all $x, y \in X$ and $\varepsilon > 0$, there exists $z \in X$ and $n, m \in \mathbb{N} \cup \{0\}$ such that $d(x, f^n(z)) < \varepsilon$ and $d(y, f^m(z)) < \varepsilon$.

Informally, given any two points, there is a special point whose orbit gets arbitrarily close to both points.

Proposition 7.1.10

$\sigma : \Sigma \rightarrow \Sigma$ is transitive.

Proof. Take z from prop. 7.1.6 such that the orbit is dense in Σ .

Then, for all $\varepsilon > 0$ and $x, y \in \Sigma$, there must exist by the definition of density n and m such that $d(x, \sigma^n(z)) < \varepsilon$ and $d(y, \sigma^m(z)) < \varepsilon$. \square

Definition 7.1.11 (sensitive dependence on initial conditions)

Let $f : X \rightarrow X$ be a dynamical system.

We say f is sensitively dependent on initial conditions (or just sensitive) if

$$\exists \beta > 0, \forall \varepsilon > 0, \forall x \in X, \exists y \in X, \exists k \in \mathbb{N}$$

such that $d(x, y) < \varepsilon$ and $d(f^k(x), f^k(y)) \geq \beta$.

Informally, there exists a “wrongness” β that can always be achieved in the orbit no matter how close two starting points are.

Proposition 7.1.12

$\sigma : \Sigma \rightarrow \Sigma$ is sensitive.

Proof. Take $\beta = 1$.

Let $\varepsilon > 0$ and let $x \in \Sigma$. Say $\frac{1}{2^n} < \varepsilon$ and pick $y \in \Sigma$ such that $0 < d(x, y) < \frac{1}{2^n}$. That is, x and y must agree on the first n terms by prop. 6.1.5, but they are not equal.

Therefore, there exists $k \geq n$ such that $x_k \neq y_k$.

In the distance $d(\sigma^k(x), \sigma^k(y)) \geq \frac{|x_k - y_k|}{2^0} \geq 1 = \beta$. □

7.2 Defining chaos

↓ Lectures 14 and 15 adapted from Imaad ↓

Lecture 14
Feb 7

Definition 7.2.1 (chaos)

A dynamical system $f : X \rightarrow X$ is chaotic if

1. the periodic points for f are dense in X ,
2. f is transitive, and
3. f is sensitive.

Theorem 7.2.2

$\sigma : \Sigma \rightarrow \Sigma$ is chaotic.

Proof. By props. 7.1.6, 7.1.10 and 7.1.12. □

Proposition 7.2.3

Let $(X, d), (Y, d')$ be metric spaces.

Suppose $f : X \rightarrow Y$ is continuous and surjective. If $A \subseteq X$ is dense in X , then $f(A)$ is dense in Y .

Proof. Let $y \in Y$ and say $y = f(x)$.

Let $\epsilon > 0$. Since f is continuous at x , there exists $\delta > 0$ such that

$$d(z, x) < \delta \implies d'(f(z), f(x)) < \epsilon$$

for any z . In particular, since A is dense in X , we may find $a \in A$ such that

$$d(a, x) < \delta \implies d'(f(a), f(x)) = d'(f(a), y) < \epsilon$$

as desired. □

Theorem 7.2.4

Let $c < \frac{-(5+2\sqrt{5})}{4}$. Then, $Q_c : \Lambda \rightarrow \Lambda$ is chaotic.

Proof. (periodic point density) Note that $Q_c^n(x) = x \iff S(Q_c^n(x)) = S(x) \iff \sigma^n(S(x)) = S(x)$.

By prop. 7.2.3 applied to $S^{-1} : \Sigma \rightarrow \Lambda$, the periodic points for Q_c are dense in Λ .

(transitivity) Take $z \in \Sigma$ from prop. 7.1.6 such that $\{\sigma^K(z) : K \in \mathbb{N} \cup \{0\}\}$ is dense in Σ . Again by prop. 7.2.3, $\{S^{-1}(\sigma^K(z)) : K \in \mathbb{N} \cup \{0\}\}$ is dense in Λ .

Note: Say $S(x) = z$, we know $(S(Q_c^K(x))) = \sigma^K(S(x)) \iff Q_c^K(x) = S^{-1}(\sigma^K(S(x)))$

This, $\{Q_c^K(x) : K \in \mathbb{N} \cup \{0\}\}$ is dense in Λ . As in prop. 7.1.10, we have that Q_c is transitive.

(sensitivity) Recall that $\Lambda \subseteq I \setminus J_1 = I_0 \cup I_1$. Let $\beta > 0$ be less than the gap between I_0 and I_1 .

For $x, y \in \Lambda$ with $x \neq y$, suppose $S(x) \neq S(y)$. Then, there must exist a k where k^{th} term of $S(x)$ does not equal the k^{th} term of $S(y)$.

Hence, $|Q_c^k(x) - Q_c^k(y)| > \beta$ and Q_c is sensitive. □

Chapter 8

Sarkovskii's Theorem

Theorem 8.0.1 (period 3)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous. If f has a point with period 3, then f has a point with period n for all $n \in \mathbb{N}$.

Lecture 15
Feb 9

Proposition 8.0.2

Let $I \subseteq J$ be closed intervals and suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous. If $f(I) \supseteq J$, then $f(x)$ has a fixed point in I .

Proposition 8.0.3

Let I, J be closed intervals, $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous, and $f(I) \supseteq J$. Then, there exists a closed interval $I' \subseteq I$ such that $f(I') = J$.

We can now prove thm. 8.0.1.

Proof. Let $a \in \mathbb{R}$ be a period 3 point for $f(x)$. Say $f(a) = b, f(b) = c, f(c) = a$. WLOG, suppose $a < b$ and $a < c$.

Suppose $a < b < c$. The case where $a < c < b$ is left as an exercise.

Let $I = [a, b]$ and $J = [b, c]$. Then, $f(a) = b$ and $f(b) = c$ imply by IVT that $[b, c] = J \subseteq f(I)$. Likewise, $f(b) = c$ and $f(c) = a$ imply by IVT that $[a, c] = I \cup J \subseteq f(J)$.

Since $J \subseteq f(J)$, there exists a closed interval $A_1 \subseteq J$ such that $f(A_1) = J$ by prop. 8.0.3. Again, $A_1 \subseteq J = f(A_1)$, so there exists a closed interval $A_2 \subseteq A_1$ such that $f(A_2) = A_1$.

Now, fix $n > 3$. Repeating the above process, we can find $A_{n-2} \subseteq A_{n-3} \subseteq \dots \subseteq A_2 \subseteq A_1 \subseteq J$ such that $f(A_i) = A_{i-1}$. Now, $f(I) \supseteq J \supseteq A_{n-2}$ means there exists a closed interval $A_{n-1} \subseteq I$ such that $f(A_{n-1}) = A_{n-2}$.

Moreover, $f(J) \supseteq I \supseteq A_{n-1}$ which means there exists a closed interval $A_n \subseteq J$ such that $f(A_n) = A_{n-1}$.

We have $f^n(A_n) = J$ and $A_n \subseteq J$. By prop. 8.0.2, there exists $x_0 \in A_n$ such that $f^n(x_0) = x_0$.

Note: for $x_0 \in A_n$, $f(x_0) \in A_{n-1} \subseteq I$, $f^i(x_0) \in J$ for $i = 2, 3, \dots, n$.

For contradiction, suppose $f^i(x_0) = x_0$ for $i < n$.

Then, $\overbrace{f(x_0)}^{\in I} = \overbrace{f^{i+1}(x_0)}^{\in J} = b$ so $f(x_0) = b$, $f^2(x_0) = c$, and $f^3(x_0) = a$, which is a contradiction because $f^3(x_0) \in J$ but $a \notin J$. Hence, x_0 has period n .

That is, f has a periodic point with period n for all $n > 3$.

Further, $f(J) \supseteq J$ and so by prop. 8.0.2, f has a fixed point (aka period 1) in J .

Finally, $f(I) \supseteq J$ means $J = f(I')$ and $f(J) \supseteq I'$ means $f(J') = I'$. This implies $f^2(J') = f(I') = J \supset J'$. Therefore, we know there exists $x \in J'$ such that $f^2(x) = x$.

If $f(x) = x$, then $x \in J'$ and $f(x) \in I'$, meaning $x = b$. But, $f(b) \neq b = c$, contradiction.

Hence, x has period 2.


Therefore, since we already supposed f has a period 3 point, f has a period n point for all n . \square

Exercise 8.0.4. Complete the proof for the case where $a < c < b$.

↑ Lectures 14 and 15 adapted from Imaad ↑

Draw the continuous function

Lecture 16
Feb 12



Then, the orbit of 1 is $1 \mapsto 3 \mapsto 4 \mapsto 2 \mapsto 5 \mapsto 1$ and 1 has period 5.

Claim 8.0.5. f has no point with period 3.

Proof. Suppose that f has a point x with period 3. Then, $1 \leq x \leq 5$.

Suppose $x \in [1, 2]$. Then, $x \in [1, 2] \cap f^3([1, 2])$ since $x = f^3(x)$. But $f^3([1, 2]) = [2, 5]$, so $x = 2$. However, 2 has period 5 since it is on the same 5-cycle given above.

Suppose instead that $x \in [2, 3]$. Then, $x \in [2, 3] \cap f^3([2, 3]) = [2, 3] \cap [3, 5] = \{3\}$ which is also on the 5-cycle.

If $x \in [4, 5]$, then $x \in [4, 5] \cap f^3([4, 5]) = [4, 5] \cap [1, 4] = \{4\}$ which is, again, on the 5-cycle.

Finally, suppose that $x \in [3, 4]$. Then, $f([3, 4]) = [2, 4]$ and it is strictly decreasing. Further, $f([2, 4]) = [2, 5]$ and it is also strictly decreasing. Once more, $f([2, 5]) = [1, 5]$ and it is again strictly decreasing. Since f^3 is strictly decreasing, it has a unique fixed point in $[3, 4]$, but it is just the fixed point of f .

Since we have covered the entire interval $[1, 5]$, x must not exist. \square

Example 8.0.6. The function $f(x) = \begin{cases} 1 & x < -1 \\ -x & -1 \leq x \leq 1 \\ 1 & x > 1 \end{cases}$ has a period 1 point at $x = 0$, period 2 points $[-1, 1] \setminus \{0\}$, and no other periodic points.

Definition 8.0.7 (Sarkovskii ordering)

Start by ordering the odd numbers $3 \prec 5 \prec 7 \prec 9 \prec \dots$

Then, all those are $\dots \prec 2 \cdot 3 \prec 2 \cdot 5 \prec 2 \cdot 7 \prec \dots$

All those are $\dots \prec 2^2 \cdot 3 \prec 2^2 \cdot 5 \prec 2^2 \cdot 7 \prec \dots$

Complete the ordering as $\dots \prec 2^n \prec 2^{n-1} \prec \dots \prec 2^2 \prec 2 \prec 1$.

This is a total order on the natural numbers.

Example 8.0.8.

- $26 = 2 \cdot 13 \prec 2^2 \cdot 5 = 40$ because the exponent of 2 is smaller.
- $3072 = 2^{10} \cdot 3 \prec 2^5 = 32$ because powers of 2 are big.
- $n \prec 1$ for all n .
- $2^{15} \prec 2^3$ since the powers of 2 are ordered backwards.

Theorem 8.0.9 (Sarkovskii's theorem)

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Suppose $n \prec m$ in the Sarkovskii ordering. Then, if f has a point with period n , then it has a point with period m .

Chapter 9

Fractals

Definition 9.0.1

Define a few things from topology.

- For $\mathbf{x} \in \mathbb{R}^n$, the norm $\|\mathbf{x}\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$
- $d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|$ is our default metric on \mathbb{R}^n
- For $\mathbf{x} \in \mathbb{R}^n$, $\varepsilon > 0$, the open ball of radius ε centered at x is $B_\varepsilon(\mathbf{x}) = \{\mathbf{y} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{y}\| < \varepsilon\}$
- We say $U \subseteq \mathbb{R}^n$ is open if for all $\mathbf{x} \in U$, there exists $\varepsilon > 0$ such that $B_\varepsilon(\mathbf{x}) \subseteq U$.
- The boundary $\delta(A)$ of a set $A \subseteq \mathbb{R}^n$ is the closure of A without the interior of A .

Lecture 17
Feb 14

Definition 9.0.2 (topological dimension (zero case))


We say $S \subseteq \mathbb{R}^n$ has topological dimension $\dim_t S = 0$ if for all $\mathbf{x} \in S$, there exists arbitrarily small open sets $U \ni \mathbf{x}$ such that $\delta(U) \cap S = \emptyset$.

Example 9.0.3. Let $X = \{\cdot\}$. Then, since we can draw balls $B_\varepsilon(\cdot)$ separating each point, $\dim_t X = 0$.

Example 9.0.4. $X = \{\frac{1}{n} : n \in \mathbb{N}\} \cup \{0\}$ has topological dimension 0.


Definition 9.0.5 (topological dimension (non-zero case))

A set $S \subseteq \mathbb{R}^n$ has topological dimension $k \in \mathbb{N}$ if for all $\mathbf{x} \in S$, there exists arbitrarily small $U \ni \mathbf{x}$ such that $\delta(U) \cap S$ has topological dimension $k-1$, where k is minimal with this property.

Example 9.0.6. Consider a line $X = \text{---}$. Then, since any ball's boundary  creates an intersection of two points (i.e., a set with topological dimension 0), we know that $\dim_t X = 1$.


Example 9.0.7. Let X be a circle . Again, any ball's boundary  still only has two intersecting points, so $\dim_t X = 1$.

Example 9.0.8. Let X be a filled 2D region.



Then, the intersection of a ball's boundary will give either a circle or an arc, which have topological dimension 1, so the region has topological dimension 2.

Example 9.0.9. Let X be a non-filled sphere.



Then, the intersection of a 3D ball's boundary will give a circle, which has topological dimension 1, so $\dim_t X = 2$.

Example 9.0.10. Let X be a filled sphere.

Then, a 3D ball's boundary's intersection is either a hollow sphere or a spherical cap, which each have topological dimension 2, so $\dim_t X = 3$.

Definition 9.0.11 (fractal dimension)

We say $S \subseteq \mathbb{R}^n$ is self-similar if S may be divided into K congruent subsets, each of which may be magnified by a fixed M to yield S itself.

The fractal dimension of S is given by $\dim_f S = \frac{\ln K}{\ln M}$.

Definition 9.0.12 (fractal)

A fractal is a self-similar $S \subseteq \mathbb{R}^n$ such that $\dim_f S > \dim_t S$.

Example 9.0.13. Let $X = \bullet\text{---}\bullet$ be a line. Then, since we can divide it into n smaller lines each of size $\frac{1}{n}$, it has fractal dimension $\dim_f X = \frac{\ln n}{\ln n} = 1$. The topological dimension is $\dim_t X = 1$.

So this is not a fractal, and is indeed just boring (not a fractal).

Example 9.0.14 (Sierpinski triangle). Let X be the Sierpinski triangle, i.e., the limiting point of the process



Then, the topological dimension is $\dim_t X = 1$ because, in the limit, any ball will touch only single points. In particular, we can imagine balls touching the three points of a triangle.

However, the fractal dimension is $\dim_f X = \frac{\ln 3}{\ln 2} \approx 1.58 > 1$ because each step is consisted of 3 previous steps scaled by $\frac{1}{2}$. so X is a fractal!

List of Named Results

1.2.5	Proposition (convergence implies boundedness)	6
1.2.7	Proposition (limit laws)	7
1.2.10	Theorem (completeness of \mathbb{R})	8
1.3.6	Theorem (Banach contraction mapping theorem)	9
3.1.5	Theorem (attracting fixed point theorem)	17
3.1.7	Theorem (repelling fixed point theorem)	18
5.0.4	Theorem (Cantor sets are closed)	26
6.1.12	Proposition (sequential characterization of continuity in metric spaces)	31
6.2.3	Theorem (monotone convergence theorem)	31
6.2.4	Lemma (nested intervals lemma)	32
8.0.1	Theorem (period 3)	38
8.0.9	Theorem (Sarkovskii's theorem)	40

Index of Defined Terms

- bifurcation, 23
- boundary, 41
- Cantor set, 26
- Cantor space, 29
- chaos, 36
- cobweb plot, 12
- contraction, 9
- density, 34
- doubling function, 5
- dynamical system, 35
- fixed point, 4
 - attracting, 17
 - weakly, 19
 - neutral, 17
 - repelling, 17
 - weakly, 20
- fractal, 43
- fractal dimension, 42
- function
 - continuity, 8
 - homeomorphism, 31
 - interval
 - closure, 9
 - iteration, 3
 - itinerary, 28
 - metric, 29
 - metric space, 29
 - continuity, 30
 - convergence, 31
 - non-isolated, 17
 - norm, 41
 - open, 41
 - open ball, 41
 - orbit, 3
 - constant, 4
 - eventually periodic, 5
 - periodic, 5
 - period, 5
 - periodic point, 5
 - attracting, 20
 - neutral, 20
 - repelling, 20
 - periodicity, 5
 - quadratic family, 22
 - Sarkovskii ordering, 40
 - self-similar, 42
 - sensitive, 35
 - sequence
 - bounded, 6
 - Cauchy, 7
 - convergence, 6
 - shift map, 30
 - strongly-Cauchy, 10
 - tangent bifurcation, 23
 - topological dimension, 41
 - transitivity, 35