Tutorial 2

Research Methods for Political Science - PO3110

Andrea Salvi

1 & 2 October 2019

Trinity College Dublin,

https://andrsalvi.github.io/research-methods/

Table of contents

- 1. Homework 1: answer keys and review
- 2. Key Concepts from the Lecture
- 3. In-class Exercise

Homework 1: answer keys and

review

Homework 1

- · Comments?
- Feedback on previous tutorial and material?
- · Homework Walk-through

Key Concepts from the Lecture

Measures of Dispersion and CLT

- · Standard deviation: spread of the sample;
- Standard error of the mean: spread of the means of many samples. That is, standard deviation of the sampling distribution;
- Central limit theorem: The mean of a large number of random samples will be normally distributed, regardless of the underlying distribution of that variable. That is, the sampling distribution will take the form of a normal distribution!

Calculating Dispersion

- 1. Estimate Mean: $\bar{x} = \frac{\sum x}{n}$
- 2. Sum of Squared Errors (SS): $\sum (x \bar{x})^2$
- 3. Estimate Variance: $\sigma^2 = \frac{SS}{n-1} = \frac{\sum (x-\bar{x})^2}{n-1}$
- 4. Estimate Standard Deviation: $\sigma = \sqrt{\frac{\sum (x \bar{x})^2}{n-1}} = \sqrt{\sigma^2}$
 - · Remember that it is **NEVER** negative.
 - · It can be 0 though in case of a uniform distribution.
- 5. Estimate standard error of the mean: $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$

Calculating Dispersion

- 1. Estimate Mean: $\bar{x} = \frac{\sum x}{n}$
- 2. Sum of Squared Errors (SS): $\sum (x \bar{x})^2$
- 3. Estimate Variance: $\sigma^2 = \frac{SS}{n-1} = \frac{\sum (x-\bar{x})^2}{n-1}$
- 4. Estimate Standard Deviation: $\sigma = \sqrt{\frac{\sum (x \bar{x})^2}{n-1}} = \sqrt{\sigma^2}$
 - · Remember that it is **NEVER** negative.
 - · It can be 0 though in case of a uniform distribution.
- 5. Estimate standard error of the mean: $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}}$

Hands-on exercise: Estimate \bar{x} , σ , $sd(\bar{X})$ for the following sample: 9, 2, 5, 4

Mean:
$$\bar{x} = \frac{9+2+5+4}{4} = \frac{20}{4} = 5$$

Mean:
$$\bar{x} = \frac{9+2+5+4}{4} = \frac{20}{4} = 5$$

 $(9-5)^2 = 4^2 = 16$
 $(2-5)^2 = (-3)^2 = 9$
 $(5-5)^2 = (0)^2 = 0$
 $(4-5)^2 = (1)^2 = 1$

Mean:
$$\bar{x} = \frac{9+2+5+4}{4} = \frac{20}{4} = 5$$

 $(9-5)^2 = 4^2 = 16$
 $(2-5)^2 = (-3)^2 = 9$
 $(5-5)^2 = (0)^2 = 0$
 $(4-5)^2 = (1)^2 = 1$
 $16+9+0+1=26$

Mean:
$$\bar{x} = \frac{9+2+5+4}{4} = \frac{20}{4} = 5$$

 $(9-5)^2 = 4^2 = 16$
 $(2-5)^2 = (-3)^2 = 9$
 $(5-5)^2 = (0)^2 = 0$
 $(4-5)^2 = (1)^2 = 1$
 $16+9+0+1=26$
Standard deviation: $\sigma = \sqrt{\frac{\sum (x-\bar{x})^2}{n-1}} = \sqrt{\frac{26}{4-1}} = 2.94392$

Mean:
$$\bar{x} = \frac{9+2+5+4}{4} = \frac{20}{4} = 5$$

 $(9-5)^2 = 4^2 = 16$
 $(2-5)^2 = (-3)^2 = 9$
 $(5-5)^2 = (0)^2 = 0$
 $(4-5)^2 = (1)^2 = 1$
 $16+9+0+1=26$
Standard deviation: $\sigma = \sqrt{\frac{\sum(x-\bar{x})^2}{n-1}} = \sqrt{\frac{26}{4-1}} = 2.94392$
Standard error: $sd(\bar{X}) = \frac{\sigma}{\sqrt{n}} = \frac{2.94392}{\sqrt{4}} = 1.47196$

5

- Download "parlgov elections.xlsx" https://tinyurl.com/datamt2
- Download Parlgov Elections Codebook https://tinyurl.com/codebookmt2
- · You can work in pairs

- · Load the Data
- · What do they represent? What are the observations here?
- · Describe the variables we have.
- · How are missing values coded?
- Subset the data in order to select just the Irish elections.
- Arrange the data based on "left_right" from Left to Right.

- Re-code the "election_type" variable into a numeric variable called "election_type_num" where "1" is "parliament" and 2 is "ep". Assign them a label accordingly.
- Create a new binary variable (0,1) called "left_right_binary" where "right" = 1 and "left" = 0. How would you do that? (at least two ways!)
- "Split" the data-set based on the "election_type" variable (Hint: Data -> Split File). Now try to calculate the mean, standard deviation and Standard Error of the Mean for the "vote_share" variable. What happened?
- Plot a histogram of your choice that conveys meaningful information.
- How do you get rid of the "split"?