

Algebraic Foundations of Computer Science.

Computational Introduction to Number Theory (II)

Ferucio Laurențiu Tiplea

Department of Computer Science
“AL.I.Cuza” University of Iași
Iași, Romania
E-mail: ftiplea@mail.dntis.ro

Spring 2015

Outline

*Algebraic
Foundations of
Computer Science
(AFCS)*

*Prof.Dr. F.L.
Tiplea*

*Linear
congruential
equations*

*The Chinese
remainder
theorem*

*Quadratic
residues*

*The Legendre
symbol*

The Jacobi symbol

Course readings

- 1 *Linear congruential equations*
- 2 *The Chinese remainder theorem*
- 3 *Quadratic residues*
- 4 *The Legendre symbol*
- 5 *The Jacobi symbol*
- 6 *Course readings*

Linear congruential equations

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Theorem 1

Let $a, b, m \in \mathbb{Z}$ with $m \geq 1$. Then, the equation

$$ax \equiv b \pmod{m}$$

is solvable in \mathbb{Z} iff $(a, m) \mid b$. Moreover, if it is solvable, then it has exactly (a, m) solution in \mathbb{Z}_m which are of the form

$$\left(x_0 + i \frac{m}{(a, m)} \right) \pmod{m},$$

where x_0 is an arbitrary integer solution and $0 \leq i < (a, m)$.

Example 2

The equation

$$5x \equiv 25 \pmod{10}$$

has $(5, 10) = 5$ solutions in \mathbb{Z}_{10} : 1, 3, 5, 7, 9.

Linear congruential equations

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Algorithm 1: Solving linear congruential equations

input : $m \geq 1$ and $a, b \in \mathbb{Z}$;

output: all solutions modulo m of $ax \equiv b \pmod{m}$;

begin

compute $\gcd(a, m) := \alpha a + \beta m$;

if $\gcd(a, m) \mid b$ **then**

$b' := b / \gcd(a, m)$;

$x_0 := \alpha b'$;

for $i := 0$ **to** $\gcd(a, m) - 1$ **do**

print $((x_0 + im / \gcd(a, m)) \pmod{m})$

else

print “no integer solutions”

The Chinese remainder theorem

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

According to D.Wells, the following problem was posed by Sun Tsu Suan-Ching (4th century AD):

There are certain things whose number is unknown. Repeatedly divided by 3, the remainder is 2; by 5, the remainder is 3; and by 7, the remainder is 2. What will be the number?

The mathematical form of this problem is:

$$\begin{cases} x \equiv 2 \pmod{3} \\ x \equiv 3 \pmod{5} \\ x \equiv 2 \pmod{7} \end{cases}$$

This system of equations has a least integer solution which is $x = 23$.

The Chinese remainder theorem

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Theorem 3 (Chinese Remainder Theorem)

Let $k \geq 1$ and m_1, \dots, m_k be pairwise co-prime integers. Then, for any $b_1, \dots, b_k \in \mathbb{Z}$, the following system (S) of equations has a unique solution modulo $m_1 \cdots m_k$

$$(S) \begin{cases} x \equiv b_1 \pmod{m_1} \\ \dots \\ x \equiv b_k \pmod{m_k} \end{cases}$$

The solution can be obtained as follows:

- compute $c_i = \prod_{j=1, j \neq i}^k m_j$;
- compute an integer solution x_i of the equation $c_i x \equiv b_i \pmod{m_i}$, for any i ;
- $x = (c_1 x_1 + \dots + c_k x_k) \pmod{(m_1 \cdots m_k)}$ is the unique solution modulo $m_1 \cdots m_k$ of the system.

The Chinese remainder theorem: example

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Example 4

Let (S) be the system

$$(S) \begin{cases} x \equiv 2 \pmod{3} \\ x \equiv 3 \pmod{5} \\ x \equiv 2 \pmod{7} \end{cases}$$

Then:

- $c_1 = 35$, $c_2 = 21$, and $c_3 = 15$;
- $x_1 = 1$ is a solution of $35x \equiv 2 \pmod{3}$;
- $x_2 = 3$ is a solution of $21x \equiv 3 \pmod{5}$;
- $x_3 = 2$ is a solution of $15x \equiv 2 \pmod{7}$;
- $x = (35 \cdot 1 + 21 \cdot 3 + 15 \cdot 2) \pmod{105} = 128 \pmod{105} = 23$ is the unique solution modulo 105 of the system (S) .

The Chinese remainder theorem: application

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

There is an important application of CRT to the problem of solving equations of the form $f(x) \equiv 0 \pmod{m}$, where $f(x)$ is a polynomial with integer coefficients and variables x .

Theorem 5

Let $f(x)$ be a polynomial with integer coefficients, and m_1, \dots, m_k be pairwise co-prime integers. Then, $a \in \mathbb{Z}$ is a solution to the equation

$$f(x) \equiv 0 \pmod{m_1 \cdots m_k} \quad (1)$$

if and only if a is a solution to each of the equations

$$f(x) \equiv 0 \pmod{m_i}, \quad 1 \leq i \leq k. \quad (2)$$

Moreover, the number of solutions in $\mathbb{Z}_{m_1 \cdots m_k}$ of the equation (1) is the product of the numbers of solutions in \mathbb{Z}_{m_i} of the equations (2).

The Chinese remainder theorem: application

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Example 6

- ❶ The equation

$$x^2 \equiv 1 \pmod{p},$$

where $p > 2$ is a prime number, has exactly 2 solutions in \mathbb{Z}_p , namely $x = 1$ and $x = p - 1$.

- ❷ The equation

$$x^2 \equiv 1 \pmod{p_1 \cdots p_k},$$

where p_1, \dots, p_k are distinct odd primes ($k \geq 2$), has exactly 2^k solutions in $\mathbb{Z}_{p_1 \cdots p_k}$.

Quadratic residues - motivation

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Proposition 1 (Solving quadratic congruences)

Let $p > 2$ be a prime and $a, b, c \in \mathbb{Z}$ such that $(a, p) = 1$. Then, the quadratic congruence

$$ax^2 + bx + c \equiv 0 \pmod{p}$$

has

- two roots in \mathbb{Z}_p , if $\Delta \equiv y^2 \pmod{p}$ for some $y \in \mathbb{Z}$ with $p \nmid y$;
- one root in \mathbb{Z}_p , if $\Delta \equiv 0 \pmod{p}$;
- no roots, otherwise,

where $\Delta = b^2 - 4ac$.

How hard is to decide if a given $a \in \mathbb{Z}_p^*$ satisfies $a \equiv y^2 \pmod{p}$ for some $y \in \mathbb{Z}$?

Quadratic residues and non-residues

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Definition 7

Let $p > 2$ be a prime and $a \in \mathbb{Z}$ non-divisible by p . a is called a **quadratic residue modulo p** if $a \equiv x^2 \pmod{p}$ for some integer x .

If a is neither divisible by p nor a quadratic residue modulo p then a is called a **quadratic non-residue modulo p** .

Remark 1

An integer a non-divisible by a prime $p > 2$ is a quadratic (non-)residue modulo p if and only if $a \pmod{p}$ is a quadratic (non-)residue modulo p .

Denote

- $QR_p = \{a \in \mathbb{Z}_p^* | a \text{ is a quadratic residue modulo } p\}$
- $QNR_p = \{a \in \mathbb{Z}_p^* | a \text{ is a quadratic non-residue modulo } p\}$

Quadratic residues. Basic properties

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Proposition 2

Let $p > 2$ be a prime. Then, $|QR_p| = |QNR_p| = \frac{p-1}{2}$.

Proposition 3

Let $p > 2$ be a prime. Then:

- 1. $a, b \in QR_p \Rightarrow (ab \bmod p) \in QR_p$;
- 2. $a \in QR_p \wedge b \in QNR_p \Rightarrow (ab \bmod p) \in QNR_p$;
- 3. $a, b \in QNR_p \Rightarrow (ab \bmod p) \in QR_p$.

Theorem 8 (Euler's Criterion)

Let $p > 2$ be a prime and $a \in \mathbb{Z}_p^*$. Then,

- 1. $a \in QR_p$ if and only if $a^{\frac{p-1}{2}} \equiv 1 \bmod p$;
- 2. $a \in QNR_p$ if and only if $a^{\frac{p-1}{2}} \equiv -1 \bmod p$.

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Introduced by Adrien-Marie Legendre in 1798 when trying to prove the law of quadratic reciprocity.

Definition 9

Let $p > 2$ be a prime. The **Legendre symbol** of $a \in \mathbb{Z}$, denoted $\left(\frac{a}{p}\right)$, is defined by

$$\left(\frac{a}{p}\right) = \begin{cases} 0, & \text{if } p|a \\ 1, & \text{if } p \nmid a \text{ and } a \text{ is a quadratic residue modulo } p \\ -1, & \text{if } p \nmid a \text{ and } a \text{ is a quadratic non-residue modulo } p \end{cases}$$

Remark that the Legendre symbol is only defined for primes $p > 2$. For $p = 2$, all even integers are divisible by p and all odd integers are quadratic residues modulo p .

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Proposition 4

Let $p > 2$ be a prime and $a, b \in \mathbb{Z}$. If $a \equiv b \pmod{p}$ then $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$. Therefore, $\left(\frac{a}{p}\right) = \left(\frac{a \bmod p}{p}\right)$.

Proposition 5

Let $p > 2$ be a prime. Then, for any $a \in \mathbb{Z}$, $\left(\frac{a}{p}\right) \equiv a^{\frac{p-1}{2}} \pmod{p}$.

Proposition 6

Let $p > 2$ be a prime. Then, for any $a, b \in \mathbb{Z}$, $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$.

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

According to the above properties, computing the Legendre symbol modulo p comes down to computing $\left(\frac{-1}{p}\right)$ and $\left(\frac{q}{p}\right)$, for any prime q with $2 \leq q < p$.

Proposition 7

Let $p > 2$ be a prime. Then,

$$\left(\frac{-1}{p}\right) = (-1)^{\frac{p-1}{2}} = \begin{cases} 1, & \text{if } p \equiv 1 \pmod{4} \\ -1, & \text{if } p \equiv 3 \pmod{4} \end{cases}$$

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Theorem 10 (Gauss' Criterion)

Let $p > 2$ be a prime and $a \in \mathbb{Z}$ non-divisible by p . Then,
 $\left(\frac{a}{p}\right) = (-1)^r$, where

$$r = |\{i \in \{1, \dots, (p-1)/2\} \mid ia \bmod p > p/2\}|.$$

Proposition 8

Let $p > 2$ be a prime. Then,

$$\left(\frac{2}{p}\right) = (-1)^{\frac{p^2-1}{8}} = \begin{cases} 1, & \text{if } p \equiv \pm 1 \bmod 8 \\ -1, & \text{if } p \equiv \pm 3 \bmod 8 \end{cases}$$

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Theorem 11 (Quadratic reciprocity law)

Let $p, q > 2$ be distinct primes. Then,

$$\left(\frac{q}{p}\right) \left(\frac{p}{q}\right) = (-1)^{\frac{p-1}{2} \cdot \frac{q-1}{2}}.$$

Equivalently,

$$\left(\frac{q}{p}\right) = \begin{cases} -\left(\frac{p}{q}\right), & \text{if } p, q \equiv 3 \pmod{4} \\ \left(\frac{p}{q}\right), & \text{otherwise} \end{cases}$$

Example 12

$$\left(\frac{7}{59}\right) = -\left(\frac{59}{7}\right) = -\left(\frac{3}{7}\right) = \left(\frac{7}{3}\right) = \left(\frac{1}{3}\right) = 1$$

The Legendre symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Basic rules for computing the Legendre symbol (review):

• if $a \equiv b \pmod{p}$ then $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right)$

• $\left(\frac{ab}{p}\right) = \left(\frac{a}{p}\right) \left(\frac{b}{p}\right)$

• $\left(\frac{1}{p}\right) = 1$

• $\left(\frac{-1}{p}\right) = \begin{cases} 1, & \text{if } p \equiv 1 \pmod{4} \\ -1, & \text{if } p \equiv 3 \pmod{4} \end{cases}$

• $\left(\frac{2}{p}\right) = \begin{cases} 1, & \text{if } p \equiv \pm 1 \pmod{8} \\ -1, & \text{if } p \equiv \pm 3 \pmod{8} \end{cases}$

• $\left(\frac{q}{p}\right) = \begin{cases} -\left(\frac{p}{q}\right), & \text{if } p \equiv q \equiv 3 \pmod{4} \\ \left(\frac{p}{q}\right), & \text{if } p \equiv 1 \pmod{4} \text{ or } q \equiv 1 \pmod{4} \end{cases}$

for any distinct primes $p, q > 2$ and $a, b \in \mathbb{Z}$.

The Jacobi symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Introduced by Carl Gustav Jacob Jacobi in 1837 as a generalization of the Legendre symbol.

Definition 13

Let $n > 0$ be an odd integer. The **Jacobi symbol** of $a \in \mathbb{Z}$, denoted $\left(\frac{a}{n}\right)$, is defined by

$$\left(\frac{a}{n}\right) = \begin{cases} 1, & \text{if } n=1 \\ \left(\frac{a}{p_1}\right)^{e_1} \cdots \left(\frac{a}{p_k}\right)^{e_k}, & \text{otherwise} \end{cases}$$

where $n = p_1^{e_1} \cdots p_k^{e_k}$ is the prime factorization of n .

Remark 2

- The Jacobi symbol is defined only for odd integers $n > 0$.
- $(a, n) = 1$ if and only if $\left(\frac{a}{n}\right) \neq 0$, for all $a \in \mathbb{Z}$ and $n > 0$ odd.

The Jacobi symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Theorem 14

The following properties hold:

1 if $a \equiv b \pmod n$ then $\left(\frac{a}{n}\right) = \left(\frac{b}{n}\right)$

2 $\left(\frac{ab}{n}\right) = \left(\frac{a}{n}\right) \left(\frac{b}{n}\right)$

3 $\left(\frac{1}{n}\right) = 1$

4 $\left(\frac{-1}{n}\right) = \begin{cases} 1, & \text{if } n \equiv 1 \pmod 4 \\ -1, & \text{if } n \equiv 3 \pmod 4 \end{cases}$

5 $\left(\frac{2}{n}\right) = \begin{cases} 1, & \text{if } n \equiv \pm 1 \pmod 8 \\ -1, & \text{if } n \equiv \pm 3 \pmod 8 \end{cases}$

6 $\left(\frac{m}{n}\right) = \begin{cases} -\left(\frac{n}{m}\right), & \text{if } n \equiv m \equiv 3 \pmod 4 \\ \left(\frac{n}{m}\right), & \text{if } n \equiv 1 \pmod 4 \text{ or } m \equiv 1 \pmod 4 \end{cases}$

for any distinct odd integers $n, m > 0$ and $a, b \in \mathbb{Z}$.

The Jacobi symbol

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

Algorithm 2: Computing the Jacobi symbol

input : integer a and odd integer $n > 0$;

output: $\left(\frac{a}{n}\right)$

begin

$b := a \bmod n$; $c := n$; $s := 1$;

while $b \geq 2$ **do**

while $4|b$ **do** $b := b/4$;

if $2|b$ **then**

if $c \bmod 8 \in \{3, 5\}$ **then** $s := -s$;

$b := b/2$;

if $b = 1$ **then quit**;

if $b \bmod 4 = 3 = c \bmod 4$ **then**

$s := -s$;

$(b, c) := (c \bmod b, b)$;

return $s \cdot b$;

Course readings

Algebraic
Foundations of
Computer Science
(AFCS)

Prof.Dr. F.L.
Tiplea

Linear
congruential
equations

The Chinese
remainder
theorem

Quadratic
residues

The Legendre
symbol

The Jacobi symbol

Course readings

- F.L. Țiplea: *Fundamentele Algebrice ale Informaticii*, Ed. Polirom, Iași, 2006, pag. 164–172.
- M. B. Nathanson: *Elementary Methods in Number Theory*, Graduate Texts in Mathematics, Springer-Verlag, 2000, pag. 100–120.