

Advanced Signal Processing

Fourier Analysis and Wavelet Transform

Gustavo Andrade Miranda PhD

gxavier.andradem@ug.edu.ec

Facultad de Ingeniería en Sistemas, Electrónica e Industrial
Universidad Técnica de Ambato



PROGRAMA DE MAESTRÍA EN TELECOMUNICACIONES

UNIVERSIDAD
TÉCNICA DE AMBATO

Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier



Oscillations Everywhere

- sustainable dynamic systems exhibit oscillatory behaviour
- intuitively things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...



Oscillations Everywhere

- sustainable dynamic systems exhibit oscillatory behaviour
- intuitively things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...



Oscillations Everywhere

- sustainable dynamic systems exhibit oscillatory behaviour
- intuitively things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...



Oscillations Everywhere

- sustainable dynamic systems exhibit oscillatory behaviour
- intuitively things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...




Oscillations Everywhere

- sustainable dynamic systems exhibit oscillatory behaviour
- intuitively things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...




Circular motion

- period P
- frequency $f = \frac{1}{P}$



Circular Motion Cosine



UNIVERSIDAD
TÉCNICA DE AMBATO

Circular Motion Cosine-Sine



UNIVERSIDAD
TÉCNICA DE AMBATO

Circular Motion Cosine-Sine




UNIVERSIDAD
TÉCNICA DE AMBATO


Detect sinusoids

The human body has two receptors for sinusoidal

Cochlea (inner ear)



rods and cones (retina)




- air pressure sinusoids
- frequencies from 20 Hz to 20KHz

- electromagnetic sinusoids
- frequencies from 430 THz to 790 THz

Frequency Domain

- humans analyze complex signals in terms of their sinusoidal components
- we can build instruments that “resonate” at one or multiple frequencies
- the “frequency domain” seems to be as important as the time domain



UNIVERSIDAD
TÉCNICA DE AMBATO

Fourier Analysis

Can we decompose any signal into sinusoidal elements??

Yes, FOURIER ANALYSIS

analysis

- from time to frequency domain
- find the contribution of different frequencies
- discover “hidden” signal properties

synthesis

- from frequency to time domain
- create signals with known frequency content
- fit signals to specific frequency regions

Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier




Mathematical Setup

- Fourier analysis is a simple change of basis
- a change of basis is a change of perspective
- a good basis can reveal information




Unknown Signal



A
ATO

Unknown Signal



A
ATO

Fourier Basis for \mathbb{C}^N

Claim: the set of N signals in \mathbb{C}^N

$$w_k[n] = e^{\frac{2\pi}{N} nk}, \quad n, k = 0, 1, 2, \dots, N-1$$

is a orthogonal basis in \mathbb{C}^N

- k is the index of the signal
- n is the index of the element

$$\omega = \frac{2\pi k}{N}$$



Fourier Basis - Vector Notation

In vector notation

$$\{\mathbf{w}^{(k)}\}_{k=0,1,2,\dots,N-1}$$

with


$$w_k[n] = e^{\frac{2\pi}{N}nk}, \quad n, k = 0, 1, 2, \dots, N-1$$

is a orthogonal basis in \mathbb{C}^N


$$\omega = \frac{2\pi k}{N}$$




Complex Generating Machine




Complex Generating Machine




Complex Generating Machine




Complex Generating Machine




Complex Generating Machine




Complex Generating Machine




Basis vector



Basis vector




Basis vector




UNIVERSIDAD
DE AMBATO


Basis vector




Basis vector




Basis vector




Basis vector




Basis vector




Basis vector



Basis vector



Basis vector




Remarks

- N orthogonal vectors \rightarrow basis for \mathbb{C}^N
- vectors are not orthonormal. Normalization should be $1/\sqrt{N}$




Mistery signal



UNIVERSIDAD
TÉCNICA DE AMBATO

Mistery signal



Mistery signal

$$x[n] = \cos(\omega n + \phi) + \text{noise}$$

with



$$\phi = 0$$

$$\omega = \frac{2\pi}{1024} 64$$

Peak occurs in 64




Mistery signal revelead



UNIVERSIDAD
TÉCNICA DE AMBATO

Example



TÉCNICA DE AMBATO

Mistery signal

- DFT main peak $k=22$
- 22 cycles over 2904 months

$$\text{period} = \frac{2904}{22} \approx 11 \text{ years}$$



Labelling the frequency axis


if we know the clock of the system

- fastest (positive) frequency is $\omega = \pi$
- sinusoids at $\omega = \pi$ need two sample to do full revolution
- time between samples $T_s = 1/F_s$
- real-world period for fastest sinusoid $2T_s$
- real-world frequency for fastest sinusoid $F_s/2$




Train Whistle

32768 samples (the "clock" of the system $F_s = 8000\text{Hz}$)



Train Whistle


32768 samples (the "clock" of the system $F_s = 8000\text{Hz}$)



TÉCNICA DE AMBATO

Train Whistle

the "clock" of the system $F_s = 8000\text{Hz}$



UTA
UNIVERSIDAD
TÉCNICA DE AMBATO

Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier






Analysis of musical instrument




- Pitch - Frequency
- Harmonics??
- What give the timbre of an instrument??



Musical instrument



Musical instrument



Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier




Dual-tone Multi Frequency Dialing


	1209Hz	1336Hz	1477Hz
697Hz	1	2	3
770Hz	4	5	6
852Hz	7	8	9
941Hz	*	0	#



Dialing 1-5-9



Dialing 1-5-9



Problem

- time representation hide the frequency
- frequency representation hide the time, We know frequency but why don't know when it happen



Solution

Idea:

- take small signal pieces of length L
- compute DFT of each piece

$$X[m; k] = \sum_{n=0}^{L-1} x[m + n] e^{-j \frac{2\pi}{L} nk}$$



Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier




Spectrogram

Idea:

- color code the magnitude: dark is small, white is large
- use $10\log_{10}(|X[m; k]|)$
- plot spectra slices one after other



Dialing 1-5-9 -spectrogram



Labelling the Spectrogram

- width of the analysis windows??
- position of the windows (overlapping??)
- shape of the window(weighting samples??)



wideband vs narrowband


Long window: narrowband spectrogram

- ▶ long window \Rightarrow more DFT points \Rightarrow more frequency resolution
- ▶ long window \Rightarrow more “things can happen” \Rightarrow less precision in time

Short window: wideband spectrogram


- ▶ short window \Rightarrow many time slices \Rightarrow precise location of transitions
- ▶ short window \Rightarrow fewer DFT points \Rightarrow poor frequency resolution

wideband vs narrowband




UNIVERSIDAD
TÉCNICA DE AMBATO

wideband vs narrowband



wideband vs narrowband



Outline

Introduction to Fourier analysis

DFT - Change of basis

Musical Instruments

Short Time Fourier Transform

Spectrogram

Advanced concepts in Fourier




Discrete Fourier Series (DFS)

- DFS = DFT with periodicity explicit
- DFS maps an N-periodic signal onto an N-periodic sequence of Fourier coefficients
- the DFS of an N-periodic signal is mathematically equivalent to the DFT of one period
- DFS: only N Fourier coefficients capture all the information



Karplus-Strong revisited and DFS




$$y[n] = \alpha y[n - M] + x[n]$$

Karplus-Strong revisited and DFS

- ▶ choose a signal $\bar{x}[n]$ that is nonzero only for $0 \leq n < M$
- ▶ $\alpha = 1$ (for now)


$$y[n] = \underbrace{\bar{x}[0], \bar{x}[1], \dots, \bar{x}[M-1]}_{\text{1st period}}, \underbrace{\bar{x}[0], \bar{x}[1], \dots, \bar{x}[M-1]}_{\text{2nd period}}, \underbrace{\bar{x}[0], \bar{x}[1], \dots}_{\dots}$$

Karplus-Strong revisited and DFS




UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS




UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS




UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS




UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS



UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS




$k = 1$



UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS




$k = 2$



UNIVERSIDAD
TÉCNICA DE AMBATO

Karplus-Strong revisited and DFS



DFT of L periods

$$\begin{aligned}
 X_L[k] &= \sum_{n=0}^{LM-1} y[n] e^{-j \frac{2\pi}{LM} nk} \quad k = 0, 1, 2, \dots, LM-1 \\
 &= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} y[n + pM] e^{-j \frac{2\pi}{LM} (n+pM)k} \\
 &= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} y[n] e^{-j \frac{2\pi}{LM} nk} e^{-j \frac{2\pi}{L} pk} \\
 &= \left(\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} \right) \sum_{n=0}^{M-1} \bar{x}[n] e^{-j \frac{2\pi}{LM} nk}
 \end{aligned}$$



UNIVERSIDAD
TÉCNICA DE AMBATO

DFT of L periods

$$\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} = \begin{cases} L & \text{if } k \text{ multiple of } L \\ 0 & \text{otherwise} \end{cases}$$

(remember the orthogonality proof for the DFT basis)



DFT of L periods

$$X_L[k] = \begin{cases} L\bar{X}[k/L] & \text{if } k = 0, L, 2L, 3L, \dots \\ 0 & \text{otherwise} \end{cases}$$



DFT of L periods

- again, all the spectral information for a periodic signal is contained in the DFT coefficients of a single period
- to stress the periodicity of the underlying signal, we use the term DFS



Discrete-time Fourier Transform (DTFT)

- Start with the DFT. What happens when $N \rightarrow \infty$?
- $(2\pi/N)k$ becomes denser in $[0, 2\pi]$
- in the limit $(2\pi/N)k \rightarrow \omega$

$$\sum_n x[n] \exp^{-j\omega n} \quad \omega \in \mathbb{R}$$



Discrete-time Fourier Transform (DTFT)

- Start with the DFT. What happens when $N \rightarrow \infty$?
- $(2\pi/N)k$ becomes denser in $[0, 2\pi]$
- in the limit $(2\pi/N)k \rightarrow \omega$

$$\sum_n x[n] \exp^{-j\omega n} \quad \omega \in \mathbb{R}$$



Discrete-time Fourier Transform (DTFT)

Formal definition:

- ▶ $x[n] \in \ell_2(\mathbb{Z})$
- ▶ define the *function* of $\omega \in \mathbb{R}$

$$F(\omega) = \underbrace{\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}}$$

- ▶ inversion (when $F(\omega)$ exists):

$$\underbrace{x[n]}_{=} = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\omega) e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

Discrete-time Fourier Transform (DTFT)


- ▶ $F(\omega)$ is 2π -periodic
- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$

Discrete-time Fourier Transform (DTFT)

$$x[n] = \alpha^n u[n], \quad |\alpha| < 1$$



Discrete-time Fourier Transform (DTFT)

DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$



Discrete-time Fourier Transform (DTFT)

Formal definition:

- ▶ $x[n] \in \ell_2(\mathbb{Z})$
- ▶ define the *function* of $\omega \in \mathbb{R}$

$$F(\omega) = \underbrace{\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}}$$

- ▶ inversion (when $F(\omega)$ exists):

$$\underbrace{x[n]}_{=} = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\omega) e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

Discrete-time Fourier Transform (DTFT)


- ▶ $F(\omega)$ is 2π -periodic
- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$

Discrete-time Fourier Transform (DTFT)

$$x[n] = \alpha^n u[n], \quad |\alpha| < 1$$




Discrete-time Fourier Transform (DTFT)

DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$


$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$



Plotting the DTFT




Plotting the DTFT




UNIVERSIDAD
TÉCNICA DE AMBATO

Plotting the DTFT




Plotting the DTFT




UNIVERSIDAD
TÉCNICA DE AMBATO

Plotting the DTFT




Plotting the DTFT



UNIVERSIDAD
TÉCNICA DE AMBATO

Plotting the DTFT



Remarks

- The DTFT thus maps an infinite length sequence $x[n]$ onto a function of real variable ω
- the DTFT is 2π -periodic and we write $X(\exp^{j\omega})$
- by convention DTFT is defined on the $[-\pi, \pi]$




Classifying signals in frequency

Three broad categories according to where most of the spectral energy resides:


- lowpass signals (also known as “baseband” signals)
- highpass signals
- bandpass signals




Lowpass Example



Highpass Example



Bandpass Example




Sinusoidal Modulation


$$\begin{aligned}\text{DTFT} \{x[n] \cos(\omega_c n)\} &= \text{DTFT} \left\{ \frac{1}{2} e^{j\omega_c n} x[n] + \frac{1}{2} e^{-j\omega_c n} x[n] \right\} \\ &= \frac{1}{2} \left[X(e^{j(\omega-\omega_c)}) + X(e^{j(\omega+\omega_c)}) \right]\end{aligned}$$

- ▶ usually $x[n]$ baseband
- ▶ ω_c is the *carrier* frequency


Example



Example




Example



UNIVERSIDAD
TÉCNICA DE AMBATO

Example



Example

- ▶ voice and music are lowpass signals
- ▶ radio channels are bandpass, in much higher frequencies
- ▶ modulation brings the baseband signal in the transmission band
- ▶ demodulation at the receiver brings it back

Demodulation


just multiply the received signal by the carrier again

$$y[n] = x[n] \cos(\omega_c n) \quad Y(e^{j\omega}) = \frac{1}{2} [X(e^{j(\omega-\omega_c)}) + X(e^{j(\omega+\omega_c)})]$$

$$\begin{aligned}\text{DTFT } \{y[n] \cdot 2 \cos(\omega_c n)\} &= Y(e^{j(\omega-\omega_c)}) + Y(e^{j(\omega+\omega_c)}) \\ &= \frac{1}{2} [X(e^{j(\omega-2\omega_c)}) + X(e^{j(\omega)}) + X(e^{j(\omega)}) + X(e^{j(\omega+2\omega_c)})] \\ &= X(e^{j(\omega)}) + \frac{1}{2} [X(e^{j(\omega-2\omega_c)}) + X(e^{j(\omega+2\omega_c)})]\end{aligned}$$

Example-Demodulation


DTFT $\{x[n]\}$



UNIVERSIDAD
TÉCNICA DE AMBATO

Example-Demodulation


$$\text{DTFT } \{y[n]\} = \text{DTFT } \{x[n] \cos \omega_c n\}$$



UNIVERSIDAD
TÉCNICA DE AMBATO

Example-Demodulation


$$\text{DTFT } \{y[n] \cos \omega_c n\}$$



UNIVERSIDAD
TÉCNICA DE AMBATO

Example-Demodulation


$$\text{DTFT } \{y[n] \cos \omega_c n\}$$



UNIVERSIDAD
TÉCNICA DE AMBATO

Example-Demodulation

$$\text{DTFT } \{y[n] \cos \omega_c n\}$$



FFT

