



Laplacian

25.3.30



$$\begin{aligned}\nabla^2 u_{0,0} &= \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)_{0,0} \\ &= \frac{1}{h^2} (u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1} - 4u_{0,0}) + O(h^2)\end{aligned}$$


25.3.31



$$\begin{aligned}\nabla^2 u_{0,0} &= \frac{1}{12h^2} [-60u_{0,0} + 16(u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1}) \\ &\quad - (u_{2,0} + u_{0,2} + u_{-2,0} + u_{0,-2})] + O(h^4)\end{aligned}$$


Biharmonic Operator

25.3.32



$$\begin{aligned}\nabla^4 u_{0,0} &= \left(\frac{\partial^4 u}{\partial x^4} + 2 \frac{\partial^4 u}{\partial x^2 \partial y^2} + \frac{\partial^4 u}{\partial y^4} \right)_{0,0} \\ &= \frac{1}{h^4} [20u_{0,0} - 8(u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1}) \\ &\quad + 2(u_{1,1} + u_{1,-1} + u_{-1,1} + u_{-1,-1}) \\ &\quad + (u_{0,2} + u_{2,0} + u_{-2,0} + u_{0,-2})] + O(h^2)\end{aligned}$$

25.3.33



$$\begin{aligned}\nabla^4 u_{0,0} &= \frac{1}{6h^4} [-(u_{0,3} + u_{0,-3} + u_{3,0} + u_{-3,0}) \\ &\quad + 14(u_{0,2} + u_{0,-2} + u_{2,0} + u_{-2,0}) \\ &\quad - 77(u_{0,1} + u_{0,-1} + u_{1,0} + u_{-1,0}) \\ &\quad + 184u_{0,0} + 20(u_{1,1} + u_{1,-1} + u_{-1,1} + u_{-1,-1}) \\ &\quad - (u_{1,2} + u_{2,1} + u_{1,-2} + u_{2,-1} + u_{-1,2} + u_{-2,1} \\ &\quad + u_{-1,-2} + u_{-2,-1})] + O(h^4)\end{aligned}$$

25.4. Integration

Trapezoidal Rule

25.4.1

$$\begin{aligned}\int_{x_0}^{x_1} f(x) dx &= \frac{h}{2} (f_0 + f_1) - \frac{1}{2} \int_{x_0}^{x_1} (t - x_0)(x_1 - t) f''(t) dt \\ &= \frac{h}{2} (f_0 + f_1) - \frac{h^3}{12} f''(\xi) \quad (x_0 < \xi < x_1)\end{aligned}$$

Extended Trapezoidal Rule

25.4.2

$$\begin{aligned}\int_{x_0}^{x_m} f(x) dx &= h \left[\frac{f_0}{2} + f_1 + \dots + f_{m-1} + \frac{f_m}{2} \right] \\ &\quad - \frac{mh^3}{12} f''(\xi)\end{aligned}$$

Error Term in Trapezoidal Formula for Periodic Functions

If $f(x)$ is periodic and has a continuous k^{th} derivative, and if the integral is taken over a period, then

$$25.4.3 \quad |\text{Error}| \leq \frac{\text{constant}}{m^k}$$

Modified Trapezoidal Rule

25.4.4

$$\begin{aligned}\int_{x_0}^{x_m} f(x) dx &= h \left[\frac{f_0}{2} + f_1 + \dots + f_{m-1} + \frac{f_m}{2} \right] \\ &\quad + \frac{h}{24} [-f_{-1} + f_1 + f_{m-1} - f_{m+1}] + \frac{11m}{720} h^5 f^{(4)}(\xi)\end{aligned}$$

Simpson's Rule**25.4.5**

$$\begin{aligned}\int_{x_0}^{x_2} f(x) dx &= \frac{h}{3} [f_0 + 4f_1 + f_2] \\ &+ \frac{1}{6} \int_{x_0}^{x_1} (x_0 - t)^2 (x_1 - t) f^{(3)}(t) dt \\ &+ \frac{1}{6} \int_{x_1}^{x_2} (x_2 - t)^2 (x_1 - t) f^{(3)}(t) dt \\ &= \frac{h}{3} [f_0 + 4f_1 + f_2] - \frac{h^5}{90} f^{(4)}(\xi)\end{aligned}$$

Extended Simpson's Rule**25.4.6**

$$\begin{aligned}\int_{x_0}^{x_{2n}} f(x) dx &= \frac{h}{3} [f_0 + 4(f_1 + f_3 + \dots + f_{2n-1}) \\ &+ 2(f_2 + f_4 + \dots + f_{2n-2}) + f_{2n}] - \frac{nh^5}{90} f^{(4)}(\xi)\end{aligned}$$

Euler-Maclaurin Summation Formula**25.4.7**

$$\begin{aligned}\int_{x_0}^{x_n} f(x) dx &= h \left[\frac{f_0}{2} + f_1 + f_2 + \dots + f_{n-1} + \frac{f_n}{2} \right] \\ &- \frac{B_2}{2!} h^2 (f'_n - f'_0) - \dots - \frac{B_{2k} h^{2k}}{(2k)!} [f^{(2k-1)}_n - f^{(2k-1)}_0] + R_{2k} \\ R_{2k} &= \frac{\theta n B_{2k+2} h^{2k+3}}{(2k+2)!} \max_{x_0 \leq x \leq x_n} |f^{(2k+2)}(x)|, \quad (-1 \leq \theta \leq 1)\end{aligned}$$

(For B_{2k} , Bernoulli numbers, see chapter 23.)

If $f^{(2k+2)}(x)$ and $f^{(2k+4)}(x)$ do not change sign for $x_0 < x < x_n$ then $|R_{2k}|$ is less than the first neglected term. If $f^{(2k+2)}(x)$ does not change sign for $x_0 < x < x_n$, $|R_{2k}|$ is less than twice the first neglected term.

Lagrange Formula**25.4.8**

$$\int_a^b f(x) dx = \sum_{i=0}^n (L_i^{(n)}(b) - L_i^{(n)}(a)) f_i + R_n$$

(See 25.2.1.)

25.4.9

$$L_i^{(n)}(x) = \frac{1}{\pi'_n(x_i)} \int_{x_0}^x \frac{\pi_n(t)}{t - x_i} dt = \int_{x_0}^x l_i(t) dt$$

$$\mathbf{25.4.10} \quad R_n = \frac{1}{(n+1)!} \int_a^b \pi_n(x) f^{(n+1)}(\xi(x)) dx$$

Equally Spaced Abscissas**25.4.11**

$$\int_{x_0}^{x_k} f(x) dx = \frac{1}{h^n} \sum_{i=0}^n f_i \frac{(-1)^{n-i}}{i!(n-i)!} \int_{x_0}^{x_k} \frac{\pi_n(x)}{x - x_i} dx + R_n$$

$$\mathbf{25.4.12} \quad \int_{x_m}^{x_{m+1}} f(x) dx = h \sum_{i=-\lfloor \frac{n-1}{2} \rfloor}^{\lfloor \frac{n}{2} \rfloor} A_i(m) f_i + R_n \quad *$$

(See Table 25.3 for $A_i(m)$.)**Newton-Cotes Formulas (Closed Type)**

(For Trapezoidal and Simpson's Rules see 25.4.1-25.4.6.)

25.4.13 (Simpson's $\frac{3}{8}$ rule)

$$\int_{x_0}^{x_3} f(x) dx = \frac{3h}{8} (f_0 + 3f_1 + 3f_2 + f_3) - \frac{3f^{(4)}(\xi)h^5}{80}$$

25.4.14 (Bode's rule)

$$\begin{aligned}\int_{x_0}^{x_4} f(x) dx &= \frac{2h}{45} (7f_0 + 32f_1 + 12f_2 \\ &+ 32f_3 + 7f_4) - \frac{8f^{(6)}(\xi)h^7}{945}\end{aligned}$$

25.4.15

$$\begin{aligned}\int_{x_0}^{x_5} f(x) dx &= \frac{5h}{288} (19f_0 + 75f_1 + 50f_2 + 50f_3 \\ &+ 75f_4 + 19f_5) - \frac{275f^{(6)}(\xi)h^7}{12096}\end{aligned}$$

25.4.16

$$\begin{aligned}\int_{x_0}^{x_6} f(x) dx &= \frac{h}{140} (41f_0 + 216f_1 + 27f_2 + 272f_3 \\ &+ 27f_4 + 216f_5 + 41f_6) - \frac{9f^{(8)}(\xi)h^9}{1400}\end{aligned}$$

25.4.17

$$\begin{aligned}\int_{x_0}^{x_7} f(x) dx &= \frac{7h}{17280} (751f_0 + 3577f_1 + 1323f_2 \\ &+ 2989f_3 + 2989f_4 + 1323f_5 + 3577f_6 \\ &+ 751f_7) - \frac{8183f^{(8)}(\xi)h^9}{518400}\end{aligned}$$

25.4.18

$$\begin{aligned}\int_{x_0}^{x_8} f(x) dx &= \frac{4h}{14175} (989f_0 + 5888f_1 - 928f_2 \\ &+ 10496f_3 - 4540f_4 + 10496f_5 - 928f_6 + 5888f_7 \\ &+ 989f_8) - \frac{2368}{467775} f^{(10)}(\xi)h^{11}\end{aligned}$$

25.4.19

$$\begin{aligned}\int_{x_0}^{x_9} f(x) dx &= \frac{9h}{89600} \{ 2857(f_0 + f_9) \\ &+ 15741(f_1 + f_8) + 1080(f_2 + f_7) + 19344(f_3 + f_6) \\ &+ 5778(f_4 + f_5) \} - \frac{173}{14620} f^{(10)}(\xi)h^{11}\end{aligned}$$

* See page II.

25.4.20

$$\int_{x_0}^{x_{10}} f(x) dx = \frac{5h}{299376} \{16067(f_0 + f_{10}) + 106300(f_1 + f_9) - 48525(f_2 + f_8) + 272400(f_3 + f_7) - 260550(f_4 + f_6) + 427368f_5\} - \frac{1346350}{326918592} f^{(12)}(\xi) h^{13}$$

Newton-Cotes Formulas (Open Type)

25.4.21

$$\int_{x_0}^{x_3} f(x) dx = \frac{3h}{2} (f_1 + f_2) + \frac{f^{(2)}(\xi) h^3}{4}$$

25.4.22

$$\int_{x_0}^{x_4} f(x) dx = \frac{4h}{3} (2f_1 - f_2 + 2f_3) + \frac{28f^{(4)}(\xi) h^5}{90}$$

25.4.23

$$\int_{x_0}^{x_5} f(x) dx = \frac{5h}{24} (11f_1 + f_2 + f_3 + 11f_4) + \frac{95f^{(4)}(\xi) h^5}{144}$$

25.4.24

$$\int_{x_0}^{x_6} f(x) dx = \frac{6h}{20} (11f_1 - 14f_2 + 26f_3 - 14f_4 + 11f_5) + \frac{41f^{(6)}(\xi) h^7}{140}$$

25.4.25


$$\int_{x_0}^{x_7} f(x) dx = \frac{7h}{1440} (611f_1 - 453f_2 + 562f_3 + 562f_4 - 453f_5 + 611f_6) + \frac{5257}{8640} f^{(6)}(\xi) h^7$$

25.4.26

$$\int_{x_0}^{x_8} f(x) dx = \frac{8h}{945} (460f_1 - 954f_2 + 2196f_3 - 2459f_4 + 2196f_5 - 954f_6 + 460f_7) + \frac{3956}{14175} f^{(8)}(\xi) h^9$$

Five Point Rule for Analytic Functions

25.4.27



$$\int_{z_0-h}^{z_0+h} f(z) dz = \frac{h}{15} \{24f(z_0) + 4[f(z_0+h) + f(z_0-h)] - [f(z_0+ih) + f(z_0-ih)]\} + R$$

$|R| \leq \frac{|h|^7}{1890} \max_{z \in S} |f^{(6)}(z)|$, S designates the square with vertices $z_0 + i^k h$ ($k=0, 1, 2, 3$); h can be complex.

Chebyshev's Equal Weight Integration Formula

$$25.4.28 \quad \int_{-1}^1 f(x) dx = \frac{2}{n} \sum_{i=1}^n f(x_i) + R_n$$

Abscissas: x_i is the i^{th} zero of the polynomial part of

$$x^n \exp \left[\frac{-n}{2 \cdot 3x^2} - \frac{n}{4 \cdot 5x^3} - \frac{n}{6 \cdot 7x^4} - \dots \right]$$

(See Table 25.5 for x_i .)

For $n=8$ and $n \geq 10$ some of the zeros are complex.

Remainder:

$$R_n = \int_{-1}^{+1} \frac{x^{n+1}}{(n+1)!} f^{(n+1)}(\xi) dx - \frac{2}{n(n+1)!} \sum_{i=1}^n x_i^{n+1} f^{(n+1)}(\xi_i)$$

where $\xi = \xi(x)$ satisfies $0 \leq \xi \leq x$ and $0 \leq \xi_i \leq x_i$

$$(i=1, \dots, n)$$

Integration Formulas of Gaussian Type

(For Orthogonal Polynomials see chapter 22)

Gauss' Formula

$$25.4.29 \quad \int_{-1}^1 f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Legendre polynomials $P_n(x)$, $P_n(1)=1$

Abscissas: x_i is the i^{th} zero of $P_n(x)$

Weights: $w_i = 2/(1-x_i^2) [P'_n(x_i)]^2$
(See Table 25.4 for x_i and w_i .)

$$R_n = \frac{2^{2n+1} (n!)^4}{(2n+1) [(2n)!]^3} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

Gauss' Formula, Arbitrary Interval

$$25.4.30 \quad \int_a^b f(y) dy = \frac{b-a}{2} \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \left(\frac{b-a}{2} \right) x_i + \left(\frac{b+a}{2} \right)$$

Related orthogonal polynomials: $P_n(x)$, $P_n(1)=1$

Abscissas: x_i is the i^{th} zero of $P_n(x)$

* Weights: $w_i = 2/(1-x_i^2) [P'_n(x_i)]^2$

$$* R_n = \frac{(b-a)^{2n+1} (n!)^4}{(2n+1) [(2n)!]^3} f^{(2n)}(\xi)$$

Radau's Integration Formula

25.4.31

$$\int_{-1}^1 f(x) dx = \frac{2}{n^2} f_{-1} + \sum_{i=1}^{n-1} w_i f(x_i) + R_n$$

Related polynomials:

$$\frac{P_{n-1}(x) + P_n(x)}{x+1}$$

Abscissas: x_i is the i^{th} zero of

$$\frac{P_{n-1}(x) + P_n(x)}{x+1}$$

Weights:

$$w_i = \frac{1}{n^2} \frac{1-x_i}{[P_{n-1}(x_i)]^2} = \frac{1}{1-x_i} \frac{1}{[P'_{n-1}(x_i)]^2}$$

Remainder:

$$R_n = \frac{2^{2n-1} \cdot n}{[(2n-1)!]^3} [(n-1)!]^4 f^{(2n-1)}(\xi) \quad (-1 < \xi < 1)$$

Lobatto's Integration Formula

25.4.32

$$\int_{-1}^1 f(x) dx = \frac{2}{n(n-1)} [f(1) + f(-1)] + \sum_{i=2}^{n-1} w_i f(x_i) + R_n$$

Related polynomials: $P'_{n-1}(x)$

Abscissas: x_i is the $(i-1)^{\text{st}}$ zero of $P'_{n-1}(x)$

Weights:

$$w_i = \frac{2}{n(n-1)[P'_{n-1}(x_i)]^2} \quad (x_i \neq \pm 1)$$

(See Table 25.6 for x_i and w_i .)

Remainder:

$$R_n = \frac{-n(n-1)^3 2^{2n-1} [(n-2)!]^4}{(2n-1) [(2n-2)!]^3} f^{(2n-2)}(\xi) \quad (-1 < \xi < 1)$$

*See page II.

$$25.4.33 \quad \int_0^1 x^k f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$q_n(x) = \sqrt{k+2n+1} P_n^{(k,0)}(1-2x)$$

(For the Jacobi polynomials $P_n^{(k,0)}$ see chapter 22.)

Abscissas:

x_i is the i^{th} zero of $q_n(x)$

Weights:

$$w_i = \left\{ \sum_{j=0}^{n-1} [q_j(x_i)]^2 \right\}^{-1}$$

(See Table 25.8 for x_i and w_i .)

Remainder:

$$R_n = \frac{f^{(2n)}(\xi)}{(k+2n+1)(2n)!} \left[\frac{n!(k+n)!}{(k+2n)!} \right]^2 \quad (0 < \xi < 1)$$

25.4.34

$$\int_0^1 f(x) \sqrt{1-x} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{1-x}} P_{2n+1}(\sqrt{1-x}), P_{2n+1}(1)=1$$

Abscissas: $x_i = 1 - \xi_i^2$ where ξ_i is the i^{th} positive zero of $P_{2n+1}(x)$.

Weights: $w_i = 2\xi_i^2 w_i^{(2n+1)}$ where $w_i^{(2n+1)}$ are the Gaussian weights of order $2n+1$.

Remainder:

$$R_n = \frac{2^{4n+3} [(2n+1)!]^4}{(2n)!(4n+3)[(4n+2)!]^2} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

25.4.35

$$\int_a^b f(y) \sqrt{b-y} dy = (b-a)^{3/2} \sum_{i=1}^n w_i f(y_i)$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{1-x}} P_{2n+1}(\sqrt{1-x}), P_{2n+1}(1)=1$$

Abscissas: $x_i = 1 - \xi_i^2$ where ξ_i is the i^{th} positive zero of $P_{2n+1}(x)$.

Weights: $w_i = 2\xi_i^2 w_i^{(2n+1)}$ where $w_i^{(2n+1)}$ are the Gaussian weights of order $2n+1$.

$$25.4.36 \quad \int_0^1 \frac{f(x)}{\sqrt{1-x}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$P_{2n}(\sqrt{1-x}), P_{2n}(1)=1$$

Abscissas: $x_i = 1 - \xi_i^2$ where ξ_i is the i^{th} positive zero of $P_{2n}(x)$.

Weights: $w_i = 2w_i^{(2n)}$, $w_i^{(2n)}$ are the Gaussian weights of order $2n$.

Remainder:

$$R_n = \frac{2^{4n+1}}{4n+1} \frac{[(2n)!]^3}{[(4n)!]^2} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

$$25.4.37 \quad \int_a^b \frac{f(y)}{\sqrt{b-y}} dy = \sqrt{b-a} \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$P_{2n}(\sqrt{1-x}), P_{2n}(1)=1$$

Abscissas:

$x_i = 1 - \xi_i^2$ where ξ_i is the i^{th} positive zero of $P_{2n}(x)$.

Weights: $w_i = 2w_i^{(2n)}$, $w_i^{(2n)}$ are the Gaussian weights of order $2n$.

$$25.4.38 \quad \int_{-1}^{+1} \frac{f(x)}{\sqrt{1-x^2}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Chebyshev Polynomials of First Kind

$$T_n(x), T_n(1) = \frac{1}{2^{n-1}}$$

Abscissas:

$$x_i = \cos \frac{(2i-1)\pi}{2n}$$

Weights:

$$w_i = \frac{\pi}{n}$$

Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{2n-1}} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

25.4.39

$$\int_a^b \frac{f(y) dy}{\sqrt{(y-a)(b-y)}} = \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \frac{b+a}{2} + \frac{b-a}{2} x_i$$

Related orthogonal polynomials:

$$T_n(x), T_n(1) = \frac{1}{2^{n-1}}$$

Abscissas:

$$x_i = \cos \frac{(2i-1)\pi}{2n}$$

Weights:

$$w_i = \frac{\pi}{n}$$

25.4.40

$$\int_{-1}^{+1} f(x) \sqrt{1-x^2} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Chebyshev Polynomials of Second Kind

$$U_n(x) = \frac{\sin [(n+1) \arccos x]}{\sin (\arccos x)} \quad *$$

Abscissas:

$$x_i = \cos \frac{i}{n+1} \pi \quad *$$

Weights:

$$w_i = \frac{\pi}{n+1} \sin^2 \frac{i}{n+1} \pi \quad *$$

Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{2n+1}} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

25.4.41

$$\int_a^b \sqrt{(y-a)(b-y)} f(y) dy = \left(\frac{b-a}{2}\right)^2 \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \frac{b+a}{2} + \frac{b-a}{2} x_i$$

Related orthogonal polynomials:

$$U_n(x) = \frac{\sin [(n+1) \arccos x]}{\sin (\arccos x)} \quad *$$

Abscissas:

$$x_i = \cos \frac{i}{n+1} \pi \quad *$$

Weights:

$$w_i = \frac{\pi}{n+1} \sin^2 \frac{i}{n+1} \pi \quad *$$

$$25.4.42 \quad \int_0^1 f(x) \sqrt{\frac{x}{1-x}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{x}} T_{2n+1}(\sqrt{x})$$

Abscissas:

$$x_i = \cos^2 \frac{2i-1}{2n+1} \cdot \frac{\pi}{2}$$

Weights:

$$w_i = \frac{2\pi}{2n+1} x_i$$

Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{4n+1}} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

25.4.43

$$\int_a^b f(x) \sqrt{\frac{x-a}{b-x}} dx = (b-a) \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{x}} T_{2n+1}(\sqrt{x})$$

Abscissas:

$$x_i = \cos^2 \frac{2i-1}{2n+1} \cdot \frac{\pi}{2}$$

Weights:

$$w_i = \frac{2\pi}{2n+1} x_i$$

25.4.44
$$\int_0^1 \ln x f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: polynomials orthogonal with respect to the weight function $-\ln x$

Abscissas: See Table 25.7

Weights: See Table 25.7

25.4.45

$$\int_0^\infty e^{-x} f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Laguerre polynomials $L_n(x)$.

Abscissas: x_i is the i^{th} zero of $L_n(x)$

Weights:

$$* \quad w_i = \frac{x_i}{(n+1)^2 [L_{n+1}(x_i)]^2}$$

(See Table 25.9 for x_i and w_i .)

Remainder:

$$R_n = \frac{(n!)^2}{(2n)!} f^{(2n)}(\xi) \quad (0 < \xi < \infty)$$

25.4.46

$$\int_{-\infty}^\infty e^{-x^2} f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Hermite polynomials $H_n(x)$.

Abscissas: x_i is the i^{th} zero of $H_n(x)$

Weights:

$$\frac{2^{n-1} n! \sqrt{\pi}}{n^2 [H_{n-1}(x_i)]^2}$$

(See Table 25.10 for x_i and w_i .)

*See page 11.

Remainder:

$$R_n = \frac{n! \sqrt{\pi}}{2^n (2n)!} f^{(2n)}(\xi) \quad (-\infty < \xi < \infty)$$

Filon's Integration Formula³

25.4.47

$$\int_{x_0}^{x_n} f(x) \cos tx dx = h \left[\alpha(th) (f_{2n} \sin tx_{2n} - f_0 \sin tx_0) + \beta(th) \cdot C_{2n} + \gamma(th) \cdot C_{2n-1} + \frac{2}{45} th^4 S'_{2n-1} \right] - R_n$$

25.4.48

$$C_{2n} = \sum_{i=0}^n f_{2i} \cos(tx_{2i}) - \frac{1}{2} [f_{2n} \cos tx_{2n} + f_0 \cos tx_0]$$

25.4.49

$$C_{2n-1} = \sum_{i=1}^n f_{2i-1} \cos tx_{2i-1}$$

25.4.50

$$S'_{2n-1} = \sum_{i=1}^n f_{2i-1}^{(3)} \sin tx_{2i-1}$$

25.4.51

$$R_n = \frac{1}{90} nh^5 f^{(4)}(\xi) + O(th^7)$$

25.4.52

$$\alpha(\theta) = \frac{1}{\theta} + \frac{\sin 2\theta}{2\theta^2} - \frac{2 \sin^2 \theta}{\theta^3}$$

$$\beta(\theta) = 2 \left(\frac{1 + \cos^2 \theta}{\theta^2} - \frac{\sin 2\theta}{\theta^3} \right)$$

$$\gamma(\theta) = 4 \left(\frac{\sin \theta}{\theta^3} - \frac{\cos \theta}{\theta^2} \right)$$

For small θ we have

25.4.53

$$\alpha = \frac{2\theta^3}{45} - \frac{2\theta^5}{315} + \frac{2\theta^7}{4725} - \dots$$

$$\beta = \frac{2}{3} + \frac{2\theta^2}{15} - \frac{4\theta^4}{105} + \frac{2\theta^6}{567} - \dots$$

$$\gamma = \frac{4}{3} - \frac{2\theta^2}{15} + \frac{\theta^4}{210} - \frac{\theta^6}{11340} + \dots$$

25.4.54

$$\int_{x_0}^{x_{2n}} f(x) \sin tx dx = h \left[\alpha(th) (f_0 \cos tx_0 - f_{2n} \cos tx_{2n}) + \beta S_{2n} + \gamma S_{2n-1} + \frac{2}{45} th^4 C'_{2n-1} \right] - R_n$$

25.4.55

$$S_{2n} = \sum_{i=0}^n f_{2i} \sin(tx_{2i}) - \frac{1}{2} [f_{2n} \sin(tx_{2n}) + f_0 \sin(tx_0)]$$

³ For certain difficulties associated with this formula, see the article by J. W. Tukey, p. 400, "On Numerical Approximation," Ed. R. E. Langer, Madison, 1959.