

My Test Manuscript

Marc Pérez

Hier kommt das Abstract

Preface

Abstract

(Knuth 1984)

Introduction

Complexity of Phosphorous

Phosphorous displays a wide range of behaviours in soils, in places where organic, mineral and aquaeous phases interface. In phases that contain oxygen Phosphorous is almost exclusively present as several derivates of Orthophosphate PO_4^{3-} . It can be found as organic molecules as anhydric- and ester-groups, being needed by all known species as a constituent of DNA and energy transfer-processes. It can be present as anorganic Phosphate either as mono-orthophosphate PO_4^{3-} or poly-orthophosphate $HO - (PO_2)_n - OH$, where it can strongly interact with water, forming, depending on pH HPO_4^{2-} or $H_2PO_4^-$. The dissolved species of phosphate are subject to adsorption to clay- and oxide-surfaces of the solid soil-phase, they also form fallout-products such as Apatite, Vivianite ect. with the present metal-cations in the solution. While the solubility constant of most phosphate-salts are comparably low (Wert eingeben), meaning that the fallout and formation of minerals happens at low chemical activities of phosphate, phosphate often is leached from soil-surface-layers, heavily reducing the efficacy of P-fertilization and presenting a disturbance to P-limited ecosystems. Those phenomena, many of them being physico-chemically controlled, are influenced by parameters such as pH, ionic-strength, clay-content, specific-surface of the soild phase, amorphous $Fe(OH)_3$ -content amorphous $Al(OH)_3$ -content, in short the phenomena depend heavily on the composition,

distribution and geometry of the soil. Those properties are considered to be stable respectively long-term properties of a soil, when looked at it with the interest of modeling the transport processes of Phosphate in soils. Factors such as water-content, temperature, vegetation and precipitation are factors that temporally can vary fast and to a certain degree unpredictably. Organic forms of phosphates, prominently DNA or oligo-nucleotides and phytate are also subject to physico-chemical reactions, mainly decomposition, but are foremost controlled in their presence by enzymatic processes, where i.e. plants form phytates in seeds to provide the embryo a compact and specific reserve of phosphate, but many bacteria possess via Phytases the ability to hydrolise phytate and use it for their own means. To assess and cover those phenomena, models, dynamically describing the motion of Phosphorous in soils, differentiate several pools of Phosphorous, most prominently the organic-P, dissolved-P, adsorbed-P, mineral-P, where the difference in temporal behaviour, such as the mean-reside-time can lead to a differentiation between labile-P, semi-labile-P and so on.

Plants as Phosphate sinks

When a soil is used agronomically, P-sinks such as leaching and plant P-uptake

Struktur

Warum ist die Arbeit wichtig

P ist endlich, Umweltprobleme ## P ist sehr komplex

Siehe oben ## Wie wird bisher P-Ernährung angegangen

GRUD

$$\frac{A}{B}$$

Warum Bodentest


Warum kinetischer Bodentest

Ramona Test

djfqrigjriodjkroidgjroigjreiojgeriogjrigjrijqrijriohriherioügrgjriohroighruih

Method

[1] 19



Source: [Article Notebook](#)

Results

Discussion

Conclusion

Acknowledgments

Legal Disclosure

References

Knuth, Donald E. 1984. "Literate Programming." *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.

Appendix

Supplements