


Sinusoidal ICA component



90% Confidence intervals:

$$\text{Par. estimators} = \left[q_{50\%} \right]_{q_{05\%}-q_{50\%}}^{q_{95\%}-q_{50\%}}$$

$$\text{Amplitude } A = 1.267_{-0.022}^{+0.022}$$

$$\text{Period } T = 3.318_{-0.014}^{+0.014}$$

$$\text{Phase } \frac{\phi}{\pi} = 1.012_{-0.011}^{+0.012}$$


$$\text{Noise } \sigma = 0.417_{-0.011}^{+0.011}$$

True value of Period $T = 3.30$

1000 random MCMC parameter sets
 (A, T, ϕ, σ)

Maxima A Posteriori:
 $(A = 1.27, T = 3.32, \phi = 3.18, \sigma = 0.42)$

Sawtooth ICA component



90% Confidence intervals:

$$\text{Par. estimators} = \left[q_{50\%} \right]_{q_{05\%}-q_{50\%}}^{q_{95\%}-q_{50\%}}$$

$$\text{Amplitude } A = 1.704_{-0.011}^{+0.011}$$

$$\text{Period } T = 1.0001_{-0.0001}^{+0.0002}$$

$$\text{Phase } \frac{\phi}{\pi} = 0.002_{-0.001}^{+0.002}$$

$$\text{Noise } \sigma = 0.177_{-0.005}^{+0.005}$$

True value of Period $T = 1.00$

Maxima A Posteriori:
 $(A = 1.70, T = 1.00, \phi = 0.00, \sigma = 0.18)$