

Linear averaging and flow dynamics

Giacomo Como, DISMA, Politecnico di Torino

Fabio Fagnani, DISMA, Politecnico di Torino

Linear averaging and flow dynamics

French-De Groot learning model

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ weighted directed graph representing a social network



- ▶ $(i, j) \in \mathcal{E}$ if i is influenced by j
- ▶ W_{ij} *strength* of this influence
- ▶ $P_{ij} = w_i^{-1} W_{ij}$ ($w_i = \sum_k W_{ik}$)
normalized weight matrix

French-De Groot learning model

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ weighted directed graph representing a social network



- ▶ $(i, j) \in \mathcal{E}$ if i is influenced by j
- ▶ W_{ij} *strength* of this influence
- ▶ $P_{ij} = w_i^{-1} W_{ij}$ ($w_i = \sum_k W_{ik}$)
normalized weight matrix

- ▶ Each agent $i \in \mathcal{V}$ has an initial opinion on some fact or event $x_i(0) \in \mathbb{R}$.
- ▶ Learning model: agents modify their opinion in time averaging the opinions of their neighbors:

$$x_i(t+1) = \sum_j P_{ij} x_j(t)$$

Linear averaging dynamics

French-De Groot learning model

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ weighted directed graph representing a social network



- ▶ $(i, j) \in \mathcal{E}$ if i is influenced by j
- ▶ W_{ij} *strength* of this influence
- ▶ $P_{ij} = w_i^{-1} W_{ij}$ ($w_i = \sum_k W_{ik}$)
normalized weight matrix

- ▶ Each agent $i \in \mathcal{V}$ has an initial opinion on some fact or event $x_i(0) \in \mathbb{R}$.
- ▶ Learning model: agents modify their opinion in time averaging the opinions of their neighbors:

$$x_i(t+1) = \sum_j P_{ij} x_j(t)$$

Linear averaging dynamics

- ▶ Compact notation $x(t) \in \mathbb{R}^n$:

$$x(t+1) = P_X(t)$$

$$x(t) = P^t x(0)$$

- Behavior when $t \rightarrow +\infty$?

Giacomo Como, DISMA, Politecnico di Torino Fabio Fagnani, DISMA, Politecnico di Torino

A little digression: dynamical systems

- ▶ A *dynamical system*: Ω set, $f : \Omega \rightarrow \Omega$ map,
- ▶ $x(0) \in \Omega$ *initial condition*
- ▶ *Evolution*:

$$x(0) \mapsto x(1) = f(x(0)) \mapsto x(2) = f(x(1)) \dots$$

More formally, $x(t)$ is the sequence defined recursively by

$$x(t+1) = f(x(t))$$

A little digression: dynamical systems

- ▶ A *dynamical system*: Ω set, $f : \Omega \rightarrow \Omega$ map,
- ▶ $x(0) \in \Omega$ *initial condition*
- ▶ *Evolution*:

$$x(0) \mapsto x(1) = f(x(0)) \mapsto x(2) = f(x(1)) \dots$$

More formally, $x(t)$ is the sequence defined recursively by

$$x(t+1) = f(x(t))$$

- x_0 *equilibrium* if $f(x_0) = x_0$;

$$x(0) = x_0 \Rightarrow x(t) = x_0 \quad \forall t$$

A little digression: dynamical systems

- ▶ A *dynamical system*: Ω set, $f : \Omega \rightarrow \Omega$ map,
- ▶ $x(0) \in \Omega$ *initial condition*
- ▶ *Evolution*: $x(t)$ is the sequence defined recursively by

$$x(t+1) = f(x(t))$$

- ▶ x_0 *equilibrium* if $f(x_0) = x_0$;

$$x(0) = x_0 \Rightarrow x(t) = x_0 \quad \forall t$$

- ▶ Ω metric space (e.g. $\Omega = \mathbb{R}^n$), f continuous:

$$\text{If } \lim_{t \rightarrow +\infty} x(t) = \bar{x} \Rightarrow f(\bar{x}) = \bar{x} \text{ (equilibrium)}$$

Proof:

A little digression: dynamical systems

- ▶ A *dynamical system*: Ω set, $f : \Omega \rightarrow \Omega$ map,
- ▶ $x(0) \in \Omega$ *initial condition*
- ▶ *Evolution*: $x(t)$ is the sequence defined recursively by

$$x(t+1) = f(x(t))$$

- ▶ x_0 *equilibrium* if $f(x_0) = x_0$;

$$x(0) = x_0 \Rightarrow x(t) = x_0 \quad \forall t$$

- ▶ Ω metric space (e.g. $\Omega = \mathbb{R}^n$), f continuous:

If $\lim_{t \rightarrow +\infty} x(t) = \bar{x} \Rightarrow f(\bar{x}) = \bar{x}$ (equilibrium)

Proof: $x(t) \rightarrow \bar{x} \Rightarrow x(t+1) = f(x(t)) \rightarrow f(\bar{x})$

But $x(t+1) \rightarrow \bar{x}$. Hence, $f(\bar{x}) = \bar{x}$



Recap of PF theory for stochastic matrices

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$, $P = D^{-1}W$ normalized weight matrix

Theorem (Spectral properties of stochastic matrices)

- ▶ Dominant eigenvalue $\lambda_P = 1$, $P\mathbb{1} = \mathbb{1}$,
- ▶ Invariant distributions $\{\pi \in \mathbb{R}_+^n, \mathbb{1}'\pi = 1, P'\pi = \pi\}$ form a simplex in $\mathbb{R}^{\mathcal{V}}$ with $s_{\mathcal{G}}$ vertices.
- ▶ For every sink component with nodes \mathcal{W} , there exists an invariant distribution π such that $\pi_i > 0$ if and only if $i \in \mathcal{W}$.
- ▶ The invariant distribution is unique if and only if $s_{\mathcal{G}} = 1$.
- ▶ If \mathcal{G} is strongly connected, then 1 is simple and $\pi_i > 0$ for all i ;
- ▶ If \mathcal{G} is strongly connected and aperiodic, then every eigenvalue $\mu \neq 1$ is s.t. $|\mu| < 1$.

Analysis of the French-De Groot model

- ▶ Dynamical system with $\Omega = \mathbb{R}^n$, $f(x) = Px$
- ▶ Evolution from the initial condition $x(0)$: $x(t) = P^t x(0)$
- ▶ Equilibria: $Px_0 = x_0$, right eigenvectors of P relative to the dominant eigenvalue 1
- ▶ *Consensus vectors* $\alpha \mathbf{1}$ are always equilibria
- ▶ If $s_{\mathcal{G}} = 1$, the only equilibria are the consensus vectors.

Analysis of the French-De Groot model

- ▶ Dynamical system with $\Omega = \mathbb{R}^n$, $f(x) = Px$
- ▶ Evolution from the initial condition $x(0)$: $x(t) = P^t x(0)$
- ▶ Equilibria: $Px_0 = x_0$, right eigenvectors of P relative to the dominant eigenvalue 1
- ▶ *Consensus vectors* $\alpha \mathbb{1}$ are always equilibria
- ▶ If $s_G = 1$, the only equilibria are the consensus vectors.
- ▶ **IMPORTANT:** for any invariant distribution π

$$\pi'x(t+1) = \pi'Px(t) = \pi'x(t) \quad \forall t \Rightarrow \pi'x(t) = \pi'x(0) \quad \forall t$$

The quantity $\pi'x(t)$ is a *motion invariant*.

Analysis of the French-De Groot model

$$x(t) = P^t x(0)$$

Suppose that for a given $x(0)$, $x(t) \rightarrow \bar{x}$ for $t \rightarrow +\infty$

Then,

Analysis of the French-De Groot model

$$x(t) = P^t x(0)$$

Suppose that for a given $x(0)$, $x(t) \rightarrow \bar{x}$ for $t \rightarrow +\infty$

Then,

- ▶ \bar{x} is an equilibrium.
- ▶ $\pi' \bar{x} = \pi' x(0)$ for every invariant distribution π

Analysis of the French-De Groot model

$$x(t) = P^t x(0)$$

Suppose that for a given $x(0)$, $x(t) \rightarrow \bar{x}$ for $t \rightarrow +\infty$

Then,

- ▶ \bar{x} is an equilibrium.
- ▶ $\pi' \bar{x} = \pi' x(0)$ for every invariant distribution π

Proof: $\pi' x(t) \rightarrow \pi' \bar{x}$, $\pi' x(t) = \pi' x(0) \forall t$. Hence, $\pi' \bar{x} = \pi' x(0)$.

Analysis of the French-De Groot model

$$x(t) = P^t x(0)$$

Suppose that for a given $x(0)$, $x(t) \rightarrow \bar{x}$ for $t \rightarrow +\infty$

Then,

- ▶ \bar{x} is an equilibrium.
- ▶ $\pi' \bar{x} = \pi' x(0)$ for every invariant distribution π

Proof: $\pi' x(t) \rightarrow \pi' \bar{x}$, $\pi' x(t) = \pi' x(0) \forall t$. Hence, $\pi' \bar{x} = \pi' x(0)$.

- ▶ When $s_G = 1$, $x(t) \rightarrow \bar{x} = \alpha \mathbb{1}$ and

$$\alpha = \pi' \bar{x} = \pi' x(0) = \sum_{k \in \mathcal{V}} \pi_k x_k(0)$$

where π is the only invariant distribution of P


Analysis of the French-De Groot model

$$x(t) = P^t x(0)$$

When does it converge?

A counterexample


$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ 2-line simple graph:



$$W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

A counterexample

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ 2-line simple graph:




$$W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$x(0) = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad x(1) = Px(0) = \begin{pmatrix} \beta \\ \alpha \end{pmatrix}$$

A counterexample

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ 2-line simple graph:



$$W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$


$$x(0) = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad x(1) = Px(0) = \begin{pmatrix} \beta \\ \alpha \end{pmatrix}$$

$$t \text{ even } x(t) = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad t \text{ odd } x(t) = \begin{pmatrix} \beta \\ \alpha \end{pmatrix}$$

If $\alpha \neq \beta$, no convergence!

A counterexample

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ 2-line simple graph:



$$W = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$x(0) = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad x(1) = Px(0) = \begin{pmatrix} \beta \\ \alpha \end{pmatrix}$$

$$t \text{ even } x(t) = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, \quad t \text{ odd } x(t) = \begin{pmatrix} \beta \\ \alpha \end{pmatrix}$$

If $\alpha \neq \beta$, no convergence!

The graph above is not aperiodic....

Asymptotics of French-De Groot learning model

Theorem

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

$$\lim_{t \rightarrow +\infty} x(t) = \alpha \mathbb{1}$$

with
$$\alpha = \pi' x(0) = \sum_k \pi_k x_k(0) \quad \forall i$$

where $\pi = P' \pi$ is the unique invariant distribution of P .

Asymptotics of French-De Groot learning model

Theorem

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

$$\lim_{t \rightarrow +\infty} x(t) = \alpha \mathbb{1}$$


with
$$\alpha = \pi' x(0) = \sum_k \pi_k x_k(0) \quad \forall i$$

where $\pi = P' \pi$ is the unique invariant distribution of P .

- ▶ All opinions converge to a common value: **CONSENSUS**
- ▶ The consensus value $\pi' x(0)$ is a convex combination of the original opinions weighted by the *invariant distribution centralities* of the various agents.
- ▶ The invariant distribution centrality is a measure of the **social power** of an agent in the French-De Groot learning process.

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




$$P = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 \\ 1/4 & 0 & 1/4 & 1/4 & 1/4 \\ 1/3 & 1/3 & 0 & 1/3 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$P = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 \\ 1/4 & 0 & 1/4 & 1/4 & 1/4 \\ 1/3 & 1/3 & 0 & 1/3 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$


$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

- \mathcal{G} is strongly connected and aperiodic
- $\lim_{t \rightarrow +\infty} x_i(t) = \pi' x(0)$ for every i

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$P = \begin{pmatrix} 0 & 1/2 & 1/2 & 0 & 0 \\ 1/4 & 0 & 1/4 & 1/4 & 1/4 \\ 1/3 & 1/3 & 0 & 1/3 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$


$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

- ▶ \mathcal{G} is strongly connected and aperiodic
- ▶ $\lim_{t \rightarrow +\infty} x_i(t) = \pi'_i x(0)$ for every i
- ▶ \mathcal{G} is undirected, unweighted $\Rightarrow \pi = (1/6, 1/3, 1/4, 1/6, 1/12)$
- ▶ $\pi'_i x(0) = 1/3 + 1/2 + 5/6 + 1/4 = 23/12$

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




$$P = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1/3 & 1/3 & 1/3 \\ 1/2 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$P = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1/3 & 1/3 & 1/3 \\ 1/2 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$


$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

- ▶ \mathcal{G} str. conn. and aperiodic
- ▶ $\lim_{t \rightarrow +\infty} x_i(t) = \pi'_i x(0)$ for every i
- ▶ Computation of π :

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$P = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1/3 & 1/3 & 1/3 \\ 1/2 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

- ▶ \mathcal{G} str. conn. and aperiodic
- ▶ $\lim_{t \rightarrow +\infty} x_i(t) = \pi'_i x(0)$ for every i
- ▶ Computation of π :

$$\begin{cases} \pi_1 &= \pi_3/2 \\ \pi_2 &= \pi_1 + \pi_3/2 + \pi_5 \\ \pi_3 &= \pi_2/3 + \pi_4 \\ \pi_5 &= \pi_4 = \pi_2/3 \end{cases}$$

$$\pi = (1/8, 3/8, 1/4, 1/8, 1/8)$$
- ▶ $\pi'_i x(0) =$

$$1/2 + 1/2 + 5/8 + 3/8 = 2$$

Wisdom of crowds and wise societies

- ▶ Social network $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ str. connected aperiodic
- ▶ The initial opinion is a noisy measurement of a true variable μ :

$$x_i(0) = \mu + N_i, \quad N_i \text{ independent r.v. } \mathbb{E}[N_i] = 0 \text{ Var}(N_i) = \sigma^2$$

- ▶ French-De Groot learning model $x(t+1) = Px(t)$

Wisdom of crowds and wise societies

- ▶ Social network $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ str. connected aperiodic
- ▶ The initial opinion is a noisy measurement of a true variable μ :

$$x_i(0) = \mu + N_i, \quad N_i \text{ independent r.v. } \mathbb{E}[N_i] = 0 \text{ Var}(N_i) = \sigma^2$$

- ▶ French-De Groot learning model $x(t+1) = Px(t)$
- ▶ $x_i(t) \rightarrow \pi'x(0) = \sum_k \pi_k(\mu + N_k) = \mu + \sum_k \pi_k N_k$

Wisdom of crowds and wise societies

- ▶ Social network $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ str. connected aperiodic
- ▶ The initial opinion is a noisy measurement of a true variable μ :

$$x_i(0) = \mu + N_i, \quad N_i \text{ independent r.v. } \mathbb{E}[N_i] = 0 \text{ Var}(N_i) = \sigma^2$$
- ▶ French-De Groot learning model $x(t+1) = Px(t)$
- ▶ $x_i(t) \rightarrow \pi'_i x(0) = \sum_k \pi_k (\mu + N_k) = \mu + \sum_k \pi_k N_k$
- ▶ $\text{Var}(\sum \pi_k N_k) = \sigma^2 \sum \pi_k^2$
 \mathcal{G} strongly connected, $\pi_i > 0$ for all i , then, $\sum \pi_k^2 < \sum \pi_k = 1$.

Wisdom of crowds and wise societies

- ▶ Social network $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ str. connected aperiodic
- ▶ The initial opinion is a noisy measurement of a true variable μ :

$$x_i(0) = \mu + N_i, \quad N_i \text{ independent r.v. } \mathbb{E}[N_i] = 0 \text{ Var}(N_i) = \sigma^2$$

- ▶ French-De Groot learning model $x(t+1) = Px(t)$
- ▶ $x_i(t) \rightarrow \pi'x(0) = \sum_k \pi_k(\mu + N_k) = \mu + \sum_k \pi_k N_k$
- ▶ $\text{Var}(\sum \pi_k N_k) = \sigma^2 \sum \pi_k^2$
 \mathcal{G} strongly connected, $\pi_i > 0$ for all i , then, $\sum \pi_k^2 < \sum \pi_k = 1$.

$$\text{Var}(\sum \pi_k N_k) < \sigma^2 \quad \text{Crowd is wiser than a single!}$$

Wisdom of crowds and wise societies

- ▶ Social network $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ str. connected aperiodic
- ▶ The initial opinion is a noisy measurement of a true variable μ :

$$x_i(0) = \mu + N_i, \quad N_i \text{ independent r.v. } \mathbb{E}[N_i] = 0 \text{ Var}(N_i) = \sigma^2$$

- ▶ French-De Groot learning model $x(t+1) = Px(t)$

- ▶ $x_i(t) \rightarrow \pi'x(0) = \sum_k \pi_k(\mu + N_k) = \mu + \sum_k \pi_k N_k$

- ▶ $\text{Var}(\sum \pi_k N_k) = \sigma^2 \sum \pi_k^2$
 \mathcal{G} strongly connected, $\pi_i > 0$ for all i , then, $\sum \pi_k^2 < \sum \pi_k = 1$.

$$\text{Var}(\sum \pi_k N_k) < \sigma^2 \quad \text{Crowd is wiser than a single!}$$

- ▶ *Asymptotic wisdom*: $\lim_{n \rightarrow +\infty} \pi'x(0) = \mu \Leftrightarrow \lim_{n \rightarrow +\infty} \max_k \pi_k = 0$
 (Golub and Jackson, 2010)

The many applications of averaging dynamics

- ▶ A simple model for opinion fusion, social power, and consensus formation
- ▶ The basis of many distributed algorithms
 - ▶ Decentralized computation in sensor networks
 - ▶ Load balancing in computer networks
 - ▶ Clock synchronization
 - ▶ Relative localization
 - ▶ Coordination dynamics of robot networks.

Asymptotics of French-De Groot learning model

Theorem

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

$$\lim_{t \rightarrow +\infty} x(t) = \alpha \mathbb{1}$$

with
$$\alpha = \pi' x(0) = \sum_k \pi_k x_k(0) \quad \forall i$$

where $\pi = P' \pi$ is the unique invariant distribution of P .

We are left with proving convergence.

All remaining facts follow from previous arguments.

Asymptotics of French-De Groot learning model

Theorem

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

$$\lim_{t \rightarrow +\infty} x(t) = \alpha \mathbb{1}$$

with
$$\alpha = \pi' x(0) = \sum_k \pi_k x_k(0) \quad \forall i$$

where $\pi = P' \pi$ is the unique invariant distribution of P .

We are left with proving convergence.

All remaining facts follow from previous arguments.

There are extensions for more general graphs.

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

► $M = P - \mathbb{1}\pi'$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

► $M = P - \mathbb{1}\pi'$



$$\begin{aligned} M^2 &= (P - \mathbb{1}\pi')(P - \mathbb{1}\pi') \\ &= P^2 - \mathbb{1}\pi'P - P\mathbb{1}\pi' + \mathbb{1}\pi'\mathbb{1}\pi' \\ &= P^2 - \mathbb{1}\pi' - \mathbb{1}\pi' + \mathbb{1}\pi' \\ &= P^2 - \mathbb{1}\pi' \end{aligned}$$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

► $M = P - \mathbb{1}\pi'$

► $M^t = P^t - \mathbb{1}\pi'$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0, \pi'M = 0$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0$, $\pi'M = 0$
 - ▶ $Mx = \lambda x$ with $\lambda \neq 0$ and $x \neq 0$
 $0 = \pi'Mx = \lambda\pi'x$, hence $\pi'x = 0$ and $Mx = Px$,

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0$, $\pi'M = 0$
 - ▶ $Mx = \lambda x$ with $\lambda \neq 0$ and $x \neq 0$
 $0 = \pi'Mx = \lambda\pi'x$, hence $\pi'x = 0$ and $Mx = Px$,
 - ▶ $Px = \lambda x$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0$, $\pi'M = 0$
 - ▶ $Mx = \lambda x$ with $\lambda \neq 0$ and $x \neq 0$
 $0 = \pi'Mx = \lambda\pi'x$, hence $\pi'x = 0$ and $Mx = Px$,
 - ▶ $Px = \lambda x$
 - ▶ $\pi'x = 0$ implies $x \neq \alpha\mathbb{1}$. Hence $\lambda \neq 1$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0$, $\pi'M = 0$
 - ▶ $Mx = \lambda x$ with $\lambda \neq 0$ and $x \neq 0$
 $0 = \pi'Mx = \lambda\pi'x$, hence $\pi'x = 0$ and $Mx = Px$,
 - ▶ $Px = \lambda x$
 - ▶ $\pi'x = 0$ implies $x \neq \alpha\mathbb{1}$. Hence $\lambda \neq 1$
 - ▶ \mathcal{G} str. connected, aperiodic yields $|\lambda| < 1$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ Eigenvalues of M :
 - ▶ $M\mathbb{1} = P\mathbb{1} - \mathbb{1}\pi'\mathbb{1} = \mathbb{1} - \mathbb{1} = 0$, $\pi'M = 0$
 - ▶ $Mx = \lambda x$ with $\lambda \neq 0$ and $x \neq 0$
 $0 = \pi'Mx = \lambda\pi'x$, hence $\pi'x = 0$ and $Mx = Px$,
 - ▶ $Px = \lambda x$
 - ▶ $\pi'x = 0$ implies $x \neq \alpha\mathbb{1}$. Hence $\lambda \neq 1$
 - ▶ \mathcal{G} str. connected, aperiodic yields $|\lambda| < 1$
 - ▶ $\lambda \in \sigma(M) \Rightarrow |\lambda| < 1$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ $\lambda \in \sigma(M) \Rightarrow |\lambda| < 1$

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ $\lambda \in \sigma(M) \Rightarrow |\lambda| < 1$
- ▶ General fact: if M is a squared matrix with all eigenvalues λ such that $|\lambda| < 1$, then $M^t \rightarrow 0$ (proof using Jordan form)

The quickest proof of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic. $x(t) = P^t x(0)$

π the unique invariant distribution of P ($\pi = P'\pi$ or $\pi' = \pi'P$)

- ▶ $M = P - \mathbb{1}\pi'$
- ▶ $M^t = P^t - \mathbb{1}\pi'$
- ▶ $\lambda \in \sigma(M) \Rightarrow |\lambda| < 1$
- ▶ General fact: if M is a squared matrix with all eigenvalues λ such that $|\lambda| < 1$, then $M^t \rightarrow 0$ (proof using Jordan form)
- ▶ $M^t = P^t - \mathbb{1}\pi' \rightarrow 0 \Rightarrow P^t x(0) - \mathbb{1}\pi' x(0) \rightarrow 0$
Hence, $P^t x(0) \rightarrow (\pi' x(0)) \mathbb{1}$

An alternative road

Contractive properties of stochastic matrices

An alternative road

Contractive properties of stochastic matrices

Q stochastic matrix: $Q_{ij} \geq 0$, $\sum_j Q_{ij} = 1$ for every i .

Consider a vector x and $y = Qx$.

Put $x_{\max} = \max x_j$, $x_{\min} = \min x_j$.

An alternative road

Contractive properties of stochastic matrices

Q stochastic matrix: $Q_{ij} \geq 0$, $\sum_j Q_{ij} = 1$ for every i .

Consider a vector x and $y = Qx$.

Put $x_{\max} = \max x_j$, $x_{\min} = \min x_j$.

$$y_i = \sum_j Q_{ij} x_j \begin{cases} \leq \sum_j Q_{ij} x_{\max} = x_{\max} \\ \geq \sum_j Q_{ij} x_{\min} = x_{\min} \end{cases}$$

An alternative road

Contractive properties of stochastic matrices

Q stochastic matrix: $Q_{ij} \geq 0$, $\sum_j Q_{ij} = 1$ for every i .

Consider a vector x and $y = Qx$.

Put $x_{\max} = \max x_j$, $x_{\min} = \min x_j$.

$$y_i = \sum_j Q_{ij} x_j \begin{cases} \leq \sum_j Q_{ij} x_{\max} = x_{\max} \\ \geq \sum_j Q_{ij} x_{\min} = x_{\min} \end{cases}$$

$$\Rightarrow x_{\min} \leq y_{\min} \leq y_{\max} \leq x_{\max}$$

An alternative road

Contractive properties of stochastic matrices

Q stochastic matrix: $Q_{ij} \geq 0$, $\sum_j Q_{ij} = 1$ for every i .

Consider a vector x and $y = Qx$.

Put $x_{\max} = \max x_j$, $x_{\min} = \min x_j$.

$$y_i = \sum_j Q_{ij} x_j \begin{cases} \leq \sum_j Q_{ij} x_{\max} = x_{\max} \\ \geq \sum_j Q_{ij} x_{\min} = x_{\min} \end{cases}$$

$$\Rightarrow x_{\min} \leq y_{\min} \leq y_{\max} \leq x_{\max}$$

$$\Rightarrow y_{\max} - y_{\min} \leq x_{\max} - x_{\min}$$

An alternative road

Contractive properties of stochastic matrices

Q stochastic matrix: $Q_{ij} \geq 0$, $\sum_j Q_{ij} = 1$ for every i .

Consider a vector x and $y = Qx$.

Put $x_{\max} = \max x_j$, $x_{\min} = \min x_j$.

$$y_i = \sum_j Q_{ij} x_j \begin{cases} \leq \sum_j Q_{ij} x_{\max} = x_{\max} \\ \geq \sum_j Q_{ij} x_{\min} = x_{\min} \end{cases}$$

$$\Rightarrow x_{\min} \leq y_{\min} \leq y_{\max} \leq x_{\max}$$

$$\Rightarrow y_{\max} - y_{\min} \leq x_{\max} - x_{\min}$$

Not yet a real contraction... we have not used connectivity and aperiodicity so far!

An alternative road

Lemma

Let Q be a stochastic matrix for which there exist $\alpha > 0$ and an index k such that $Q_{ik} \geq \alpha$ for all i . Then, $y = Qx$ satisfies

$$y_{\max} - y_{\min} \leq (1 - \alpha)(x_{\max} - x_{\min})$$

$$\begin{aligned} y_i &= \sum_j Q_{ij}x_j = \sum_j Q_{ij}(x_j - x_{\min}) + \sum_j Q_{ij}x_{\min} \\ &\geq \alpha(x_k - x_{\min}) + x_{\min} = \alpha x_k + (1 - \alpha)x_{\min} \end{aligned}$$

$$\begin{aligned} y_i &= \sum_j Q_{ij}x_j = \sum_j Q_{ij}(x_j - x_{\max}) + \sum_j Q_{ij}x_{\max} \\ &\leq \alpha(x_k - x_{\max}) + x_{\max} = \alpha x_k + (1 - \alpha)x_{\max} \end{aligned}$$

An alternative road

Lemma

Let Q be a stochastic matrix for which there exist $\alpha > 0$ and an index k such that $Q_{ik} \geq \alpha$ for all i . Then, $y = Qx$ satisfies

$$y_{\max} - y_{\min} \leq (1 - \alpha)(x_{\max} - x_{\min})$$

For every node i :

$$y_i \geq \alpha x_k + (1 - \alpha)x_{\min}$$

$$y_i \leq \alpha x_k + (1 - \alpha)x_{\max}$$

An alternative road

Lemma

Let Q be a stochastic matrix for which there exist $\alpha > 0$ and an index k such that $Q_{ik} \geq \alpha$ for all i . Then, $y = Qx$ satisfies

$$y_{\max} - y_{\min} \leq (1 - \alpha)(x_{\max} - x_{\min})$$

For every node i :

$$y_i \geq \alpha x_k + (1 - \alpha)x_{\min}$$

$$y_i \leq \alpha x_k + (1 - \alpha)x_{\max}$$

Putting these two inequalities together gives:

$$\begin{aligned} y_{\max} - y_{\min} &\leq \alpha x_k + (1 - \alpha)x_{\max} - \alpha x_k - (1 - \alpha)x_{\min} \\ &= (1 - \alpha)(x_{\max} - x_{\min}), \end{aligned}$$

An alternative road

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), \quad P_{ij} = w_i^{-1} W_{ij}, \quad x(t) = P^t x(0)$$

Suppose there exist $\alpha > 0$ and k such that $P_{ik} \geq \alpha$ for all i

An alternative road

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), \quad P_{ij} = w_i^{-1} W_{ij}, \quad x(t) = P^t x(0)$$

Suppose there exist $\alpha > 0$ and k such that $P_{ik} \geq \alpha$ for all i

$$[x_{\max}(t+1) - x_{\min}(t+1)] \leq (1 - \alpha)[x_{\max}(t) - x_{\min}(t)]$$

$$[x_{\max}(t) - x_{\min}(t)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$$

$$\blacktriangleright \lim_{t \rightarrow +\infty} [x_{\max}(t) - x_{\min}(t)] = 0$$

An alternative road

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P_{ij} = w_i^{-1} W_{ij}, x(t) = P^t x(0)$$

Suppose there exist $\alpha > 0$ and k such that $P_{ik} \geq \alpha$ for all i

$$[x_{\max}(t+1) - x_{\min}(t+1)] \leq (1 - \alpha)[x_{\max}(t) - x_{\min}(t)]$$

$$[x_{\max}(t) - x_{\min}(t)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$$

$$\blacktriangleright \lim_{t \rightarrow +\infty} [x_{\max}(t) - x_{\min}(t)] = 0$$

$$\blacktriangleright x_{\min}(t) \leq x_{\min}(t+1) \leq x_{\max}(t+1) \leq x_{\max}(t) \text{ monotonicity}$$

$$\lim_{t \rightarrow +\infty} x_{\max}(t) = \bar{x} = \lim_{t \rightarrow +\infty} x_{\min}(t)$$

An alternative road

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P_{ij} = w_i^{-1} W_{ij}, x(t) = P^t x(0)$$

Suppose there exist $\alpha > 0$ and k such that $P_{ik} \geq \alpha$ for all i

$$[x_{\max}(t+1) - x_{\min}(t+1)] \leq (1 - \alpha)[x_{\max}(t) - x_{\min}(t)]$$

$$[x_{\max}(t) - x_{\min}(t)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$$

$$\blacktriangleright \lim_{t \rightarrow +\infty} [x_{\max}(t) - x_{\min}(t)] = 0$$

$$\blacktriangleright x_{\min}(t) \leq x_{\min}(t+1) \leq x_{\max}(t+1) \leq x_{\max}(t) \text{ monotonicity}$$

$$\lim_{t \rightarrow +\infty} x_{\max}(t) = \bar{x} = \lim_{t \rightarrow +\infty} x_{\min}(t) \Rightarrow \lim_{t \rightarrow +\infty} x_i(t) = \bar{x} \quad \forall i \text{ consensus}$$

An alternative road


$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

When P satisfies the assumption of the Lemma: $P_{ik} \geq \alpha \forall i$?

An alternative road

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

When P satisfies the assumption of the Lemma: $P_{ik} \geq \alpha \forall i$?



There must exist a *global influencer*, a node to which all others are directly connected!

This is a quite strong assumption!

An alternative road

However,

Lemma

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, P)$ strongly connected, aperiodic. There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .

The alternative proof

- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.

The alternative proof

- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.
- ▶ Apply the Contraction lemma to P^m :

The alternative proof

- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.
- ▶ Apply the Contraction lemma to P^m :
 - ▶ $[x_{\max}(tm) - x_{\min}(tm)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$

The alternative proof

- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.
- ▶ Apply the Contraction lemma to P^m :
 - ▶ $[x_{\max}(tm) - x_{\min}(tm)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$
 - ▶ $\lim_{t \rightarrow +\infty} [x_{\max}(tm) - x_{\min}(tm)] = 0$

The alternative proof


- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.
- ▶ Apply the Contraction lemma to P^m :
 - ▶ $[x_{\max}(tm) - x_{\min}(tm)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$
 - ▶ $\lim_{t \rightarrow +\infty} [x_{\max}(tm) - x_{\min}(tm)] = 0$
- ▶ $x_{\min}(t) \leq x_{\min}(t+1) \leq x_{\max}(t+1) \leq x_{\max}(t)$ monotonicity

The alternative proof

- ▶ There exists $m \in \mathbb{N}$ such that $P_{ij}^m > 0$ for every i, j .
- ▶ Put $\alpha = \min_{ij} P_{ij}^m$.
- ▶ Apply the Contraction lemma to P^m :
 - ▶ $[x_{\max}(tm) - x_{\min}(tm)] \leq (1 - \alpha)^t [x_{\max}(0) - x_{\min}(0)]$
 - ▶ $\lim_{t \rightarrow +\infty} [x_{\max}(tm) - x_{\min}(tm)] = 0$
- ▶ $x_{\min}(t) \leq x_{\min}(t+1) \leq x_{\max}(t+1) \leq x_{\max}(t)$ monotonicity
- ▶ $\lim_{t \rightarrow +\infty} x_{\max}(t) = \bar{x} = \lim_{t \rightarrow +\infty} x_{\min}(t)$
- ▶ $\Rightarrow \lim_{t \rightarrow +\infty} x_i(t) = \bar{x}$

Extensions of the main theorem

\mathcal{G} possesses a globally reachable connected component \mathcal{C}_0 (equivalently, the condensation graph has just one sink $s_{\mathcal{G}} = 1$) that is aperiodic.



- ▶ Global reachability: Fix $k \in \mathcal{C}_0$. For every i , $P_{ik}^{m_i} > 0$ for some m_i .
- ▶ Aperiodicity lemma $P_{kk}^q > 0$ for every $q \geq m$
- ▶ $P_{ik}^s > 0$ for every i , where $s = \max m_i + m$.
- ▶ Apply the contractive lemma again!

The most general result

Theorem

If $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ possesses a globally reachable aperiodic component \mathcal{C}_0 , then

$$\lim_{t \rightarrow +\infty} x_i(t) = \pi'x(0) \quad \forall i$$

where $\pi = P'\pi$ is the invariant measure centrality of the graph

π has support only on the globally reachable component.

The opinions of agents not belonging to the globally reachable component have no influence on the final consensus value.


A remark

$$P^t \rightarrow \mathbb{1}\pi'$$

Namely, P^t converges to a matrix where the rows are all the same, equal to the invariant distribution centrality π .

The most general result

No further generalization is possible:



No influence between nodes in \mathcal{C}' and \mathcal{C}'' .

Nodes in \mathcal{C}' and \mathcal{C}'' (if the components are aperiodic) will reach separated consensus depending on their own initial opinions.


No global consensus

The most general result

- ▶ Lack of periodicity \Rightarrow no convergence.
- ▶ More than one sink in the condensation graph \Rightarrow no consensus.

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




$$P = ???$$

$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



- \mathcal{G} has an aperiodic globally reachable component


$$P = ???$$

$$x(t) = P^t x(0), x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

Example

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$P = ???$$

$$x(t) = P^t x(0), \quad x(0) = (0, 1, 2, 5, 3)$$

$$\lim_{t \rightarrow +\infty} x(t)?$$

- ▶ \mathcal{G} has an aperiodic globally reachable component
- ▶ $\lim_{t \rightarrow +\infty} x_i(t) = \pi' x(0)$ for every i
- ▶ $\pi = (1/3, 1/3, 1/3, 0, 0)$
- ▶ $\pi' x(0) = 1$

Other applications of the linear averaging dynamics

The basis of many distributed algorithms

- ▶ Decentralized computation in sensor networks
- ▶ Load balancing in computer networks
- ▶ Clock synchronization
- ▶ Relative localization
- ▶ Coordination dynamics of robot networks.


Other applications of the linear averaging dynamics

The basis of many distributed algorithms

- ▶ Decentralized computation in sensor networks
- ▶ Load balancing in computer networks
- ▶ Clock synchronization
- ▶ Relative localization
- ▶ Coordination dynamics of robot networks.

Decentralized computation of global functions


$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state

Decentralized computation of global functions

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network




$i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute some global function
 $f(x_i \mid i \in \mathcal{V})$

Decentralized computation of global functions

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state


Goal: compute some global function
 $f(x_i \mid i \in \mathcal{V})$

Constraints:

- ▶ no supervision, decentralized design;
- ▶ use only the available communication links;
- ▶ time and computation complexity scaling well w.r. to size $n = |\mathcal{V}|$;

Decentralized computation of global functions

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state


Goal: compute some global function
 $f(x_i \mid i \in \mathcal{V})$

Examples:

x_i internal state (e.g. level of energy) or the result of a measurement (temperature, presence detection, fire detection)

Decentralized computation of global functions

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute some global function
 $f(x_i \mid i \in \mathcal{V})$

Examples:

x_i internal state (e.g. level of energy) or the result of a measurement (temperature, presence detection, fire detection)

f computing some statistics of the data:

- ▶ average value $\bar{x} = n^{-1} \sum x_i$
- ▶ variance $n^{-1} \sum (x_i - \bar{x})^2$
- ▶ $\max x_i$, $\min x_i$
- ▶ fraction of nodes s.t. $x_i \geq \alpha$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

Important remark: in this case, there is not an a-priori choice for the matrix P : this just becomes a *design* choice.

Given an $n \times n$ matrix P , we can consider the associated graph $\mathcal{G}_P = (\mathcal{V}, \mathcal{E})$ where

$$\mathcal{V} = \{1, \dots, n\}, \mathcal{E} = \{(i, j) \mid P_{ij} > 0\}$$

Notice that, with the choice above, $\mathcal{G}_P = \mathcal{G} \cup \{\text{selfloops}\}$: to implement P we only need to communicate along the edges of the graph \mathcal{G} ; use of self-loops is not an issue.

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

$$x(t+1)_i = \frac{1}{2}x(t)_i + \frac{1}{2w_i} \sum_{j \in N_i} x(t)_j$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

$\mathcal{G}_P = \mathcal{G} \cup \{\text{selfloops}\}$ str. connected, aperiodic

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

$\mathcal{G}_P = \mathcal{G} \cup \{\text{selfloops}\}$ **str. connected, aperiodic**

$$\Rightarrow x(t)_i \rightarrow x^* = \sum \pi_j x_j \quad \pi_j = w_j / |\mathcal{E}| \text{ centrality}$$

$$((D^{-1}W)'\pi = \pi \Leftrightarrow P'\pi = \pi)$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

$\mathcal{G}_P = \mathcal{G} \cup \{\text{selfloops}\}$ **str. connected, aperiodic**

$$\Rightarrow x(t)_i \rightarrow x^* = \sum \pi_j x_j \quad \pi_j = w_j / |\mathcal{E}| \text{ centrality}$$

$$((D^{-1}W)'\pi = \pi \Leftrightarrow P'\pi = \pi)$$

$$x^* = \bar{x} \Leftrightarrow \mathcal{G} \text{ is regular}$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$ **Idea:** use averaging dynamics.

W adjacency matrix of \mathcal{G} , $P = (D^{-1}W + I)/2$

$$x(t+1) = Px(t), \quad x(0)_i = x_i$$

$\mathcal{G}_P = \mathcal{G} \cup \{\text{selfloops}\}$ str. connected, aperiodic

$$\Rightarrow x(t)_i \rightarrow x^* = \sum \pi_j x_j \quad \pi_j = w_j / |\mathcal{E}| \text{ centrality}$$

$$((D^{-1}W)'\pi = \pi \Leftrightarrow P'\pi = \pi)$$

$$x^* = \bar{x} \Leftrightarrow \mathcal{G} \text{ is regular} \quad \text{What if } \mathcal{G} \text{ is not regular?}$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$

Possible solutions:

- Use P as follows

$$\begin{cases} x(t+1) = Px(t), & x(0)_i = x_i/w_i \\ y(t+1) = Py(t), & y(0)_i = 1/w_i \end{cases}$$

Check (exercise): $\frac{x(t)_i}{y(t)_i} \rightarrow \bar{x}$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ undirected str. connected, $i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$

Possible solutions:

- Use P as follows

$$\begin{cases} x(t+1) = Px(t), & x(0)_i = x_i/w_i \\ y(t+1) = Py(t), & y(0)_i = 1/w_i \end{cases}$$


Check (exercise): $\frac{x(t)_i}{y(t)_i} \rightarrow \bar{x}$

- Find Q stochastic symmetric s.t. $\mathcal{G}_Q \subseteq \mathcal{G} \cup \{\text{selfloops}\}$, \mathcal{G}_Q aperiodic (exercise)

$$x(t+1) = Qx(t), \quad x(0)_i = x_i, \quad x(t)_i \rightarrow \bar{x}$$

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state


Goal: compute $\bar{x} = n^{-1} \sum x_i$

Using average consensus algorithms:

- ▶ no supervision, decentralized design;
- ▶ use only the available communication links;

Average computation

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ set of units (sensors)
connected through a network



$i \in \mathcal{V} \rightarrow x_i$ state

Goal: compute $\bar{x} = n^{-1} \sum x_i$

Using average consensus algorithms:

- ▶ no supervision, decentralized design;
- ▶ use only the available communication links;
- ▶ time and computation complexity?;

Speed of convergence

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}) \text{ str. connected, } P = (D^{-1}W + I)/2$$

$$x(t+1) = Px(t), \quad x(t) \rightarrow \mathbb{1}(\pi'x(0))$$

How fast $x(t)$ converges to consensus?

Speed of convergence

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ str. connected, $P = (D^{-1}W + I)/2$

$x(t+1) = Px(t), \quad x(t) \rightarrow \mathbb{1}(\pi'x(0))$

How fast $x(t)$ converges to consensus?

Theorem

\mathcal{G} undirected str. connected. $P = (D^{-1}W + I)/2$.

► Eigenvalues of P : $1 = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_n \geq 0$

►

$$\|x(t) - \mathbb{1}\pi'x(0)\|_2 \leq \sqrt{\frac{\max \pi_i}{\min \pi_i}} \lambda_2^t \|x(0)\|_2$$

Speed of convergence and computational complexity

$$\|x(t) - \mathbb{1}\pi'x(0)\|_2 \leq \frac{\max \sqrt{\pi_i}}{\min \sqrt{\pi_i}} \lambda_2^t \|x(0)\|_2$$

Speed of convergence and computational complexity

$$\|x(t) - \mathbb{1}\pi'x(0)\|_2 \leq \frac{\max \sqrt{\pi_i}}{\min \sqrt{\pi_i}} \lambda_2^t \|x(0)\|_2$$

$$\sqrt{\frac{\max \pi_i}{\min \pi_i}} \lambda_2^t \leq \epsilon \Leftrightarrow t \geq \frac{\log(\epsilon^{-1} \max \pi_i / \min \pi_i)}{2 \log \lambda_2^{-1}}$$

Speed of convergence and computational complexity

$$\|x(t) - \mathbb{1}\pi'x(0)\|_2 \leq \frac{\max \sqrt{\pi_i}}{\min \sqrt{\pi_i}} \lambda_2^t \|x(0)\|_2$$

$$\sqrt{\frac{\max \pi_i}{\min \pi_i}} \lambda_2^t \leq \epsilon \Leftrightarrow t \geq \frac{\log(\epsilon^{-1} \max \pi_i / \min \pi_i)}{2 \log \lambda_2^{-1}}$$

► *Convergence time:* $\tau_{conv}(\epsilon) := \frac{\log(\epsilon^{-1} \max \pi_i / \min \pi_i)}{2 \log \lambda_2^{-1}}$

$$\log \lambda_2^{-1} \sim 1 - \lambda_2 \text{ spectral gap (for } \lambda_2 \rightarrow 1)$$

Speed of convergence and computational complexity

$$\|x(t) - \mathbb{1}\pi'x(0)\|_2 \leq \frac{\max \sqrt{\pi_i}}{\min \sqrt{\pi_i}} \lambda_2^t \|x(0)\|_2$$

$$\sqrt{\frac{\max \pi_i}{\min \pi_i}} \lambda_2^t \leq \epsilon \Leftrightarrow t \geq \frac{\log(\epsilon^{-1} \max \pi_i / \min \pi_i)}{2 \log \lambda_2^{-1}}$$

► *Convergence time:* $\tau_{conv}(\epsilon) := \frac{\log(\epsilon^{-1} \max \pi_i / \min \pi_i)}{2 \log \lambda_2^{-1}}$

$$\log \lambda_2^{-1} \sim 1 - \lambda_2 \text{ spectral gap (for } \lambda_2 \rightarrow 1)$$

► *Computation complexity per node:* $\gamma(\epsilon) = \frac{\tau_{conv}(\epsilon)|\mathcal{E}|}{|\mathcal{V}|}$

Performance comparison

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ family of graphs with increasing size $n = |\mathcal{V}|$

	$1 - \lambda_2$	τ_{conv}	diam	γ
Line, cycle	C/n^2	Cn^2	Cn	Cn^2
d-dimensional Grids	$Cn^{2/d}$	$n^{2/d}$	$Cn^{1/d}$	$Cdn^{2/d}$
Complete	$1/2$	1	1	Cn
Expanders	C	$C \log n$	$C \log n$	$C \log n$

Performance comparison

$\mathcal{G} = (\mathcal{V}, \mathcal{E})$ family of graphs with increasing size $n = |\mathcal{V}|$

	$1 - \lambda_2$	τ_{conv}	diam	γ
Line, cycle	C/n^2	Cn^2	Cn	Cn^2
<i>d</i>-dimensional Grids	$Cn^{2/d}$	$n^{2/d}$	$Cn^{1/d}$	$Cdn^{2/d}$
Complete	$1/2$	1	1	Cn
Expanders	C	$C \log n$	$C \log n$	$C \log n$

Expanders \supseteq Random graphs, Barabasi, Small world

Network flow dynamics

- ▶ Models of transport phenomena
- ▶ They find applications in several fields: infrastructure networks, epidemiology, ecology, pharmacokinetics
- ▶ In some of the literature they are referred to as *compartmental systems*

Network flow dynamics

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ physical network

- ▶ \mathcal{V} cells containing a homogeneous mass of some matter,
- ▶ \mathcal{E} physical constraints: the matter can flow directly from cell i to cell j if $(i, j) \in \mathcal{E}$,
- ▶ $y_i(t)$ mass present in node i at time t .
- ▶ $f_{ij}(t)$ mass flowing from i to j at time t ($f_{ij}(t) = 0$ if $(i, j) \notin \mathcal{E}$)

Mass conservation law prescribes that

$$y_i(t+1) = y_i(t) + \sum_j f_{ji}(t) - \sum_j f_{ij}(t)$$

Linear network flow dynamics

$$y_i(t+1) = y_i(t) + \sum_j f_{ji}(t) - \sum_j f_{ij}(t)$$

We study the *linear* case: $f_{ij}(t) = y_i(t)P_{ij}$ where $P = D^{-1}W$.

Linear network flow dynamics

$$y_i(t+1) = y_i(t) + \sum_j f_{ji}(t) - \sum_j f_{ij}(t)$$

We study the *linear* case: $f_{ij}(t) = y_i(t)P_{ij}$ where $P = D^{-1}W$.

The amount of matter flowing from i to j is proportional to the amount present in i .

$$y_i(t+1) = y_i(t) + \sum_j y_j(t)P_{ji} - \sum_j y_i(t)P_{ij} = \sum_j y_j(t)P_{ji}$$

Linear network flow dynamics

$$y_i(t+1) = y_i(t) + \sum_j f_{ji}(t) - \sum_j f_{ij}(t)$$

We study the *linear* case: $f_{ij}(t) = y_i(t)P_{ij}$ where $P = D^{-1}W$.

The amount of matter flowing from i to j is proportional to the amount present in i .

$$y_i(t+1) = y_i(t) + \sum_j y_j(t)P_{ji} - \sum_j y_i(t)P_{ij} = \sum_j y_j(t)P_{ji}$$

Compact notation:

$$y(t+1) = P'y(t)$$

- ▶ Equilibria: $P'y = y$ eigenvectors of P' of eigenvalue 1
- ▶ Invariant distributions are equilibria (this is why they are called invariant!)
- ▶ $\mathbb{1}'y(t+1) = \mathbb{1}'P'y(t) = \mathbb{1}'y(t)$ **Total mass is a motion invariant**

Linear network flow dynamics

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

$P = D^{-1}W$, $\pi = P'\pi$ invariant distribution centrality

Linear network flow dynamics

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

$P = D^{-1}W$, $\pi = P'\pi$ invariant distribution centrality

The only equilibria are scalar multiples of π : $y = a\pi$.

Linear network flow dynamics

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

$P = D^{-1}W$, $\pi = P'\pi$ invariant distribution centrality

The only equilibria are scalar multiples of π : $y = a\pi$.

$$\lim_{t \rightarrow +\infty} P^t = \mathbb{1}\pi'$$

Linear network flow dynamics

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ strongly connected, aperiodic.

$P = D^{-1}W$, $\pi = P'\pi$ invariant distribution centrality

The only equilibria are scalar multiples of π : $y = a\pi$.

$$\lim_{t \rightarrow +\infty} P^t = \mathbb{1}\pi'$$

$$\lim_{t \rightarrow +\infty} y(t) = \lim_{t \rightarrow +\infty} P'^t y(0) = \pi \mathbb{1}' y(0)$$

Asymptotically, mass distributes according to π

Continuous time dynamics models

The linear averaging and the linear network flow dynamics on a graph $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ have an analogue in continuous time:

$$\dot{x} = -Lx, \quad \dot{y} = -L'y$$

where $L = D - W$ is the Laplacian of \mathcal{G} .

Theorem

Let $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ be a graph and let L be its Laplacian. If $s_{\mathcal{G}} = 1$, then

$$\lim_{t \rightarrow +\infty} x(t) = \mathbb{1}\bar{\pi}'x(0) \quad (1)$$

$$\lim_{t \rightarrow +\infty} y(t) = \bar{\pi}\mathbb{1}'y(0) \quad (2)$$

where $\bar{\pi}$ is the unique Laplace invariant probability distribution of \mathcal{G} ($L'\bar{\pi} = 0$, $\mathbb{1}'\bar{\pi} = 1$).