

# Algebraic graph theory and centrality measures

Giacomo Como, DISMA, Politecnico di Torino

Fabio Fagnani, DISMA, Politecnico di Torino

## Weight of walks and powers of the weight matrix

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$$\gamma = (i_0, i_1, \dots, i_l) \text{ walk, } W_\gamma = \prod_{1 \leq h \leq l} W_{i_{h-1}i_h}$$

# Weight of walks and powers of the weight matrix

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$$\gamma = (i_0, i_1, \dots, i_l) \text{ walk, } W_\gamma = \prod_{1 \leq h \leq l} W_{i_{h-1}i_h}$$

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ .  $l = 1$  trivial.

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ . Assume it true for  $l$ .

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ . Assume it true for  $l$ .

$$\sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l + 1}} W_\gamma$$

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ . Assume it true for  $l$ .

$$\sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l+1}} W_\gamma = \sum_k \sum_{\substack{\tilde{\gamma} \text{ walk from } i \text{ to } k \\ l(\tilde{\gamma}) = l}} W_{\tilde{\gamma}} W_{kj}$$



# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ . Assume it true for  $l$ .

$$\begin{aligned} \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l+1}} W_\gamma &= \sum_k \sum_{\substack{\tilde{\gamma} \text{ walk from } i \text{ to } k \\ l(\tilde{\gamma}) = l}} W_{\tilde{\gamma}} W_{kj} \\ &= \sum_k (W^l)_{ik} W_{kj} = (W^{l+1})_{ij} \end{aligned}$$

# Weight of walks and powers of the weight matrix

Theorem

$$(W^l)_{ij} = \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l}} W_\gamma$$

**Proof** By induction on  $l \geq 1$ . Assume it true for  $l$ .

$$\begin{aligned} \sum_{\substack{\gamma \text{ walk from } i \text{ to } j \\ l(\gamma) = l+1}} W_\gamma &= \sum_k \sum_{\substack{\tilde{\gamma} \text{ walk from } i \text{ to } k \\ l(\tilde{\gamma}) = l}} W_{\tilde{\gamma}} W_{kj} \\ &= \sum_k (W^l)_{ik} W_{kj} = (W^{l+1})_{ij} \end{aligned}$$

It is proven for  $l+1$ . By induction result is proven.

# Properties of the products of the weight matrix

## Theorem

Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  be a graph. Then,

1.  $(W^l)_{ij} > 0$  if and only if there exists a walk of length  $l$  from  $i$  to  $j$ ;
2.  $\mathcal{G}$  is strongly connected iff for every  $i, j \in \mathcal{V}$ , there exists  $l > 0$  such that  $(W^l)_{ij} > 0$ .
3.  $\mathcal{G}$  is strongly connected and aperiodic iff there exists  $N > 0$  such that  $(W^N)_{ij} > 0$  for every  $i, j \in \mathcal{V}$ .

# Properties of the products of the weight matrix

## Theorem

Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  be a graph. Then,


1.  $(W^l)_{ij} > 0$  if and only if there exists a walk of length  $l$  from  $i$  to  $j$ ;
2.  $\mathcal{G}$  is strongly connected iff for every  $i, j \in \mathcal{V}$ , there exists  $l > 0$  such that  $(W^l)_{ij} > 0$ .
3.  $\mathcal{G}$  is strongly connected and aperiodic iff there exists  $N > 0$  such that  $(W^N)_{ij} > 0$  for every  $i, j \in \mathcal{V}$ .

Comments on the proof:

- ▶ 1. and 2. are consequences of previous theorem.
- ▶ 3. is more involved.


## Examples

$C_5$



# Examples


$C_5$



$$W = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

# Examples

$C_5$




$$W = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

$$W^2 = \begin{pmatrix} 2 & 0 & 1 & 1 & 0 \\ 0 & 2 & 0 & 1 & 1 \\ 1 & 0 & 2 & 0 & 1 \\ 1 & 1 & 0 & 2 & 0 \\ 0 & 1 & 1 & 0 & 2 \end{pmatrix}$$

# Examples

$C_5$




$$W = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

$$W^3 = \begin{pmatrix} 0 & 3 & 1 & 1 & 3 \\ 3 & 0 & 3 & 1 & 1 \\ 1 & 3 & 0 & 3 & 1 \\ 1 & 1 & 3 & 0 & 3 \\ 3 & 1 & 1 & 3 & 0 \end{pmatrix}$$



# Examples

$C_5$




$$W = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

$$W^4 = \begin{pmatrix} 6 & 1 & 4 & 4 & 1 \\ 1 & 6 & 1 & 4 & 4 \\ 4 & 1 & 6 & 1 & 4 \\ 4 & 4 & 1 & 6 & 1 \\ 1 & 4 & 4 & 1 & 6 \end{pmatrix}$$


## Examples

$C_4$



## Examples


$C_4$



$$W = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

# Examples

$C_4$




$$W = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

$$W^2 = \begin{pmatrix} 2 & 0 & 2 & 0 \\ 0 & 2 & 0 & 2 \\ 2 & 0 & 2 & 0 \\ 0 & 2 & 0 & 2 \end{pmatrix}$$

## Examples

$C_4$




$$W = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

$$W^3 = \begin{pmatrix} 0 & 4 & 0 & 4 \\ 4 & 0 & 4 & 0 \\ 0 & 4 & 0 & 4 \\ 4 & 0 & 4 & 0 \end{pmatrix}$$

# Examples

$C_4$



$$W = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

$$W^4 = \begin{pmatrix} 8 & 0 & 8 & 0 \\ 0 & 8 & 0 & 8 \\ 8 & 0 & 8 & 0 \\ 0 & 8 & 0 & 8 \end{pmatrix}$$

# Counting objects in simple graphs

## Corollary

*For a simple graph  $\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$ , we have that:*

- (i)  $(W^2)_{ii} = w_i$  for every  $i \in \mathcal{V}$ ;*
- (ii)  $Tr(W^2) = |\mathcal{E}|$ ;*
- (iii)  $Tr(W^3) = 6 \cdot \text{number of triangles}.$*

## The normalized weight matrix

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  graph,  $w_i = \sum_j W_{ij}$  out-degrees

$P_{ij} = w_i^{-1} W_{ij}$  *normalized weight matrix* of  $\mathcal{G}$ .

More compactly,  $P = D^{-1}W$  where  $D$  is diagonal with  $D_i = w_i$

$P$  is a *stochastic* matrix:  $P_{ij} \geq 0$ ,  $P\mathbb{1} = \mathbb{1}$



# The normalized weight matrix

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  graph,  $w_i = \sum_j W_{ij}$  out-degrees

$P_{ij} = w_i^{-1} W_{ij}$  *normalized weight matrix* of  $\mathcal{G}$ .

More compactly,  $P = D^{-1}W$  where  $D$  is diagonal with  $D_i = w_i$

$P$  is a *stochastic* matrix:  $P_{ij} \geq 0$ ,  $P\mathbb{1} = \mathbb{1}$

- ▶ Topology of  $\mathcal{G} \leftrightarrow$  Spectral properties of  $P$ ;
- ▶ Through  $P$  we can describe interesting dynamical systems over  $\mathcal{G}$ ;
- ▶  $P$  can be interpreted as the transition matrix of a Markov chain, a *random walk* over  $\mathcal{G}$ .

# The Laplacian matrix

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  graph,  $D$  diagonal matrix with  $D_i = w_i$  out-degrees

$$L = D - W$$

$$L_{ij} = \begin{cases} -W_{ij} & \text{if } i \neq j \\ w_i - W_{ii} & \text{if } i = j \end{cases}$$

# The Laplacian matrix

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  graph,  $D$  diagonal matrix with  $D_i = w_i$  out-degrees

$$L = D - W$$

$$L_{ij} = \begin{cases} -W_{ij} & \text{if } i \neq j \\ w_i - W_{ii} & \text{if } i = j \end{cases}$$

$-L$  is called a Metzler matrix (non-negative off diagonal),  $L\mathbb{1} = 0$

# The Laplacian matrix

$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$  graph,  $D$  diagonal matrix with  $D_i = w_i$  out-degrees

$$L = D - W$$


$$L_{ij} = \begin{cases} -W_{ij} & \text{if } i \neq j \\ w_i - W_{ii} & \text{if } i = j \end{cases}$$

$-L$  is called a Metzler matrix (non-negative off diagonal),  $L\mathbb{1} = 0$

- ▶ Topology of  $\mathcal{G} \leftrightarrow$  Spectral properties of  $L$ ;
- ▶ Through  $L$  we can describe interesting dynamical systems over  $\mathcal{G}$  in continuous time;

# Perron-Frobenius theory

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




## Theorem (**Perron-Frobenius**)

*There exists  $\lambda_W \geq 0$  and non-negative vectors  $x \neq 0$ ,  $y \neq 0$  s.t.*

- ▶  $Wx = \lambda_W x$ ,  $W'y = \lambda_W y$ ;
- ▶ *every eigenvalue  $\mu$  of  $W$  is such that  $|\mu| \leq \lambda_W$ ;*

# Perron-Frobenius theory

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




## Theorem (**Perron-Frobenius**)

*There exists  $\lambda_W \geq 0$  and non-negative vectors  $x \neq 0$ ,  $y \neq 0$  s.t.*

- ▶  $Wx = \lambda_W x$ ,  $W'y = \lambda_W y$ ;
- ▶ *every eigenvalue  $\mu$  of  $W$  is such that  $|\mu| \leq \lambda_W$ ;*
- ▶ *If  $\mathcal{G}$  is strongly connected, then  $\lambda_W$  is simple;*

# Perron-Frobenius theory

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




## Theorem (**Perron-Frobenius**)

*There exists  $\lambda_W \geq 0$  and non-negative vectors  $x \neq 0$ ,  $y \neq 0$  s.t.*

- ▶  $Wx = \lambda_W x$ ,  $W'y = \lambda_W y$ ;
- ▶ *every eigenvalue  $\mu$  of  $W$  is such that  $|\mu| \leq \lambda_W$ ;*
- ▶ *If  $\mathcal{G}$  is strongly connected, then  $\lambda_W$  is simple;*
- ▶ *If  $\mathcal{G}$  is strongly connected and aperiodic, then every eigenvalue  $\mu \neq \lambda_W$  is s.t.  $|\mu| < \lambda_W$ .*

# Perron-Frobenius theory

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



## Theorem (**Perron-Frobenius**)

*There exists  $\lambda_W \geq 0$  and non-negative vectors  $x \neq 0$ ,  $y \neq 0$  s.t.*

- ▶  $Wx = \lambda_W x$ ,  $W'y = \lambda_W y$ ;
- ▶ *every eigenvalue  $\mu$  of  $W$  is such that  $|\mu| \leq \lambda_W$ ;*
- ▶ *If  $\mathcal{G}$  is strongly connected, then  $\lambda_W$  is simple;*
- ▶ *If  $\mathcal{G}$  is strongly connected and aperiodic, then every eigenvalue  $\mu \neq \lambda_W$  is s.t.  $|\mu| < \lambda_W$ .*

$\lambda_W$  *dominant eigenvalue of  $W$ .*



## Perron-Frobenius theory applied to graphs


$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$$P = D^{-1}W \text{ normalized weight matrix}$$

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$$P = D^{-1}W \text{ normalized weight matrix}$$




## Theorem (Spectral properties of stochastic matrices)

- ▶  $\lambda_P = 1$ ;
- ▶  $P\mathbb{1} = \mathbb{1}$ ,  $\exists \pi \geq 0$  s.t.  $\mathbb{1}'\pi = 1$  and  $P'\pi = \pi$ ;
- ▶ every eigenvalue  $\mu$  of  $P$  is such that  $|\mu| \leq 1$ ;

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$P = D^{-1}W$  normalized weight matrix




## Theorem (Spectral properties of stochastic matrices)

- ▶  $\lambda_P = 1$ ;
- ▶  $P\mathbb{1} = \mathbb{1}$ ,  $\exists \pi \geq 0$  s.t.  $\mathbb{1}'\pi = 1$  and  $P'\pi = \pi$ ;
- ▶ every eigenvalue  $\mu$  of  $P$  is such that  $|\mu| \leq 1$ ;
- ▶ If  $\mathcal{G}$  is strongly connected, then 1 is simple and  $\pi_i > 0$  for all  $i$ ;
- ▶ If  $\mathcal{G}$  is strongly connected and aperiodic, then every eigenvalue  $\mu \neq 1$  is s.t.  $|\mu| < 1$ .

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$P = D^{-1}W$  normalized weight matrix




## Theorem (Spectral properties of stochastic matrices)

- ▶  $\lambda_P = 1$ ;
- ▶  $P\mathbb{1} = \mathbb{1}$ ,  $\exists \pi \geq 0$  s.t.  $\mathbb{1}'\pi = 1$  and  $P'\pi = \pi$ ;
- ▶ every eigenvalue  $\mu$  of  $P$  is such that  $|\mu| \leq 1$ ;
- ▶ If  $\mathcal{G}$  is strongly connected, then 1 is simple and  $\pi_i > 0$  for all  $i$ ;
- ▶ If  $\mathcal{G}$  is strongly connected and aperiodic, then every eigenvalue  $\mu \neq 1$  is s.t.  $|\mu| < 1$ .

$\pi$  invariant distribution

## A digression: convex sets...


A subset  $\mathcal{K} \subseteq \mathbb{R}^n$  is called *convex* if given any two points in  $\mathcal{K}$ , the segment joining them is all inside  $\mathcal{K}$



## polytopes...

A *polytope* is a convex subset that can be obtained by taking convex combinations of  $k + 1$  vectors  $x^1, \dots, x^{k+1} \in \mathbb{R}^n$ ,

$$\mathcal{K} = \{x = \sum_i \lambda_i x^i \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$



$x^i$  extremal points of the polytope.

# Simplexes


A *k-simplex* is a convex subset that can be obtained by taking convex combinations of  $k + 1$  vectors  $x^1, \dots, x^{k+1} \in \mathbb{R}^n$ , **s.t.**  $x^i - x^1$  are independent vectors

$$\mathcal{K} = \{x = \sum_i \lambda_i x^i \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$


# Simplexes

A  $k$ -simplex is a convex subset that can be obtained by taking convex combinations of  $k + 1$  vectors  $x^1, \dots, x^{k+1} \in \mathbb{R}^n$ , s.t.  $x^i - x^1$  are independent vectors

$$\mathcal{K} = \{x = \sum_i \lambda_i x^i \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$



polytope, not a simplex





simplex



# Simplexes

A *k-simplex* is a convex subset that can be obtained by taking convex combinations of  $k + 1$  vectors  $x^1, \dots, x^{k+1} \in \mathbb{R}^n$ , s.t.  $x^i - x^1$  are independent vectors

$$\mathcal{K} = \{x = \sum_i \lambda_i x^i \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$



- ▶ 0-simplex is a point,
- ▶ 1-simplex is a segment,
- ▶ 2-simplex is a triangle,
- ▶ ...

# Simplexes

Example:  $x^i = e^i \in \mathbb{R}^n$  canonical basis

$$\mathcal{K} = \{x = (\lambda_1, \dots, \lambda_n); \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$

$(n - 1)$ -simplex of probability vectors in  $\mathbb{R}^n$

# Simplexes

Example:  $x^i = e^i \in \mathbb{R}^n$  canonical basis

$$\mathcal{K} = \{x = (\lambda_1, \dots, \lambda_n); \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$

$(n - 1)$ -simplex of probability vectors in  $\mathbb{R}^n$

$P$  stochastic matrix:  $P_{ij} \geq 0, \sum_j P_{ij} = 1$

$$p \in \mathcal{K} \Rightarrow P'p \in \mathcal{K}$$

# Simplexes

Example:  $x^i = e^i \in \mathbb{R}^n$  canonical basis

$$\mathcal{K} = \{x = (\lambda_1, \dots, \lambda_n); \mid \lambda_i \geq 0, \sum_i \lambda_i = 1\}$$

$(n - 1)$ -simplex of probability vectors in  $\mathbb{R}^n$

$P$  stochastic matrix:  $P_{ij} \geq 0, \sum_j P_{ij} = 1$

$$p \in \mathcal{K} \Rightarrow P'p \in \mathcal{K}$$

**Perron-Frobenius theory**  $\Rightarrow \exists$  a fixed point  $\pi \in \mathcal{K}$ :  $P'\pi = \pi$

## Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A more refined result.

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A more refined result.

$\mathcal{H}_{\mathcal{G}}$  condensation graph,  $s_{\mathcal{G}}$  number of sinks in  $\mathcal{H}_{\mathcal{G}}$ .

## Theorem

- ▶ *Invariant distributions  $\pi$*

$$\pi \geq 0, \mathbb{1}'\pi = 1, P'\pi = \pi$$

*form a simplex in  $\mathbb{R}^{\mathcal{V}}$  with  $s_{\mathcal{G}}$  vertices.*

- ▶ *For every sink component with nodes  $\mathcal{W}$ , there exists an invariant distribution  $\pi$  such that  $\pi_i > 0$  if and only if  $i \in \mathcal{W}$ .*
- ▶ *The invariant distribution is unique if and only if  $s_{\mathcal{G}} = 1$ .*

## The special case of balanced graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A remarkable fact.

Suppose  $\mathcal{G}$  is balanced ( $w_i = w_i^-$  for every  $i \in \mathcal{V}$ ). Then,

## The special case of balanced graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A remarkable fact.

Suppose  $\mathcal{G}$  is balanced ( $w_i = w_i^-$  for every  $i \in \mathcal{V}$ ). Then,

$$\sum_i w_i P_{ij} = \sum_i w_i w_i^{-1} W_{ij} = \sum_i W_{ij} = w_j^- = w_j$$



## The special case of balanced graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A remarkable fact.

Suppose  $\mathcal{G}$  is balanced ( $w_i = w_i^-$  for every  $i \in \mathcal{V}$ ). Then,

$$\sum_i w_i P_{ij} = \sum_i w_i w_i^{-1} W_{ij} = \sum_i W_{ij} = w_j^- = w_j$$

$$P'w = w$$

# The special case of balanced graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W), P = D^{-1}W.$$

A remarkable fact.

Suppose  $\mathcal{G}$  is balanced ( $w_i = w_i^-$  for every  $i \in \mathcal{V}$ ). Then,

$$\sum_i w_i P_{ij} = \sum_i w_i w_i^{-1} W_{ij} = \sum_i W_{ij} = w_j^- = w_j$$


$$P'w = w$$

- ▶ the out-degree vector  $w$  is an eigenvector of eigenvalue 1
- ▶  $\pi = \frac{1}{|\mathcal{E}|} w$  is an invariant distribution of  $\mathcal{G}$ !
- ▶ If the graph is strongly connected, this is the unique invariant distribution.

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$L = D - W$  Laplacian matrix




## Theorem (Spectral properties of the Laplacian)

- ▶  $L\mathbb{1} = 0$ , 0 is an eigenvalue of  $L$
- ▶  $\exists \bar{\pi} \geq 0$  s.t.  $\mathbb{1}'\bar{\pi} = 1$  and  $L'\bar{\pi} = 0$ ;
- ▶ All other eigenvalues  $\lambda$  have  $\text{Re}(\lambda) > 0$ ;
- ▶ If  $\mathcal{G}$  is strongly connected, then 0 is simple and  $\bar{\pi}_i > 0$  for all  $i$ ;
- ▶  $L'\bar{y} = 0 \Leftrightarrow P'(D\bar{y}) = (D\bar{y})$ ;

# Perron-Frobenius theory applied to graphs

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$

$L = D - W$  Laplacian matrix




## Theorem (Spectral properties of the Laplacian)

- ▶  $L\mathbb{1} = 0$ ,  $0$  is an eigenvalue of  $L$
- ▶  $\exists \bar{\pi} \geq 0$  s.t.  $\mathbb{1}'\bar{\pi} = 1$  and  $L'\bar{\pi} = 0$ ;
- ▶ All other eigenvalues  $\lambda$  have  $\mathcal{R}e(\lambda) > 0$ ;
- ▶ If  $\mathcal{G}$  is strongly connected, then  $0$  is simple and  $\bar{\pi}_i > 0$  for all  $i$ ;
- ▶  $L'\bar{y} = 0 \Leftrightarrow P'(D\bar{y}) = (D\bar{y})$ ;

$\bar{\pi}$  Laplace-invariant distribution


# Who is the most central node?



'Lorenzo de' Medici'  
G. Vasari, 1534

## Network centralities


$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$z_i \geq 0 \text{ centrality of node } i$$

# Degree centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




$$z_i = w_i^- \text{ in-degree of node } i \text{ *degree centrality*}$$

Example: number of citations of an article, of followers on Twitter

## Degree centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




The two nodes with the highest degree centrality.



# Eigenvector centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




**Drawback** of degree-centrality: all incoming links are considered the same.

A different approach:  $z_i \propto \sum_{j \in N_i^-} z_j$

# Eigenvector centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




**Drawback** of degree-centrality: all incoming links are considered the same.

A different approach: 
$$z_i \propto \sum_{j \in N_i^-} z_j = \sum_j W_{ji} z_j$$

# Eigenvector centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




**Drawback** of degree-centrality: all incoming links are considered the same.

A different approach:  $z_i \propto \sum_{j \in N_i^-} z_j = \sum_j W_{ji} z_j$

$\lambda z = W'z$  with  $\lambda = \lambda_W$  *eigenvector centrality*

# Eigenvector centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



$$\lambda z = W'z \text{ with } \lambda = \lambda_W$$


Remark: If all nodes have the same in-degree:  $w_i^- = \delta$  for all  $i$

$$W'\mathbb{1} = \delta\mathbb{1} \Rightarrow z = \mathbb{1}$$

**Drawback:** node  $j$  contributes proportional to its out-degree  $w_j$

## Invariant distribution centrality


$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



Another centrality measure:  $z_i \propto \sum_{j \in N_i^-} \frac{1}{w_j} z_j$

# Invariant distribution centrality


$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



Another centrality measure: 
$$z_i \propto \sum_{j \in N_i^-} \frac{1}{w_j} z_j = \sum_j P_{ji} z_j$$

# Invariant distribution centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




Another centrality measure: 
$$z_i \propto \sum_{j \in N_i^-} \frac{1}{w_j} z_j = \sum_j P_{ji} z_j$$

$$z = P' z \Rightarrow z = \pi \text{ invariant distribution centrality}$$

# Invariant distribution centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




Another centrality measure:  $z_i \propto \sum_{j \in N_i^-} \frac{1}{w_j} z_j = \sum_j P_{ji} z_j$

$z = P'z \Rightarrow z = \pi$  *invariant distribution centrality*

If  $\mathcal{G}$  is balanced,  $\pi = w = w^- \Rightarrow \text{inv. dist. centr.} = \text{deg centr.}$



# A comparison of the various centralities




	Deg	Eig	Inv. dist.
1	0.0345	0.0348	0.0313
2	0.0517	0.0581	0.0451
3	0.0517	0.0664	0.0613
4	0.0517	0.0689	0.0680
5	0.0517	0.0680	0.0869
6	0.0517	0.0430	0.0490


7	0.0690	0.0678	0.0514
8	0.0517	0.0661	0.0444
9	0.0517	0.0659	0.0491
10	0.0517	0.0627	0.0761
11	0.0345	0.0226	0.0324
12	0.0345	0.0215	0.0240
13	0.0517	0.0399	0.0317
14	0.0690	0.0640	0.0548
15	0.0517	0.0613	0.0464
16	0.0517	0.0484	0.0817
17	0.0517	0.0225	0.0481
18	0.0345	0.0215	0.0240
19	0.0345	0.0307	0.0300
20	0.0517	0.0492	0.0441
21	0.0172	0.0166	0.0204

# A comparison of the various centralities

Eigenvalue centrality



Invariant distribution centrality




## Drawbacks of eigenvalue and invariant distribution centr.

In general, they are not uniquely defined if the graph has more than one sink component

if the graph has just one sink component, only nodes in that component have non zero centrality

# Katz centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




A convex comb. of *network centrality* and *intrinsic centrality*

$$z = \frac{1 - \beta}{\lambda_W} W' z + \beta \mu$$

$$z = (I - (1 - \beta) \lambda_W^{-1} W')^{-1} \beta \mu \quad \text{Katz centrality}$$

# Bonacich centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$




A convex comb. of *network centrality* and *intrinsic centrality*

$$z = (1 - \beta)P'z + \beta\mu$$

$$z = (I - (1 - \beta)P')^{-1}\beta\mu \quad \text{Bonacich centrality}$$

# Bonacich centrality

$$\mathcal{G} = (\mathcal{V}, \mathcal{E}, W)$$



A convex comb. of *network centrality* and *intrinsic centrality*

$$z = (1 - \beta)P'z + \beta\mu$$

$$z = (I - (1 - \beta)P')^{-1}\beta\mu \quad \text{Bonacich centrality}$$

Page-rank (Google):  $\mu = n^{-1}\mathbb{1}$ ,  $\beta \sim 0.15$

## The structure of Bonacich centrality


$$z = (I - (1 - \beta)P')^{-1}\beta\mu$$

$$z_i = \beta \sum_{k=0}^{\infty} (1 - \beta)^k \sum_j \mu_j (P^k)_{ji}$$

# The structure of Bonacich centrality

$$z = (I - (1 - \beta)P')^{-1}\beta\mu$$

$$z_i = \beta \sum_{k=0}^{\infty} (1 - \beta)^k \sum_j \mu_j (P^k)_{ji}$$




$$= \beta [\mu_i + (1 - \beta)(\mu_{j_1} P_{j_1 i} + \mu_{j_2} P_{j_2 i}) + (1 - \beta)^2 (\mu_{j_3} P_{j_3 j_1} P_{j_1 i} + \mu_{j_4} P_{j_4 j_2} P_{j_2 i} + \mu_{j_5} P_{j_5 j_2} P_{j_2 i})] + \dots$$



# The structure of Bonacich centrality

$$z = (I - (1 - \beta)P')^{-1}\beta\mu$$

$$z_i = \beta \sum_{k=0}^{\infty} (1 - \beta)^k \sum_j \mu_j (P^k)_{ji}$$




$$= \beta [\mu_i + (1 - \beta)(\mu_{j_1} P_{j_1 i} + \mu_{j_2} P_{j_2 i}) + (1 - \beta)^2 (\mu_{j_3} P_{j_3 j_1} P_{j_1 i} + \mu_{j_4} P_{j_4 j_2} P_{j_2 i} + \mu_{j_5} P_{j_5 j_2} P_{j_2 i})] + \dots$$

# The structure of Bonacich centrality

$$z = (I - (1 - \beta)P')^{-1}\beta\mu$$

$$z_i = \beta \sum_{k=0}^{\infty} (1 - \beta)^k \sum_j \mu_j (P^k)_{ji}$$



$$= \beta [\mu_i + (1 - \beta)(\mu_{j_1} P_{j_1 i} + \mu_{j_2} P_{j_2 i}) + (1 - \beta)^2 (\mu_{j_3} P_{j_3 j_1} P_{j_1 i} + \mu_{j_4} P_{j_4 j_2} P_{j_2 i} + \mu_{j_5} P_{j_5 j_2} P_{j_2 i})] + \dots$$

- ▶ The Bonacich centrality is uniquely defined for every graph  $\mathcal{G}$
- ▶  $\beta = 0 \Rightarrow$  Intrinsic centrality (no network)
- ▶  $\beta \rightarrow 1 \Rightarrow$  Invariant distribution centrality (only network)

# The various centralities

Eigenvector centrality

$$z = \frac{1}{\lambda_W} W' z$$

Invariant distribution

$$z = P' z$$

Katz centrality

$$z = \frac{1 - \beta}{\lambda_W} W' z + \beta \mu$$

Bonacich centrality

$$z = (1 - \beta) P' z + \beta \mu$$

## Some references

- ▶ L. Katz; A new status index derived from sociometric analysis, *Psychometrika*, 18, 3–43, 1953.
- ▶ P. Bonacich; Power and Centrality: A Family of Measures, *American Journal of Sociology*, 1987.