

Enhancing Solar Panel Maintenance: A CNN-GAN Approach for Dust Detection and Classification

Shiva Mehta

Centre for Research Impact &
Outcome, Chitkara University Institute
of Engineering and Technology
Chitkara University
Punjab, India
shiva.2387@chitkara.edu.in

Rajat Saini

Chitkara Centre for Research and
Development,
Chitkara University, Himachal
Pradesh, 174103, India
rajat.saini.orp@chitkara.edu.in

Abstract— The problem of dust settlement on solar panels is vital, likely decreasing their performance by 30%, which negatively affects energy yield and economic viability. Conventional dust detection approaches, such as optical sensors and basic image processing, frequently do not perform well in different environmental conditions. This investigation presents a pioneering hybrid framework that combines Convolutional Neural Networks (CNN) with Generative Adversarial Networks (GAN) to increase the detection and classification accuracy of dust on solar panels. CNN-GAN makes use of CNN's extraction of features and the data augmentation talents of GAN to create a sturdy structure that generalizes widely across multiple situations. The model was trained and assessed using an extensive dataset of 30,000 photos, divided into three categories: There is no Dust, some Light Dust, and a lot of Heavy Dust. This proposed model exceeded traditional models, including VGG-16 (accuracy of 89.2%) and InceptionV3 (accuracy of 90.5%), by achieving an accuracy of 95.4%. The CNN-GAN model suggested that accuracy stood at 96.2%, recall reached 94.7%, and the F1-score was an outstanding 95.4%. The Area Under the ROC Curve (AUC) was 0.97, implicating a powerful capability for discrimination. Investigations performed qualitatively indicated that pictures produced by GANs enhance CNN's educational process. The study points out the capability of the CNN-GAN model for immediate monitoring and automatic care of solar panels, ensuring maximum performance and energy efficiency. This research will also focus on expanding this methodology to reveal a larger range of abnormalities and implementing it into IoT devices for proactive maintenance solutions.

Keywords: CNN-GAN, solar panels, dust detection, image analysis, classification, Convolutional Neural Networks (CNN), Generative Adversarial Networks (GAN), Data Augmentation, Deep Learning.

I. INTRODUCTION

With a growing concern towards renewable energy, solar panels are being incorporated globally in energy-saving strategies at a faster pace. Sunlight is abundant and renewable, making it critical to wean the world's population off fossil fuel-based energy and arrest climate change [1]. The failure rates of solar panels vary due to different factors, such as environmental conditions where dust accumulation is of great concern [2]. The dust accumulates on the surface of the solar panels, reducing the amount of light that is reflected to the panel and leading to as much as 30% loss in energy conversion. Such a decline in efficiency not only affects the efficiency of individual panels but also has significant consequences on energy production and the economy, especially in regions with high dust density [3]. It has been established that proper maintenance of solar panels, especially cleaning, is very important if they are to

perform as expected. Current methods of detecting dust, such as snapshot inspection and simple image processing methods, are slow, time-consuming and unreliable because of their inability to compensate for the amount or type of light incident on the object's surface and the surrounding environment. Automated dust detection has been explored using Convolutional Neural Networks (CNNs) to some level of success [4]. However, CNNs could be faced with problems of generalization and how accurate they will be in IOT conditions, which are very complicated conditions like changes in dust densities, varieties of lighting angles, and other forms of occlusions. This study proposes a new model that combines the benefits of CNN and GAN to overcome these shortcomings [5]. The CNN-GAN combines CNN's powerful feature extractor in dust detection with GAN's unique ability to generate realistic fake pictures, therefore enhancing the model's robustness in different environments. The suggested strategy is to enhance the level of dust identification on solar panels to maximize the utilization of solar energy by keeping the surface clean. One of the typical problems found throughout the world is dust, which affects panels' efficiency and maximum energy production. Traditional approaches for detecting and classifying dust, including visual examination and simple image processing techniques, may be insufficient at times because of their inability to adapt to changes in ambient conditions, including lighting conditions, the orientation of the object being examined, and dust concentration ratios [6]. Conventional machine learning techniques like CNNs show improvements in performance; still, they lack versatility. This constraint is quite a challenge to the maintenance and the real-time monitoring of the solar panels, which is very crucial to ensure they work to the optimal level and for the intended longer life span. This work proposes a new CNN-GAN model that can enhance the detection and classification of dust on solar panels. In addition to this, the CNN-GAN model capitalizes on CNN's ability to perform feature extraction while using the GAN to create fake pictures. This study makes three key contributions: Among them, I propose (1) a novel CNN-GAN designed to provide significantly improved dust detection performance over standard CNN models, (2) a large database of various dust conditions and environmental factors for training and testing the model, and (3) a comprehensive analysis of the proposed method's applicability in various real-world scenarios. These contributions create the basis for the implementation of an accurate, self-sustaining dust-detecting system that can be integrated into the existing IoT-based smart grid to provide predictive solar panel maintenance, thus maximising energy yield and reducing operating costs [7].

II. LITERATURE REVIEW

2.1 Dust Detection Techniques

Accumulating dust is ordinarily seen as a significant element that affects the performance of solar panels. A variety of methodologies exist to pinpoint and quantify dust deposition. Optical sensors have been in use for a long time and operate by assessing the reduction in light transmission that is caused by dust [8]. When optical sensor technology is used in places where the weather is not controlled or where brightness is variable, inaccuracies can arise. Investigations into infrared imaging have involved looking into heat differences between dust-free and dusty zones to discover dust collection [9]. This method requires complicated facilities and is vulnerable to variations in external temperature, making it unsuitable for ordinary use. Machine learning has become adept at utilizing algorithms to classify dust conditions according to their individual visual properties [10]. For this reason, the methods comprise Support Vector Machines (SVM), Decision Trees, and k-nearest Neighbours (k-NN). Even though they allow for greater user adaptability in comparison to traditional methods, their effectiveness is heavily tied to feature engineering and may prove to be less than ideal in different environmental situations. The limitations in these methodologies indicate the urgent necessity for models that are designed to learn from various situations with restricted human input [7].

2.2 Deep Learning in Dust Detection

Convolutional Neural Networks (CNN), especially in the domain of deep learning, have captured interest due to their proficiency in detecting sophisticated features from data on images. CNNs, when left to their own devices, can detect patterns and abnormalities in photos, which cuts the need for manual feature identification. According to research, artificial neural networks have demonstrated their effectiveness in grading dust concentrations and pinpointing surface irregularities on solar panels [11]. Evaluations indicate that CNN models have been established to be effective in classifying dust patterns with premium accuracy, applying specific classifications. Distinct from these achievements, the ongoing complications regarding CNNs' generalization are caused by their requirement for abundant and diverse training data. Changes can notably influence the effectiveness of these mitigation techniques in illumination, the angles of camera capture, and variations in dust properties. When it comes to practical uses, this constraint is particularly difficult due to the considerably fluctuating ambient conditions that solar panels encounter. Therefore, the necessity of improving the resilience and flexibility of CNN models is growing to ensure trustworthy dust detection in a variety of contexts [12].

2.3 Generative Adversarial Networks (GAN) for Data Augmentation

The development of Generative Adversarial Networks (GANs) has resulted in a useful tool for data augmentation under circumstances where high-quality and diverse data is difficult to come by. GANs include two networks: a

generator that creates synthetic data and a discriminator for assessing the authenticity of the data it creates [13]. The competitive interactions between these two networks produce synthetic visuals with an exceptionally authentic look. In the realm of solar panel dust-detection tech, GANs (Generative Adversarial Networks) enable the development of diverse spectra of images that show how dust impacts mimic real circumstances, enhancing the training dataset. Such supplementary data escalates CNN's ability to acquire a wide variety of characteristics, in turn advancing its ability to generalize in several scenarios [14]. GAN, paired with CNN, intends to improve feature learning and significantly reduce the obstacles triggered by data scarcity and imbalance, hindering the effective training of deep learning models for dust detection [15]. Researchers have demonstrated how GANs can create synthetic data, improving model resistance and illustrating their potential for improved dust detection systems. Ultimately, though conventional dust detection techniques and CNN-based approaches have formed a solid basis for automated surveillance, they need to address the difficulties associated with varied real-world situations. The application of GANs can bring stability and enhance the performance of solar panel dust detection platforms by enriching the data, thereby overcoming current challenges. This exploration underscores the capabilities of a hybrid technique that blends CNN-GAN methodologies in order to further dust detection technology for the improvement of solar energy system maintenance [16].

III. METHODOLOGY

This part provides a concise step-by-step process of developing and validating a CNN-GAN model that can detect and classify dust on solar panels. The process is organized into four primary components: Data collection, CNN structure development, generator and discriminator structure, and learning and assessment stages, as shown in Figure 1.




Fig1. Phases in the Process Approach

A. Data Information, Augmentation

To perform this investigation, a thorough assortment of solar panel photographs was brought together, encompassing a range of climatic conditions and different dust levels. Various lighting situations, camera angles, and

conditions related to weather are vital for producing a selection of high-quality photographs. The photos were labelled and classified into three separate categories: Dust-Free: It is paramount to keep solar panels dust and grim-free. Light Dust: Panels which reveal just enough dust accumulation to slightly decrease clarity without entirely blocking sunlight from getting through. Severe Dust Accumulation: Panels that are dusty due to a thick buildup, hindering sunlight penetration substantially and resulting in reduced panel efficiency, as shown in the table 1.

TABLE 1: DATASET DISTRIBUTION

Category	Original Images	Augmented Images	Total Images
No Dust	1,500	8,500	10,000
Light Dust	1,500	8,500	10,000
Heavy Dust	1,500	8,500	10,000
Total	4,500	25,500	30,000

To enhance the dataset and increase model generalization, several data augmentation strategies were employed: Rotation: Images were rotated to different angles that varied from -45 to +45 degrees. Scaling: Sizes of images were decreased or increased by a factor ranging between 0.8 and 1.2. Brightness Modifications: The intensity of photos was randomly changed to resemble many lighting conditions. This generated a well-balanced dataset comprising 30,000 images, involving the production of 10,000 photographs for each category, as shown in Figure 2.




Fig 2. Various Solar Panels Dust Detection Images

B. Feature Extraction using CNN and GAN

The design of the CNN-GAN model aspires to make use of the advantages of Convolutional Neural Networks (CNN) together with Generative Adversarial Networks (GAN). The architecture has two primary components: This feature extractor receives training with the ResNet-50 model prior to training. The model is refined using the solar panel dataset to extract the unique characteristics of dust

collection. The information induced by the final convolutional layer is flattened and fed through fully connected layers to classify the images according to one of three dust classifications. ResNet-50 model architecture is what the feature extractor for CNN uses. The Generative Adversarial Network (GAN) comprises two neural networks: a generator and a discriminator, both trained within a competitive environment. The generator network is conditioned to generate convincing images of solar panels demonstrating a variety of dust patterns, whereas the discriminator network is taught to identify both genuine and fabricated photographs. Generator Network: Engages random noise as input and generates images of solar panels filled with dust. Discriminator Network: This site accepts photos that are either real or artificial and categorizes them as authentic or not genuine. Adversarial training strengthens the feature extractor of CNN by letting it take knowledge from the images it produces, as shown in Figure 3.




Fig 3 . CNN Architecture Images.

C. Model Evaluation

The loss function of the CNN-GAN model is training it using a union of adversarial loss generated by the GAN along with classification loss arising from the CNN. The training procedure occurs in two stages: The generator and discriminator perform concurrent training in a minimax framework until the generator generates synthetic pictures of high quality.

TABLE 2: MODEL TRAINING PARAMETERS

Parameter	Value
Learning Rate	0.001
Batch Size	64
Number of Epochs	50
Optimizer	Adam
Loss Function (CNN)	Cross-Entropy
Loss Function (GAN)	Binary Cross-Entropy
λ lambda λ (Adversarial Loss Weight)	0.5

The generator looks to minimize the discriminator's loss in tandem with the discriminator's intention to refine its ability to differentiate between real and fake images. In order to measure the effectiveness of the model, we apply

metrics of classical classification, comprising accuracy, precision, recall, F1-score, and the area under the ROC curve (AUC). The model's resilience is assessed by applying it to unfamiliar images in a range of environmental situations. Findings show that the CNN-GAN model greatly outperforms traditional strategies for both accuracy and resilience across different scenarios. This structure furnishes the groundwork for automatic, real-time surveillance of solar panel health, likely resulting in more efficient maintenance and greater energy generation, as shown in Figure 5.




Fig4 . Training Validation Loss and Accuracy Images.

IV. RESULTS

The suggested CNN-GAN architecture produced important progress over traditional models in recognizing and categorizing dust on solar panels. This model reached an overall accuracy of 95.4%, a precision of 96.2%, a recall of 94.7%, along with an F1-score of 95.4%. These metrics demonstrate the model's proficiency in differentiating among the three dust classifications: No Dust, Moderate Dust, and Widespread Dust. The model revealed strong abilities in properly defining dust levels, qualifying with an Area Under the ROC Curve (AUC) of 0.97.




Fig5 . Detection Accuracy per Dust Type

The higher AUC value confirms the model's effectiveness and durability, especially in response to changing environmental conditions that frequently pose hardships for conventional models. The boost in diversity of the training set came from the GAN component in the model, which trained to create visuals of solar panels covered in

differing levels of dust. A visual assessment of the produced images indicated their detailed quality and realism, helping CNN secure a broader and more representative spectrum of dust deposition attributes.




Fig6 . Confusion Matrix.

This model showed remarkable capability across diverse illumination circumstances, angles, and dust arrangements. It extensively exceeded traditional CNN-based systems, which often suffered from problems related to generalization and adaptation in fresh settings. The CNN-GAN model adopted the generative properties of GANs, which improved feature learning and enhanced performance in practical environments. The study performed a comparison by using models at baseline, including VGG-16, InceptionV3, and solo CNN, to evaluate the performance of the suggested CNN-GAN model.




Fig7 . Detection Accuracy per Dust Type

The resulting data is presented in the following table, along with various performance graphs. The ROC curve illustrates the true positive rate related to the false positive rate for both the CNN-GAN model and a comparison model. The CNN-GAN model, seen as the light blue line, attains an AUC value of 0.97, greatly superior to the AUC of 0.92 (represented by the red, dashed line) for the baseline model; this reflects a rise in both models regarding effectiveness in understanding dust classifications. The confusion matrix illustrates the efficacy of the CNN-GAN model in accurately categorizing pictures into three

classifications: No Dust, some Light Dust, and Much Heavy Dust. In all the categories, the model exhibited strong performance, with the largest misclassifications taking place between Light Dust and Heavy Dust, which look alike. The bar chart shows the performance measures of a number of models. The CNN-GAN model repeatedly outshone the baseline models on all criteria, proving the viability of the combination of CNN and GAN for the detection of dust on solar panels.




Fig8 . Dust Type Detection Distribution.

V. CONCLUSION

In this paper, an innovative CNN-GAN model is presented for better detection and classification of dust on solar panels, solving the flaws found in typical methods and independent CNN models. The recommended method incorporates the strong feature extraction skills of Convolutional Neural Networks (CNN) combined with the data augmentation features of Generative Adversarial Networks (GAN), thereby attaining resilient performance in different environmental Setups. Accuracy at 95.4% shows that the CNN-GAN model is superior to VGG-16 (89.2%) and InceptionV3 (90.5%) baseline models. Furthermore, the model exhibited exceptional accuracy, recall, and F1-scores of 96.2%, 94.7%, and 95.4%, respectively, underscoring its efficacy in differentiating among the three dust classifications: No Dust, just Light Dust, and substantial Heavy Dust. The Area Under the Curve (AUC) score of 0.97 points out the outstanding discriminative potential of the model. The GAN-produced images in the qualitative study confirmed their realism, enabling CNN to improve feature learning and generalization. The proposed model exhibited robustness by maintaining exceptional performance under a variety of illumination conditions and dust levels, exceeding the generalization problems faced by individual CNN models. By using this merged detection system, the reliability of dust detection systems increases, and a solid footing is developed for real-time monitoring solutions in solar panel maintenance, thus enhancing energy efficiency and lowering operating costs. Subsequent research goals

include expanding this method to identify anomaly categories further and their integration with IoT-enabled smart grids for automated, preventative maintenance.

REFERENCES

- [1] R. Al-Mashhadani *et al.*, "Deep learning methods for solar fault detection and classification: a review," *Sol. cells*, vol. 11, p. 12, 2021.
- [2] E. Jiménez-Delgado, A. Méndez-Porras, and J. Alfaro-Velasco, "Generative Adversarial Network for Synthetic Imaging Data of Sub-optimal Conditions in Photovoltaic Panels," in *Information Technology and Systems*, 2022, pp. 206–218.
- [3] R. F. Pamungkas, I. B. K. Y. Utama, and Y. M. Jang, "A Novel Approach for Efficient Solar Panel Fault Classification Using Coupled UDenseNet," *Sensors*, vol. 23, no. 10, 2023.
- [4] G. Tanda and M. Migliazzi, "Infrared thermography monitoring of solar photovoltaic systems: A comparison between UAV and aircraft remote sensing platforms," *Therm. Sci. Eng. Prog.*, vol. 48, p. 102379, 2024.
- [5] C. Yang *et al.*, "A Survey of Photovoltaic Panel Overlay and Fault Detection Methods," *Energies*, vol. 17, no. 4, 2024, doi: 10.3390/en17040837.
- [6] S. Oh, B. W. Figgis, and S. Rashkeev, "Effects of thermophoresis on dust accumulation on solar panels," *Sol. Energy*, vol. 211, pp. 412–417, 2020.
- [7] R. J. Mustafa, M. R. Goma, M. Al-Dhaifallah, and H. Rezk, "Environmental Impacts on the Performance of Solar Photovoltaic Systems," *Sustainability*, vol. 12, no. 2, 2020, doi: 10.3390/su12020608.
- [8] F. Mejia, J. Kleissl, and J. L. Bosch, "The Effect of Dust on Solar Photovoltaic Systems," *Energy Procedia*, vol. 49, pp. 2370–2376, 2014.
- [9] S. Mehta and D. Kundra, "Improving Drone Operational Efficiency through IoT-Blockchain Integration," in *2023 4th International Conference on Intelligent Technologies (CONIT)*, 2024, pp. 1–5.
- [10] V. Bhardwaj, S. Bala, V. Kadyan, and V. Kukreja, "Development of Robust Automatic Speech Recognition System for Children's using Kaldi Toolkit," in *2020 Second International Conference on Inventive Research in Computing Applications (ICIRCA)*, 2020, pp. 10–13.
- [11] A. Suryavanshi, S. Mehta, S. Chattopadhyay, and M. Aeri, "Navigating Water Scarcity with IoT: A Smart Management System Approach," in *2023 4th International Conference on Intelligent Technologies (CONIT)*, 2024, pp. 1–5.
- [12] H. Lu, B. He, and W. Zhao, "Experimental study on the super-hydrophobic coating performance for solar photovoltaic modules at different wind directions," *Sol. Energy*, vol. 249, pp. 725–733, 2023.
- [13] Sakshi, V. Kukreja, and S. Ahuja, "Recognition and classification of mathematical expressions using machine learning and deep learning methods," in *9th International Conference on Reliability, Infocom Technologies and Optimization (Trends and Future Directions) (ICRITO)*, 2021, pp. 1–5.
- [14] M. Katoch, K. Kumar, and V. Dahiya, "Dust accumulation and reduction in electrical performance of solar PV panels," *Mater. Today Proc.*, vol. 46, pp. 6608–6612, 2021.
- [15] R. K. Saini *et al.*, "Effects of dust on the performance of solar panels – a review update from 2015–2020," *Energy & Environ.*, vol. 34, no. 6, pp. 2110–2162, 2023.
- [16] J. Pedro Lemos Morais, L. Antônio Barros Herênio Madeira, F. Perin Gasparin, A. Krenzinger, F. Mendes, and A. Angélica Mathias Macêdo, "Analysis of mesh screens for outdoor i-v curve measurements of pv modules at several irradiance levels," *Sol. Energy*, vol. 249, pp. 152–162, 2023.