

An Introduction to Quantum Computing

Lecture 16:

Density Operators

Paolo Zuliani

Dipartimento di Informatica
Università di Roma “La Sapienza”, Rome, Italy



SAPIENZA
UNIVERSITÀ DI ROMA

Agenda

- Mixed states and traces
- Density matrices
- The rules of Quantum Mechanics with density matrices
- The von Neumann Entropy and the Holevo bound

Definition

A ket $|\nu\rangle \in \mathcal{H}$ is a *pure state*.

Mixed States

Definition

A ket $|v\rangle \in \mathcal{H}$ is a *pure state*.

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state* ρ is the set:

$$\rho = \{|v_i\rangle, w_i\}_{i=1}^N$$

A mixed state tells us that our quantum system is in pure state $|v_i\rangle$ with ‘classical’ probability w_i .

Rule 3: when measuring observable A on (pure) state $|\nu\rangle$ we have that

$$\text{Prob}(A = \lambda_i; |\nu\rangle) = \langle \nu | P_i | \nu \rangle \quad (\text{equivalently } \langle A \rangle_\nu = \langle \nu | A | \nu \rangle)$$

where P_i is the projector associated to the i -th eigenspace of A .

Rule 3: when measuring observable A on (pure) state $|v\rangle$ we have that

$$\text{Prob}(A = \lambda_i; |v\rangle) = \langle v | P_i | v \rangle \quad (\text{equivalently } \langle A \rangle_v = \langle v | A | v \rangle)$$

where P_i is the projector associated to the i -th eigenspace of A .

How about measuring A on a mixed state $\rho = \{|v_i\rangle, w_i\}_{i=1}^N$?

Rule 3: when measuring observable A on (pure) state $|\nu\rangle$ we have that

$$\text{Prob}(A = \lambda_i; |\nu\rangle) = \langle \nu | P_i | \nu \rangle \quad (\text{equivalently } \langle A \rangle_\nu = \langle \nu | A | \nu \rangle)$$

where P_i is the projector associated to the i -th eigenspace of A .

How about measuring A on a mixed state $\rho = \{|\nu_i\rangle, w_i\}_{i=1}^N$?

Assuming the quantum and classical probabilities are independent, we get

$$\text{Prob}(A = \lambda_i; \rho) = \sum_{j=1}^N w_j \text{Prob}(A = \lambda_i; |\nu_j\rangle) = \sum_{j=1}^N w_j \langle \nu_j | P_i | \nu_j \rangle$$

Trace of a Linear Operator

Definition

Let $\{e_i\}$ be an orthonormal basis for an Hilbert space \mathcal{H} , and A be an operator on \mathcal{H} . The *trace* of A is defined

$$\text{tr}(A) = \sum_i \langle e_i | A e_i \rangle$$

Trace of a Linear Operator

Definition

Let $\{e_i\}$ be an orthonormal basis for an Hilbert space \mathcal{H} , and A be an operator on \mathcal{H} . The *trace* of A is defined

$$\text{tr}(A) = \sum_i \langle e_i | A e_i \rangle$$

Note: it is a generalisation of the usual trace of a matrix.

Exercise: show that $\text{tr}(A)$ is independent of the basis chosen (hence it is well defined!)

Trace of a Linear Operator

Definition

Let $\{e_i\}$ be an orthonormal basis for an Hilbert space \mathcal{H} , and A be an operator on \mathcal{H} . The *trace* of A is defined

$$\text{tr}(A) = \sum_i \langle e_i | A e_i \rangle$$

Note: it is a generalisation of the usual trace of a matrix.

Exercise: show that $\text{tr}(A)$ is independent of the basis chosen (hence it is well defined!)

Let us rework Rule 3:

$$\langle v | A v \rangle = \langle v | A \sum_i |e_i\rangle\langle e_i| v \rangle = \sum_i \langle v | A |e_i\rangle\langle e_i| v \rangle$$

Trace of a Linear Operator

Definition

Let $\{e_i\}$ be an orthonormal basis for an Hilbert space \mathcal{H} , and A be an operator on \mathcal{H} . The *trace* of A is defined

$$\text{tr}(A) = \sum_i \langle e_i | A e_i \rangle$$

Note: it is a generalisation of the usual trace of a matrix.

Exercise: show that $\text{tr}(A)$ is independent of the basis chosen (hence it is well defined!)

Let us rework Rule 3:

$$\begin{aligned} \langle v | A v \rangle &= \langle v | A \sum_i |e_i\rangle\langle e_i| v \rangle = \sum_i \langle v | A |e_i\rangle\langle e_i| v \rangle \\ &= \sum_i \langle e_i | v \rangle \langle v | A e_i \rangle = \sum_i \langle e_i | P_v A e_i \rangle = \text{tr}(P_v A) \end{aligned}$$

where $P_v = |v\rangle\langle v|$.

Mixed State Operator

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state operator* ρ is defined:

$$\rho = \sum_i w_i P_{v_i} = \sum_i w_i |v_i\rangle\langle v_i|$$

Mixed State Operator

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state operator* ρ is defined:

$$\rho = \sum_i w_i P_{v_i} = \sum_i w_i |v_i\rangle\langle v_i|$$

Let us now compute the probability of measuring λ_i on the mixed state ρ :

$$\text{Prob}(A = \lambda_i; \rho) = \sum_{j=1} w_j \langle v_j | P_i | v_j \rangle$$

Mixed State Operator

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state operator* ρ is defined:

$$\rho = \sum_i w_i P_{v_i} = \sum_i w_i |v_i\rangle\langle v_i|$$

Let us now compute the probability of measuring λ_i on the mixed state ρ :

$$\text{Prob}(A = \lambda_i; \rho) = \sum_{j=1} w_j \langle v_j | P_i | v_j \rangle = \sum_j w_j \text{tr}(P_{v_j} P_i)$$

Mixed State Operator

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state operator* ρ is defined:

$$\rho = \sum_i w_i P_{v_i} = \sum_i w_i |v_i\rangle\langle v_i|$$

Let us now compute the probability of measuring λ_i on the mixed state ρ :

$$\begin{aligned} \text{Prob}(A = \lambda_i; \rho) &= \sum_{j=1} w_j \langle v_j | P_i | v_j \rangle = \sum_j w_j \text{tr}(P_{v_j} P_i) \\ &= \text{tr}\left(\sum_j w_j P_{v_j} P_i\right) \end{aligned}$$

Mixed State Operator

Definition

Let $\{|v_1\rangle, \dots, |v_N\rangle\}$ be pure states and w_1, \dots, w_N be positive numbers such that $\sum_i w_i = 1$. The *mixed state operator* ρ is defined:

$$\rho = \sum_i w_i P_{v_i} = \sum_i w_i |v_i\rangle\langle v_i|$$

Let us now compute the probability of measuring λ_i on the mixed state ρ :

$$\begin{aligned} \text{Prob}(A = \lambda_i; \rho) &= \sum_{j=1} w_j \langle v_j | P_i | v_j \rangle = \sum_j w_j \text{tr}(P_{v_j} P_i) \\ &= \text{tr}\left(\sum_j w_j P_{v_j} P_i\right) \\ &= \text{tr}(\rho P_i) \end{aligned}$$

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Let us show that a positive operator ρ is self-adjoint:

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Let us show that a positive operator ρ is self-adjoint:

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$$

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Let us show that a positive operator ρ is self-adjoint:

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0 \quad \implies$$

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle = \langle v | \rho v \rangle^* = \langle \rho v | v \rangle = \langle \rho^\dagger v | v \rangle$$

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Let us show that a positive operator ρ is self-adjoint:

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0 \quad \implies$$

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle = \langle v | \rho v \rangle^* = \langle \rho v | v \rangle = \langle \rho^\dagger v | v \rangle \quad \implies$$

$$\forall v \in \mathcal{H}, \langle \rho v | v \rangle = \langle \rho^\dagger v | v \rangle$$

Density Matrices

Definition

A *density matrix* is an operator ρ satisfying:

- ρ is positive (i.e., $\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0$);
- $\text{tr}(\rho) = 1$;
- $[\rho = \rho^\dagger]$ (but this follows from positivity).

Let us show that a positive operator ρ is self-adjoint:

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle \geq 0 \quad \implies$$

$$\forall v \in \mathcal{H}, \langle v | \rho v \rangle = \langle v | \rho v \rangle^* = \langle \rho v | v \rangle = \langle \rho^\dagger v | v \rangle \quad \implies$$

$$\forall v \in \mathcal{H}, \langle \rho v | v \rangle = \langle \rho^\dagger v | v \rangle \quad \implies \text{(show that } \forall z, \langle x | z \rangle = \langle y | z \rangle \implies x = y)$$

$$\forall v \in \mathcal{H}, \rho v = \rho^\dagger v$$

Density Matrices

Exercise: prove that a mixed state operator is a density matrix.

Density Matrices

Exercise: prove that a mixed state operator is a density matrix.

Any density matrix ρ is self-adjoint, hence can apply the spectral theorem and write:

$$\rho = \sum_i y_i P_{\psi_i}$$

where the $|\psi_i\rangle$'s are an orthonormal basis and the y_i 's are probabilities summing to 1.

Density Matrices

Exercise: prove that a mixed state operator is a density matrix.

Any density matrix ρ is self-adjoint, hence can apply the spectral theorem and write:

$$\rho = \sum_i y_i P_{\psi_i}$$

where the $|\psi_i\rangle$'s are an orthonormal basis and the y_i 's are probabilities summing to 1.

Note that orthonormality is not required when modelling classically weighted sum of states (*i.e.*, when the system is in state $|\psi_i\rangle$ with probability y_i).

Density Matrices

A density matrix does NOT admit a unique decomposition as a mixed state operator!

Density Matrices

A density matrix does NOT admit a unique decomposition as a mixed state operator!

Example:

Consider density matrix $\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. It is easy to show that:

$$\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |1\rangle\langle 1|$$

which suggests that our system is in pure state $|0\rangle$ with (classical) probability 50% and in pure state $|1\rangle$ again with (classical) probability 50%.

Density Matrices

A density matrix does NOT admit a unique decomposition as a mixed state operator!

Example:

Consider density matrix $\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. It is easy to show that:

$$\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |1\rangle\langle 1|$$

which suggests that our system is in pure state $|0\rangle$ with (classical) probability 50% and in pure state $|1\rangle$ again with (classical) probability 50%. However:

$$\rho = \frac{1}{2} \left[\frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right] + \frac{1}{2} \left[\frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right]$$

Density Matrices

A density matrix does NOT admit a unique decomposition as a mixed state operator!

Example:

Consider density matrix $\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. It is easy to show that:

$$\rho = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |1\rangle\langle 1|$$

which suggests that our system is in pure state $|0\rangle$ with (classical) probability 50% and in pure state $|1\rangle$ again with (classical) probability 50%. However:

$$\rho = \frac{1}{2} \left[\frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \right] + \frac{1}{2} \left[\frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \right]$$

Exercise: verify that the two matrices above are projectors (recall a projector P satisfies $P^2 = P$ and $P = P^\dagger$).

Proposition

The set of density matrices is convex, i.e., given density matrices ρ_1, \dots, ρ_N and positive numbers w_1, \dots, w_N such that $\sum_i w_i = 1$, then $\sum_i w_i \rho_i$ is a density matrix.

Exercise: prove the proposition above.

Proposition

The set of density matrices is convex, i.e., given density matrices ρ_1, \dots, ρ_N and positive numbers w_1, \dots, w_N such that $\sum_i w_i = 1$, then $\sum_i w_i \rho_i$ is a density matrix.

Exercise: prove the proposition above.

Density matrices of the form $|\psi\rangle\langle\psi|$ are called *pure states*, the rest are *mixed*.

Proposition

The set of density matrices is convex, i.e., given density matrices ρ_1, \dots, ρ_N and positive numbers w_1, \dots, w_N such that $\sum_i w_i = 1$, then $\sum_i w_i \rho_i$ is a density matrix.

Exercise: prove the proposition above.

Density matrices of the form $|\psi\rangle\langle\psi|$ are called *pure states*, the rest are *mixed*.

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Proposition

The set of density matrices is convex, i.e., given density matrices ρ_1, \dots, ρ_N and positive numbers w_1, \dots, w_N such that $\sum_i w_i = 1$, then $\sum_i w_i \rho_i$ is a density matrix.

Exercise: prove the proposition above.

Density matrices of the form $|\psi\rangle\langle\psi|$ are called *pure states*, the rest are *mixed*.

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Thus, a state ρ is mixed iff $\text{tr}(\rho^2) < 1$.

Density Matrices

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Let us prove it.

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Let us prove it.

ρ is positive hence self-adjoint, so by the spectral theorem we can write:

$$\rho = \sum_i \lambda_i P_i$$

where the λ_i 's are the eigenvalues and the P_i 's are the eigenspace projectors of ρ , respectively.

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Let us prove it.

ρ is positive hence self-adjoint, so by the spectral theorem we can write:

$$\rho = \sum_i \lambda_i P_i$$

where the λ_i 's are the eigenvalues and the P_i 's are the eigenspace projectors of ρ , respectively. The eigenvalues are all positive and satisfy $\sum_i \lambda_i = 1$ since $\text{tr}(\rho) = 1$.

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Let us prove it.

ρ is positive hence self-adjoint, so by the spectral theorem we can write:

$$\rho = \sum_i \lambda_i P_i$$

where the λ_i 's are the eigenvalues and the P_i 's are the eigenspace projectors of ρ , respectively. The eigenvalues are all positive and satisfy $\sum_i \lambda_i = 1$ since $\text{tr}(\rho) = 1$.

$$\text{tr}(\rho\rho) = \text{tr}\left(\rho \sum_i \lambda_i P_i\right) = \text{tr}\left(\sum_{i,j} \lambda_i \lambda_j P_i P_j\right) = \text{tr}\left(\sum_i \lambda_i^2 P_i\right) = \sum_i \lambda_i^2 \leq 1.$$

Density Matrices

Proposition

Let ρ be a density matrix. Then $\text{tr}(\rho^2) \leq 1$ and $\text{tr}(\rho^2) = 1$ iff ρ is a pure state.

Let us prove it.

ρ is positive hence self-adjoint, so by the spectral theorem we can write:

$$\rho = \sum_i \lambda_i P_i$$

where the λ_i 's are the eigenvalues and the P_i 's are the eigenspace projectors of ρ , respectively. The eigenvalues are all positive and satisfy $\sum_i \lambda_i = 1$ since $\text{tr}(\rho) = 1$.

$$\text{tr}(\rho\rho) = \text{tr}\left(\rho \sum_i \lambda_i P_i\right) = \text{tr}\left(\sum_{i,j} \lambda_i \lambda_j P_i P_j\right) = \text{tr}\left(\sum_i \lambda_i^2 P_i\right) = \sum_i \lambda_i^2 \leq 1.$$

Now, $\sum_i \lambda_i^2 = 1$ iff there is only one term in the sum, *i.e.*, ρ is a pure state.

The Rules of Quantum Mechanics with Density Matrices

Rule 1: The state space of Quantum Mechanics is a Hilbert space \mathcal{H} (the “physical” state of a system is a **density matrix**).

The Rules of Quantum Mechanics with Density Matrices

Rule 1: The state space of Quantum Mechanics is a Hilbert space \mathcal{H} (the “physical” state of a system is a **density matrix**).

Rule 2: A closed system evolves over time under **unitary transformations**. Applying unitary operator U to state ρ has the effect

$$\rho \rightarrow U\rho U^\dagger.$$

The Rules of Quantum Mechanics with Density Matrices

Rule 1: The state space of Quantum Mechanics is a Hilbert space \mathcal{H} (the “physical” state of a system is a **density matrix**).

Rule 2: A closed system evolves over time under **unitary transformations**. Applying unitary operator U to state ρ has the effect

$$\rho \rightarrow U\rho U^\dagger.$$

Here is “why”:

$$\{|\psi_i\rangle, w_i\} \xrightarrow{U} \{U|\psi_i\rangle, w_i\}$$

The Rules of Quantum Mechanics with Density Matrices

Rule 1: The state space of Quantum Mechanics is a Hilbert space \mathcal{H} (the “physical” state of a system is a **density matrix**).

Rule 2: A closed system evolves over time under **unitary transformations**. Applying unitary operator U to state ρ has the effect

$$\rho \rightarrow U\rho U^\dagger.$$

Here is “why”:

$$\begin{aligned} \{|\psi_i\rangle, w_i\} &\xrightarrow{U} \{U|\psi_i\rangle, w_i\} \\ \sum_i w_i |\psi_i\rangle\langle\psi_i| &\xrightarrow{U} \sum_i w_i U|\psi_i\rangle (U|\psi_i\rangle)^\dagger = \sum_i w_i U|\psi_i\rangle\langle\psi_i| U^\dagger = U\rho U^\dagger \end{aligned}$$

The Rules of Quantum Mechanics with Density Matrices

Rule 1: The state space of Quantum Mechanics is a Hilbert space \mathcal{H} (the “physical” state of a system is a **density matrix**).

Rule 2: A closed system evolves over time under **unitary transformations**. Applying unitary operator U to state ρ has the effect

$$\rho \rightarrow U\rho U^\dagger.$$

Here is “why”:

$$\begin{aligned} \{|\psi_i\rangle, w_i\} &\xrightarrow{U} \{U|\psi_i\rangle, w_i\} \\ \sum_i w_i |\psi_i\rangle\langle\psi_i| &\xrightarrow{U} \sum_i w_i U|\psi_i\rangle (U|\psi_i\rangle)^\dagger = \sum_i w_i U|\psi_i\rangle\langle\psi_i| U^\dagger = U\rho U^\dagger \end{aligned}$$

The (linear) function that maps ρ to $U\rho U^\dagger$ is called a *superoperator*.

The Rules of Quantum Mechanics with Density Matrices

Rule 3: A measurement is described by a collection $M = \{M_m\}$ of **measurement operators** that satisfy $\sum_m M_m^\dagger M_m = I$ (where m indexes the measurement outcomes).

The Rules of Quantum Mechanics with Density Matrices

Rule 3: A measurement is described by a collection $M = \{M_m\}$ of **measurement operators** that satisfy $\sum_m M_m^\dagger M_m = I$ (where m indexes the measurement outcomes).

The probability of measuring outcome m in state ρ is:

$$\text{Prob}(M = m; \rho) = \text{tr}(M_m^\dagger M_m \rho)$$

The Rules of Quantum Mechanics with Density Matrices

Rule 3: A measurement is described by a collection $M = \{M_m\}$ of **measurement operators** that satisfy $\sum_m M_m^\dagger M_m = I$ (where m indexes the measurement outcomes).

The probability of measuring outcome m in state ρ is:

$$\text{Prob}(M = m; \rho) = \text{tr}(M_m^\dagger M_m \rho)$$

The state ρ_m resulting after obtaining measurement outcome m is:

$$\rho_m = \frac{M_m \rho M_m^\dagger}{\text{tr}(M_m^\dagger M_m \rho)}$$

The Rules of Quantum Mechanics with Density Matrices

Rule 3: A measurement is described by a collection $M = \{M_m\}$ of **measurement operators** that satisfy $\sum_m M_m^\dagger M_m = I$ (where m indexes the measurement outcomes).

The probability of measuring outcome m in state ρ is:

$$\text{Prob}(M = m; \rho) = \text{tr}(M_m^\dagger M_m \rho)$$

The state ρ_m resulting after obtaining measurement outcome m is:

$$\rho_m = \frac{M_m \rho M_m^\dagger}{\text{tr}(M_m^\dagger M_m \rho)}$$

Exercise: verify that the probability Rule 3 above reduces to the usual rule for a self-adjoint observable (the measurement operators are simply the projectors on the eigenspaces).

Shannon Entropy

After Claude Shannon (electrical engineer; 1916-2001).

We have a source that produces symbols (from a finite alphabet) via a random process:

$$\textit{symb}_1 @ p_1, \textit{symb}_2 @ p_2, \dots, \textit{symb}_n @ p_n$$

The source can be thus thought of as a random variable.

How can we quantify the resources needed to store the symbols generated?

Shannon Entropy

After Claude Shannon (electrical engineer; 1916-2001).

We have a source that produces symbols (from a finite alphabet) via a random process:

$$\text{symp}_1 @ p_1, \text{symp}_2 @ p_2, \dots, \text{symp}_n @ p_n$$

The source can be thus thought of as a random variable.

How can we quantify the resources needed to store the symbols generated?

Definition

Given a random variable $X = \{s_i @ p_i\}$, the Shannon entropy is

$$H(X) = - \sum_i p_i \log p_i$$

Note: if $p_i = 0$ then we assume $p_i \log p_i = 0$.

Shannon Entropy

Example:

Consider a binary source, *i.e.*, it emits only two symbols:

$$\{\text{Pistacchio, @ } p; \text{ Stracciatella @ } (1 - p)\}$$

Shannon Entropy

Example:

Consider a binary source, *i.e.*, it emits only two symbols:

$$\{\text{Pistacchio, @ } p; \text{ Stracciatella @ } (1 - p)\}$$

$$H(p) = -p \log p - (1 - p) \log(1 - p)$$

$H(p)$ is largest for $p = \frac{1}{2}$ and smallest for $p = 0$ (or $p = 1$).

Shannon Entropy

Definition

The joint entropy of random variables X, Y is:

$$H(X, Y) = - \sum_{x,y} p(x, y) \log p(x, y)$$

Definition

The mutual information of random variables X, Y is:

$$H(X : Y) = H(X) + H(Y) - H(X, Y)$$

$H(X : Y)$ is informally the amount of information that X and Y have in common.

The von Neumann Entropy

After John (János) von Neumann (mathematician; 1903-1957).

Definition

The entropy of a density matrix ρ is:

$$S(\rho) = -\text{tr}(\rho \log \rho)$$

or equivalently:

$$S(\rho) = -\sum_i \lambda_i \log \lambda_i$$

where the λ_i 's are the eigenvalues of ρ .

The von Neumann Entropy

After John (János) von Neumann (mathematician; 1903-1957).

Definition

The entropy of a density matrix ρ is:

$$S(\rho) = -\text{tr}(\rho \log \rho)$$

or equivalently:

$$S(\rho) = -\sum_i \lambda_i \log \lambda_i$$

where the λ_i 's are the eigenvalues of ρ .

Example: consider the density matrix I/n in an n -dimensional Hilbert space.

$$S(I/n) = -\sum_{i=1}^n \frac{1}{n} \log \frac{1}{n} = -\frac{1}{n} \log \frac{1}{n} \sum_{i=1}^n = -\log \frac{1}{n} = \log n$$

In fact, I/n is the state with highest entropy.

Holevo Bound

Context: a game between two parties, Alice and Bob.

Alice has a random source of symbols $X = 1, \dots, n$ with probability p_1, \dots, p_n . Bob aims at determining the value of X .

Holevo Bound

Context: a game between two parties, Alice and Bob.

Alice has a random source of symbols $X = 1, \dots, n$ with probability p_1, \dots, p_n . Bob aims at determining the value of X .

Alice prepares a state ρ_X chosen from ρ_1, \dots, ρ_n and gives it to Bob, who performs a quantum measurement, denoted Y , to guess X as best as he can.

Holevo Bound

Context: a game between two parties, Alice and Bob.

Alice has a random source of symbols $X = 1, \dots, n$ with probability p_1, \dots, p_n . Bob aims at determining the value of X .

Alice prepares a state ρ_X chosen from ρ_1, \dots, ρ_n and gives it to Bob, who performs a quantum measurement, denoted Y , to guess X as best as he can.

Classically: $H(X : Y) \leq H(X)$, so Bob “wins” if $H(X : Y) = H(X)$. (And classical states can be easily distinguished.)

Holevo Bound

Context: a game between two parties, Alice and Bob.

Alice has a random source of symbols $X = 1, \dots, n$ with probability p_1, \dots, p_n . Bob aims at determining the value of X .

Alice prepares a state ρ_X chosen from ρ_1, \dots, ρ_n and gives it to Bob, who performs a quantum measurement, denoted Y , to guess X as best as he can.

Classically: $H(X : Y) \leq H(X)$, so Bob “wins” if $H(X : Y) = H(X)$. (And classical states can be easily distinguished.)

Quantumly: non-orthogonal quantum states cannot be distinguished with certainty! (Remember the BB84 quantum key-distribution protocol.)

Holevo Bound

After Alexander Holevo (physicist; 1943 -).

Theorem

Suppose Alice prepares ρ_X for $X = 1, \dots, n$ with probability p_1, \dots, p_n . Suppose Bob performs a measurement on ρ_X with outcome Y . Then:

$$H(X : Y) \leq S(\rho) - \sum_i p_i \log S(\rho_i)$$

where $\rho = \sum_i p_i \rho_i$.

Holevo Bound

After Alexander Holevo (physicist; 1943 -).

Theorem

Suppose Alice prepares ρ_X for $X = 1, \dots, n$ with probability p_1, \dots, p_n . Suppose Bob performs a measurement on ρ_X with outcome Y . Then:

$$H(X : Y) \leq S(\rho) - \sum_i p_i \log S(\rho_i)$$

where $\rho = \sum_i p_i \rho_i$.

One can also show that $S(\rho) - \sum_i p_i \log S(\rho_i) \leq H(X)$ (with equality iff the states are orthogonal). Thus, in general Bob cannot learn X with certainty.