Prediction-error distributions in every test fold

a DYZ3

b Yq12

Repeat 1
Repeat 2
Repeat 3
Repeat 4
Repeat 5
Repeat 6
Repeat 7
Repeat 8
Repeat 9
Repeat 10

0 2 5 10 20 30

Individual absolute error (%)

Within each repeat, folds 1–5 run top to bottom. Boxes: 25th–75th percentiles; median; whiskers: 10th–90th.

Points retain errors beyond the whiskers. Axes are logarithmic above 5%. Repeated individuals are not independent.

Prediction-error distributions in every test fold

c DYZ1

d DYZ2

Repeat 1
Repeat 2
Repeat 3
Repeat 4
Repeat 5
Repeat 6
Repeat 7
Repeat 8
Repeat 9
Repeat 10

0 2 5 10 20 30

Individual absolute error (%)


Within each repeat, folds 1–5 run top to bottom. Boxes: 25th–75th percentiles; median; whiskers: 10th–90th.

Points retain errors beyond the whiskers. Axes are logarithmic above 5%. Repeated individuals are not independent.

Prediction-error distributions in every test fold

e DYZ18S

f YqHet2.7



Within each repeat, folds 1–5 run top to bottom. Boxes: 25th–75th percentiles; median; whiskers: 10th–90th.

Points retain errors beyond the whiskers. Axes are logarithmic above 5%. Repeated individuals are not independent.

Prediction-error distributions in every test fold

g DYZ18sub

h DYZ19


Repeat 1
Repeat 2
Repeat 3
Repeat 4
Repeat 5
Repeat 6
Repeat 7
Repeat 8
Repeat 9
Repeat 10

0 2 5 10 20 30

Individual absolute error (%)

Within each repeat, folds 1–5 run top to bottom. Boxes: 25th–75th percentiles; median; whiskers: 10th–90th.

Points retain errors beyond the whiskers. Axes are logarithmic above 5%. Repeated individuals are not independent.



Within each repeat, folds 1–5 run top to bottom. Boxes: 25th–75th percentiles; median; whiskers: 10th–90th.

Points retain errors beyond the whiskers. Axes are logarithmic above 5%. Repeated individuals are not independent.