
Formatting instructions for NIPS 2018

Anonymous Author(s)

Affiliation

Address

email

Abstract

1 The abstract paragraph should be indented $\frac{1}{2}$ inch (3 picas) on both the left- and
2 right-hand margins. Use 10 point type, with a vertical spacing (leading) of 11 points.
3 The word **Abstract** must be centered, bold, and in point size 12. Two line spaces
4 precede the abstract. The abstract must be limited to one paragraph.

5 1 Introduction

6 2 Preliminaries

7 We define a Markov decision process (MDP) as a tuple $\mathcal{M} = \langle \mathcal{S}, \mathcal{A}, \mathcal{P}, \mathcal{R}, p_0, \gamma \rangle$, where \mathcal{S} is
8 the state-space, \mathcal{A} is a finite set of actions, $\mathcal{P}(\cdot|s, a)$ is the distribution of the next state s' given
9 that action a is taken in state s , $\mathcal{R} : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R}$ is the reward function, p_0 is the initial-state
10 distribution, and $\gamma \in [0, 1)$ is the discount factor. We assume the reward function to be uniformly
11 bounded by a constant $R_{max} > 0$. A deterministic policy $\pi : \mathcal{S} \rightarrow \mathcal{A}$ is a mapping from states
12 to actions. At the beginning of each episode of interaction, the initial state s_0 is drawn from p_0 .
13 Then, the agent takes the action $a_0 = \pi(s_0)$, receives a reward $\mathcal{R}(s_0, a_0)$, transitions to the next
14 state $s_1 \sim \mathcal{P}(\cdot|s_0, a_0)$, and the process is repeated. The goal is to find the policy maximizing the
15 long-term return over a possibly infinite horizon: $\max_{\pi} J(\pi) \triangleq \mathbb{E}[\sum_{t=0}^{\infty} \gamma^t r_t | \mathcal{M}, \pi]$. To this end,
16 we define the optimal value function $Q^*(s, a)$ as the expected return obtained by taking action a
17 in state s and following an optimal policy thereafter. Then, an optimal policy π^* is a policy that
18 is greedy with respect to the optimal value function, i.e., $\pi^*(s) = \operatorname{argmax}_a Q^*(s, a)$ for all states
19 s . It can be shown (e.g., [1]) that Q^* is the unique fixed-point of the optimal Bellman operator T
20 defined by $TQ(s, a) = \mathcal{R}(s, a) + \gamma \mathbb{E}_{\mathcal{P}}[\max_{a'} Q(s', a')]$ for any value function Q . From now on, we
21 adopt the term Q -function to denote any plausible value function, i.e., any function $Q : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R}$
22 uniformly bounded by $\frac{R_{max}}{1-\gamma}$.

23 When learning the optimal value function, a quantity of interest is how close a given Q -function
24 is to the fixed-point of the Bellman operator. This is given by its Bellman residual, defined by
25 $B(Q) \triangleq TQ - Q$. Notice that Q is optimal if, and only if, $B(Q)(s, a) = 0$ for all s, a . Furthermore,
26 if we assume the existence of a distribution ν over $\mathcal{S} \times \mathcal{A}$, the squared Bellman error of Q is
27 defined as the expected squared Bellman residual of Q under ν , $\|B(Q)\|_{\nu}^2 = \mathbb{E}_{\mu}[B^2(Q)]$. Although
28 minimizing the empirical Bellman error is an appealing objective, it is well-known that an unbiased
29 estimator requires two independent samples of the next state s' of each s, a (e.g., [2]). In practice,
30 the empirical Bellman error is typically replaced by the TD error, which approximates the former
31 using a single transition sample. Given a dataset of N samples, the TD error is computed as
32 $\|B(Q)\|_D^2 = \frac{1}{N} \sum_{i=1}^N (r_i + \gamma \max_{a'} Q(s'_i, a') - Q(s_i, a_i))^2$.

cite Maillard

3 Variational Transfer Learning

In this section, we describe our variational approach to transfer in RL. In Section 3.1, we start by introducing our algorithm from a high-level perspective, in such a way that it can be used for any choice of prior and posterior distributions. Then, in Sections 3.2 and 3.3, we propose practical implementations based on Gaussian prior/posterior and mixture of Gaussian prior/posterior, respectively.

3.1 Algorithm

We begin with a simple consideration: the distribution \mathcal{D} over tasks clearly induces a distribution over optimal Q -functions. Since, for any MDP, learning its optimal Q -function is sufficient for solving the problem, one can safely replace the distribution over tasks with the distribution over their optimal value functions. Furthermore, assume we know such distribution and we are given a new task τ to solve. Our goal is to design an algorithm that efficiently explores τ so as to quickly adapt the prior distribution in a Bayesian fashion to put all probability mass over the optimal Q -function of τ .

We consider a parametric family of Q -functions, $\mathcal{Q} = \{Q_{\mathbf{w}} : \mathcal{S} \times \mathcal{A} \rightarrow \mathbb{R} \mid \mathbf{w} \in \mathbb{R}^K\}$. For simplicity, we assume each function in \mathcal{Q} to be uniformly bounded by $\frac{R_{max}}{1-\gamma}$. Then, we can reduce our prior distribution over Q -functions to a prior distribution over weights $p(\mathbf{w})$. Assume that we are given a dataset $D = \{(s_i, a_i, s'_i, r_i) \mid i = 1, 2, \dots, N\}$ of samples from some task τ that we want to solve. Then, the posterior distribution over weights given such dataset can be computed by applying Bayes theorem as $p(\mathbf{w}|D) \propto p(D|\mathbf{w})p(\mathbf{w})$. Unfortunately, this cannot be directly used in practice since we do not have a model of the likelihood $p(D|\mathbf{w})$. In such case, it is very common to make strong assumptions on the MDPs or the Q -functions so as to get tractable posteriors []. However, in our transfer settings all distributions involved depend on the family of tasks under consideration, and making such assumptions is likely to limit the applicability of the approach. Thus, we take a different approach to derive a more general and meaningful solution. Recall that our final goal is move all probability mass over the weights minimizing some empirical loss measure, which in our case is the TD error $\|B(\mathbf{w})\|_D^2$. Then, given a prior $p(\mathbf{w})$, we know from PAC-Bayesian theory that the optimal Gibbs posterior takes the form []:

Cite somebody

Cite Catoni 2007

$$q(\mathbf{w}) = \frac{e^{-\Lambda \|B(\mathbf{w})\|_D^2} p(\mathbf{w})}{\int e^{-\Lambda \|B(\mathbf{w}')\|_D^2} p(d\mathbf{w}')} \quad (1)$$

for some parameter $\Lambda > 0$. Since Λ is typically chosen to increase with the number of samples N , in the remaining we set it to $\lambda^{-1}N$, for some constant $\lambda > 0$. Notice that, whenever the term $e^{-\Lambda \|B(\mathbf{w})\|_D^2}$ can be interpreted as the actual likelihood of D , q becomes a classic Bayesian posterior. Although we now have an appealing distribution, the integral at the denominator of (1) is intractable to compute even for simple Q -function models. Thus, we propose a variational approximation q_{ξ} by considering a simpler family of distributions parameterized by $\xi \in \Xi$. Then, our problem reduces to finding the variational parameters ξ such that q_{ξ} minimizes the Kullback-Leibler (KL) divergence w.r.t. the Gibbs posterior q . From the theory of variational inference (e.g., []), this can be shown to be equivalent to minimizing the well-known (negative) *evidence lower bound* (ELBO):

cite

$$\min_{\xi \in \Xi} \mathcal{L}(\xi) = \mathbb{E}_{\mathbf{w} \sim q_{\xi}} [\|B(\mathbf{w})\|_D^2] - \frac{\lambda}{N} KL(q_{\xi}(\mathbf{w}) \parallel p(\mathbf{w})) \quad (2)$$

Intuitively, the approximate posterior trades-off between placing probability mass over those weights \mathbf{w} that have low TD error (first term), and staying close to the prior distribution (second term). Assuming that we are able to compute the gradients of (2) w.r.t. the variational parameters ξ , our objective can be easily optimized with any stochastic optimization algorithm.

We now highlight our general transfer procedure in Alg. 1, while deferring a description of practical implementations with specific choices for the distributions involved to the next two sections. Given a set of weights \mathcal{W}_s from the source tasks, we start by estimating the prior distribution (line 1) and we initialize the variational parameters by minimizing the KL divergence w.r.t. such distribution² (line

¹In practice, this is easily achieved by truncation.

²If the prior and approximate posterior were in the same family of distributions we could simply set ξ to the prior parameters, however this does not always hold in practice.

Algorithm 1 Variational Transfer

Require: Target task τ , source Q -function weights \mathcal{W}_s , batch sizes M_D and $M_{\mathcal{W}}$, prior weight λ

```
1: Estimate prior  $p(\mathbf{w})$  from  $\mathcal{W}_s$ 
2: Initialize variational parameters:  $\xi \leftarrow \operatorname{argmin}_{\xi} KL(q_{\xi} || p)$ 
3: Initialize replay buffer:  $D = \emptyset$ 
4: repeat
5:   Sample initial state:  $s_0 \sim p_0^{(\tau)}$ 
6:   while  $s_h$  is not terminal do
7:     Sample weights:  $\mathbf{w} \sim q_{\xi}(\mathbf{w})$ 
8:     Take action  $a_h = \operatorname{argmax}_a Q_{\mathbf{w}}(s_h, a)$ 
9:     Observe transition  $s_{h+1} \sim \mathcal{P}^{(\tau)}(\cdot | s_h, a_h)$ 
10:    Collect reward  $r_h = \mathcal{R}^{(\tau)}(s_h, a_h)$ 
11:    Add sample to the replay buffer:  $D \leftarrow D \cup \langle s_h, a_h, r_h, s_{h+1} \rangle$ 
12:    Sample batch  $D' = \langle s_i, a_i, r_i, s'_i \rangle_{i=1}^{M_D}$  from  $D$  and  $\mathcal{W} = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{M_{\mathcal{W}}}\}$  from  $q_{\xi}$ 
13:    Approximate objective:  $\mathcal{L}(\xi) = \frac{1}{M_{\mathcal{W}}} \sum_{\mathbf{w} \in \mathcal{W}} \|B(\mathbf{w})\|_{D'}^2 - \frac{\lambda}{N} KL(q_{\xi} || p)$ 
14:    Compute the gradient  $\nabla_{\xi} \mathcal{L}(\xi)$ 
15:    Update  $\xi$  in the direction of  $\nabla_{\xi} \mathcal{L}(\xi)$  using any stochastic optimizer (e.g., ADAM)
16:  end while
17: until forever
```

2). Then, at each time step of interaction, we re-sample the weights from the current approximate posterior and act greedily w.r.t. the corresponding Q -function (lines 7,8). This resembles the well-known Thompson sampling approach adopted in multi-armed bandits []and allows our algorithm to efficiently explore the target task. In some sense, at each time we guess what is the task we are trying to solve based on our current belief and we act as if such guess were actually true. After collecting and storing the new experience (lines 9-11), we draw a batch of samples from the replay buffer and a batch of weights from the posterior (line 12). We use these to approximate the negative ELBO, compute its gradient, and finally update the variational parameters (lines 13-15).

Cite

The main advantage of our approach is that it exploits knowledge from the source tasks to perform an efficient adaptive exploration. Intuitively, during the first steps of interaction, our algorithm has no idea about what is the current task. However, it can rely on the learned prior to take early informed decisions. As the learning process goes on, it will quickly figure out which task is being solved, thus moving all probability mass over the weights minimizing the TD error. From that point, sampling from the posterior is approximately equivalent to deterministically taking such weights, and no more exploration will be performed. Finally, notice the generality of the proposed approach: as far as the objective \mathcal{L} is differentiable in the variational parameters ξ , and its gradients can be efficiently computed, any function approximator for the Q -functions and any family for the prior and posterior distributions can be adopted. For the latter, we describe two practical choices in the next two sections.

3.2 Gaussian Variational Transfer

We now restrict ourselves to a specific choice of the prior and posterior families that makes our algorithm very efficient and easy to implement. We assume that optimal Q -functions according to our task distribution (or better, their weights) follow a multivariate Gaussian law. That is, we model the prior as $p(\mathbf{w}) = \mathcal{N}(\boldsymbol{\mu}_p, \boldsymbol{\Sigma}_p)$ and we learn its parameters from the set of source weights using, e.g., maximum likelihood estimation (with small regularization to make sure the covariance is positive definite). Then, our variational family is the set of all well-defined Gaussian distributions, i.e., the variational parameters are $\Xi = \{(\boldsymbol{\mu}, \boldsymbol{\Sigma}) \mid \boldsymbol{\mu} \in \mathbb{R}^K, \boldsymbol{\Sigma} \in \mathbb{R}^{K \times K}, \boldsymbol{\Sigma} \succ 0\}$. To prevent the covariance from going not positive definite, we consider its Cholesky decomposition $\boldsymbol{\Sigma} = \mathbf{L}\mathbf{L}^T$ and learn the lower-triangular Cholesky factor \mathbf{L} instead. Under Gaussian distributions, all quantity of interest for using Alg. 1 can be computed very easily. The KL divergence between the prior and approximate posterior can be computed in closed-form as:

$$KL(q_{\xi}(\mathbf{w}) || p(\mathbf{w})) = \frac{1}{2} \left(\log \frac{|\boldsymbol{\Sigma}_p|}{|\boldsymbol{\Sigma}|} + \operatorname{Tr}(\boldsymbol{\Sigma}_p^{-1} \boldsymbol{\Sigma}) + (\boldsymbol{\mu} - \boldsymbol{\mu}_p)^T \boldsymbol{\Sigma}_p^{-1} (\boldsymbol{\mu} - \boldsymbol{\mu}_p) - K \right) \quad (3)$$

for $\xi = (\mu, L)$ and $\Sigma = LL^T$. Its gradients with respect to the variational parameters are []:

$$\nabla_{\mu} KL(q_{\xi}(\mathbf{w}) \parallel p(\mathbf{w})) = \Sigma_p^{-1}(\mu - \mu_p) \quad (4)$$

Cite matrix
cookbook

$$\nabla_L KL(q_{\xi}(\mathbf{w}) \parallel p(\mathbf{w})) = \Sigma_p^{-1}L - (L^{-1})^T \quad (5)$$

Finally, the gradients w.r.t. the expected likelihood term of the variational objective (2) can be computed using the reparameterization trick (e.g., []):

$$\nabla_{\mu} \mathbb{E}_{\mathbf{w} \sim \mathcal{N}(\mu, LL^T)} [\|B(\mathbf{w})\|_D^2] = \mathbb{E}_{\mathbf{v} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [\nabla_{\mathbf{w}} \|B(\mathbf{w})\|_D^2] \text{ for } \mathbf{w} = L\mathbf{v} + \mu \quad (6)$$

Cite deep-
mind and
another

$$\nabla_L \mathbb{E}_{\mathbf{w} \sim \mathcal{N}(\mu, LL^T)} [\|B(\mathbf{w})\|_D^2] = \mathbb{E}_{\mathbf{v} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [\nabla_{\mathbf{w}} \|B(\mathbf{w})\|_D^2 \cdot \mathbf{v}^T] \text{ for } \mathbf{w} = L\mathbf{v} + \mu \quad (7)$$

3.3 Mixture of Gaussian Variational Transfer

Although the Gaussian assumption of the previous section is very appealing as it allows for a simple and efficient way of computing the variational objective and its gradients, we believe that such assumption almost never holds in practice. In fact, even for families of tasks in which the reward and transition models follow a Gaussian law, the Q -values might be far from it. Depending on the family of tasks under consideration and, since we are learning a distribution over weights, on the chosen function approximator, the prior might have arbitrarily complex shapes. When the information loss due to the Gaussian approximation becomes too severe, the algorithm is likely to fail at transferring knowledge, thus reducing to almost random exploration. We now propose a variant to successfully solve this problem, while keeping the algorithm simple and efficient enough to be applied in practice. In order to capture arbitrarily complex distributions, we use a kernel estimator [] for learning our prior. Assume we are given a set \mathcal{W}_s of weights from the source tasks. Then, our estimated prior places a single isotropic Gaussian over each weight: $p(\mathbf{w}) = \frac{1}{|\mathcal{W}_s|} \sum_{\mathbf{w}_s \in \mathcal{W}_s} \mathcal{N}(\mathbf{w} | \mathbf{w}_s, \sigma_p^2 \mathbf{I})^3$. This takes the form of a mixture of Gaussians with equally weighted components. Consistently with the prior, we model our approximate posterior as a mixture of Gaussians. However, we allow a different number of components (typically much less than the prior's) and we adopt full covariances instead of only diagonals, so that our posterior has the potential to match complex distributions with less components. Using C components, our posterior is $q_{\xi}(\mathbf{w}) = \frac{1}{C} \sum_{i=1}^C \mathcal{N}(\mathbf{w} | \mu_i, \Sigma_i)$, with variational parameters $\xi = (\mu_1, \dots, \mu_C, \Sigma_1, \dots, \Sigma_C)$. Once again, we learn Cholesky factors instead of full covariances.

Although this new model has the potential to capture much more complex distributions, it poses a major complication: the KL divergence between two mixture of Gaussians is well-known to have no closed-form equation. To solve this issue, we can rely on an upper bound to such quantity, so that negative ELBO we are optimizing still represents an upper bound on the KL between the approximate and true posterior. However, this turns out to be non-trivial as well. In fact, it is very easy to bound the KL between two mixtures with the KLs between each couple of components. However, the loss of information is such that minimizing the upper bound via gradient methods converges to a local optimum in which all components tend to go to the same point, thus almost reducing to the single Gaussian case. To solve this issue, we adopt the variational upper bound proposed in [], which we found to be able to preserve the needed information. We report it here for the sake of completeness. See the original paper for the proof.

Cite UB

Theorem 1. Let $p = \sum_i c_i^{(p)} f_i^{(p)}$ and $q = \sum_j c_j^{(q)} f_j^{(q)}$ be two mixture of Gaussian distributions, where $f_i^{(p)} = \mathcal{N}(\mu_i^{(p)}, \Sigma_i^{(p)})$ denotes the i -th component of p , $c_i^{(p)}$ denotes its weight, and similarly for q . Introduce two vectors $\chi^{(1)}$ and $\chi^{(2)}$ such that $c_i^{(p)} = \sum_j \chi_{j,i}^{(2)}$ and $c_j^{(q)} = \sum_i \chi_{i,j}^{(1)}$. Then:

$$KL(p \parallel q) \leq KL(\chi^{(2)} \parallel \chi^{(1)}) + \sum_{i,j} \chi_{j,i}^{(2)} KL(f_i^{(p)} \parallel f_j^{(q)}) \quad (8)$$

Our new algorithm replaces the KL with the above-mentioned upper bound. Each time we require its value, we have to recompute the parameters $\chi^{(1)}$ and $\chi^{(2)}$ that tighten the bound. As shown in [], this can be achieved by a simple fixed-point procedure. Furthermore, both terms in the approximate negative ELBO are now linear combinations of functions of the variational parameters for different components, thus their gradients can be straightforwardly derived from the ones of the Gaussian case.

cite them
again

³Notice that this is slightly different than the typical kernel estimator (e.g., [])

150 3.4 Optimizing the TD error

151 From Sections 3.2 and 3.3, we know that differentiating the negative ELBO \mathcal{L} w.r.t. ξ requires
 152 differentiating $\|B(w)\|_D^2$ w.r.t. w . Unfortunately, the TD error is well-known to be non-differentiable
 153 due to the presence of the max operator. This rarely represents a problem since typical value-based
 154 algorithm are actually semi-gradient methods, i.e., they do not differentiate the targets (see, e.g.,
 155 Chapter 11 of []). However, in our case ...

Cite Sutton

156 What is a good motivation for the fact that we need a residual algorithm?

157 To solve this issue, we replace the optimal Bellman operator with the mellow Bellman operator
 158 introduced in [], which adopts a softened version of max called *mellowmax*:

Cite MM

$$\text{mm}_a Q_w(s, a) = \frac{1}{\kappa} \log \frac{1}{|\mathcal{A}|} \sum_a e^{\kappa Q_w(s, a)} \quad (9)$$

159 where κ is a hyperparameter and $|\mathcal{A}|$ is the number of actions. We defer a theoretical analysis of the
 160 consequences of this approximation to Section 4. The mellow Bellman operator, which we denote
 161 as \tilde{T} , has several appealing properties that make it suitable for our settings: (i) it converges to the
 162 maximum as $\kappa \rightarrow \infty$, (ii) it has a unique fixed point, and (iii) it is *differentiable*. Denoting by
 163 $\tilde{B}(w) = \tilde{T}Q_w - Q_w$ the Bellman residual w.r.t. the mellow Bellman operator \tilde{T} , we have that the
 164 corresponding TD error, $\|\tilde{B}(w)\|_D^2$, is now differentiable with respect to w .

165 Here we need to talk about residual algorithms and how to improve their gradient

166 4 Theoretical Analysis

167 In this section, we theoretically analyze our variational transfer algorithm...

168 A first important question that we need to answer is whether replacing max with mellow-max in
 169 the Bellman operator constitutes a strong approximation or not. It has been proved [] that the
 170 mellow Bellman operator is a contraction under the L_∞ -norm and, thus, has a unique fixed-point.
 171 However, how such fixed-point differs from the one of the optimal Bellman operator remains an open
 172 question. Since mellow-max monotonically converges to max as $\kappa \rightarrow \infty$, it would be desirable if
 173 the corresponding operator also monotonically converged to the optimal one. We confirm that this
 174 property actually holds in the following theorem.

Cite MM

175 **Theorem 2.** *Let V be the fixed-point of the optimal Bellman operator T , and Q the corresponding*
 176 *action-value function. Define the action-gap function $g(s)$ as the difference between the value of*
 177 *the best action and the second best action at each state s . Let \tilde{V} be the fixed-point of the mellow*
 178 *Bellman operator \tilde{T} with parameter $\kappa > 0$ and denote by $\beta > 0$ the inverse temperature of the*
 179 *induced Boltzmann distribution (as in []). Let ν be a probability measure over the state-space. Then,*
 180 *for any $p \geq 1$:*

Cite MM

$$\|V - \tilde{V}\|_{\nu, p}^p \leq \frac{2R_{max}}{(1 - \gamma)^2} \left\| 1 - \frac{1}{1 + |\mathcal{A}| e^{-\beta g}} \right\|_{\nu, p}^p \quad (10)$$

181 **5 Related Works**

182 **6 Experiments**

183 **6.1 Gridworld**

184 **6.2 Classic Control**

185 **6.3 Maze Navigation**

186 **7 Conclusion**

187 **References**

- 188 [1] Martin L. Puterman. *Markov Decision Processes: Discrete Stochastic Dynamic Programming*. John Wiley
189 & Sons, Inc., New York, NY, USA, 1994.

190 A Proofs

191 **Theorem 2.** *Let V be the fixed-point of the optimal Bellman operator T , and Q the corresponding*
 192 *action-value function. Define the action-gap function $g(s)$ as the difference between the value of*
 193 *the best action and the second best action at each state s . Let \tilde{V} be the fixed-point of the mellow*
 194 *Bellman operator \tilde{T} with parameter $\kappa > 0$ and denote by $\beta > 0$ the inverse temperature of the*
 195 *induced Boltzmann distribution (as in []). Let ν be a probability measure over the state-space. Then,*
 196 *for any $p \geq 1$:*

Cite MM

$$\|V - \tilde{V}\|_{\nu,p}^p \leq \frac{2R_{max}}{(1-\gamma)^2} \left\| 1 - \frac{1}{1 + |\mathcal{A}| e^{-\beta g}} \right\|_{\nu,p}^p \quad (10)$$

197 *Proof.* We begin by noticing that:

$$\begin{aligned} \|V - \tilde{V}\|_{\nu,p}^p &= \|TV - \tilde{T}\tilde{V}\|_{\nu,p}^p \\ &= \|TV - \tilde{T}V + \tilde{T}V - \tilde{T}\tilde{V}\|_{\nu,p}^p \\ &\leq \|TV - \tilde{T}V\|_{\nu,p}^p + \|\tilde{T}V - \tilde{T}\tilde{V}\|_{\nu,p}^p \\ &\leq \|TV - \tilde{T}V\|_{\nu,p}^p + \gamma \|V - \tilde{V}\|_{\nu,p}^p \end{aligned}$$

198 where the first inequality follows from Minkowsky's inequality and the second one from the contrac-
 199 tion property of the mellow Bellman operator. This implies that:

$$\|V - \tilde{V}\|_{\nu,p}^p \leq \frac{1}{1-\gamma} \|TV - \tilde{T}V\|_{\nu,p}^p \quad (11)$$

200 Let us bound the norm on the right-hand side separately. In order to do that, we will bound the
 201 function $|TV(s) - \tilde{T}V(s)|$ point-wisely for any state s . By applying the definition of the optimal
 202 and mellow Bellman operators, we obtain:

$$\begin{aligned} |TV(s) - \tilde{T}V(s)| &= \left| \max_a \{R(s, a) + \gamma \mathbb{E}[V(s')]\} - \min_a \{R(s, a) + \gamma \mathbb{E}[V(s')]\} \right| \\ &= \left| \max_a Q(s, a) - \min_a Q(s, a) \right| \end{aligned}$$

203 Recall that applying the mellow-max is equivalent to computing an expectation under a Boltzmann
 204 distribution with inverse temperature β induced by Q []. Thus, we can write:

Cite MM

$$\begin{aligned} \left| \max_a Q(s, a) - \min_a Q(s, a) \right| &= \left| \sum_a \pi^*(a|s) Q(s, a) - \sum_a \pi_\beta(a|s) Q(s, a) \right| \\ &= \left| \sum_a Q(s, a) (\pi^*(a|s) - \pi_\beta(a|s)) \right| \\ &\leq \sum_a |Q(s, a)| |\pi^*(a|s) - \pi_\beta(a|s)| \\ &\leq \frac{R_{max}}{1-\gamma} \sum_a |\pi^*(a|s) - \pi_\beta(a|s)| \end{aligned} \quad (12)$$

205 where π^* is the optimal (deterministic) policy w.r.t. Q and π_β is the Boltzmann distribution induced
 206 by Q with inverse temperature β :

$$\pi_\beta(a|s) = \frac{e^{\beta Q(s,a)}}{\sum_{a'} e^{\beta Q(s,a')}}$$

207 Denote by $a_1(s)$ the optimal action for state s under Q . We can then write:

$$\begin{aligned}
\sum_a |\pi^*(a|s) - \pi_\beta(a|s)| &= |\pi^*(a_1(s)|s) - \pi_\beta(a_1(s)|s)| + \sum_{a \neq a_1(s)} |\pi^*(a|s) - \pi_\beta(a|s)| \\
&= |1 - \pi_\beta(a_1(s)|s)| + \sum_{a \neq a_1(s)} |\pi_\beta(a|s)| \\
&= 2 |1 - \pi_\beta(a_1(s)|s)|
\end{aligned} \tag{13}$$

208 Finally, let us bound this last term:

$$\begin{aligned}
|1 - \pi_\beta(a_1(s)|s)| &= \left| 1 - \frac{e^{\beta Q(s, a_1(s))}}{\sum_{a'} e^{\beta Q(s, a')}} \right| \\
&= \left| 1 - \frac{e^{\beta(Q(s, a_1(s)) - Q(s, a_2(s)))}}{\sum_{a'} e^{\beta(Q(s, a') - Q(s, a_2(s)))}} \right| \\
&= \left| 1 - \frac{e^{\beta g(s)}}{\sum_{a'} e^{\beta(Q(s, a') - Q(s, a_2(s)))}} \right| \\
&= \left| 1 - \frac{e^{\beta g(s)}}{e^{\beta g(s)} + \sum_{a' \neq a_1(s)} e^{\beta(Q(s, a') - Q(s, a_2(s)))}} \right| \\
&\leq \left| 1 - \frac{e^{\beta g(s)}}{e^{\beta g(s)} + |\mathcal{A}|} \right| \\
&= \left| 1 - \frac{1}{1 + |\mathcal{A}| e^{-\beta g(s)}} \right|
\end{aligned} \tag{14}$$

209 Combining Eq. (12), (13), and (14), we obtain:

$$\left| \max_a Q(s, a) - \min_a Q(s, a) \right| \leq \frac{2R_{max}}{1 - \gamma} \left| 1 - \frac{1}{1 + |\mathcal{A}| e^{-\beta g(s)}} \right|$$

210 Taking the norm and plugging this into Eq. (11) concludes the proof. \square

211 **Lemma 1.** Let p and ν denote probability measures over Q -functions and state-action pairs, respectively. Assume Q^* is the unique fixed-point of the optimal Bellman operator T . Then, for any $\delta > 0$,
212 with probability at least $1 - \delta$ over the choice of a Q -function Q , the following holds:
213

$$\|Q - Q^*\|_\nu^2 \leq \frac{\mathbb{E}_p [\|B(Q)\|_\nu^2]}{(1 - \gamma)\delta} \tag{15}$$

214 *Proof.* First notice that:

$$\begin{aligned}
\|Q - Q^*\| &= \|Q + TQ - TQ - TQ^*\| \\
&\leq \|Q - TQ\| + \|TQ - TQ^*\| \\
&\leq \|Q - TQ\| + \gamma \|Q - Q^*\| \\
&= \|B(Q)\| + \gamma \|Q - Q^*\|
\end{aligned}$$

215 which implies that:

$$\|Q - Q^*\| \leq \frac{1}{1 - \gamma} \|B(Q)\|$$

216 Then we can write:

$$P(\|Q - Q^*\| > \epsilon) \leq P(\|B(Q)\| > \epsilon(1 - \gamma)) \leq \frac{\mathbb{E}_p [\|B(Q)\|_\nu^2]}{(1 - \gamma)\epsilon}$$

217 Settings the right-hand side equal to δ and solving for ϵ concludes the proof. \square

218 **Corollary 1.** Let p and ν denote probability measures over Q -functions and state-action pairs,
 219 respectively. Assume \tilde{Q} is the unique fixed-point of the mellow Bellman operator \tilde{T} . Then, for any
 220 $\delta > 0$, with probability at least $1 - \delta$ over the choice of a Q -function Q , the following holds:

$$\|Q - \tilde{Q}\|_\nu^2 \leq \frac{\mathbb{E}_p \left[\|\tilde{B}(Q)\|_\nu^2 \right]}{(1 - \gamma)\delta} \quad (16)$$

221 **Lemma 2.** Assume Q -functions belong to a parametric space of functions bounded by $\frac{R_{max}}{1-\gamma}$. Let p
 222 and q be arbitrary distributions over the parameter space \mathcal{W} , and ν be a probability measure over
 223 $\mathcal{S} \times \mathcal{A}$. Consider a dataset D of N samples and define $v(\mathbf{w}) \triangleq \mathbb{E}_\nu [\text{Var}_{\mathcal{P}} [b(\mathbf{w})]]$. Then, for any
 224 $\delta > 0$, with probability at least $1 - \delta$, the following two inequalities hold simultaneously:

$$\mathbb{E}_q \left[\|B(\mathbf{w})\|_\nu^2 \right] \leq \mathbb{E}_q \left[\|B(\mathbf{w})\|_D^2 \right] - \mathbb{E}_q [v(\mathbf{w})] + \frac{\lambda}{N} KL(q||p) + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \quad (17)$$

225

$$\mathbb{E}_q \left[\|B(\mathbf{w})\|_D^2 \right] \leq \mathbb{E}_q \left[\|B(\mathbf{w})\|_\nu^2 \right] + \mathbb{E}_q [v(\mathbf{w})] + \frac{\lambda}{N} KL(q||p) + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \quad (18)$$

226 *Proof.* From Hoeffding's inequality we have:

$$P \left(\left| \mathbb{E}_{\nu, \mathcal{P}} \left[\|B(\mathbf{w})\|_D^2 \right] - \|B(\mathbf{w})\|_D^2 \right| > \epsilon \right) \leq 2 \exp \left(- \frac{2N\epsilon^2}{\left(2 \frac{R_{max}}{1-\gamma} \right)^4} \right)$$

227 which implies that, for any $\delta > 0$, with probability at least $1 - \delta$:

$$\left| \mathbb{E}_{\nu, \mathcal{P}} \left[\|B(\mathbf{w})\|_D^2 \right] - \|B(\mathbf{w})\|_D^2 \right| \leq 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}}$$

228 Under independence assumptions, the expected TD error can be re-written as:

$$\begin{aligned} \mathbb{E}_{\nu, \mathcal{P}} \left[\|B(\mathbf{w})\|_D^2 \right] &= \mathbb{E}_{\nu, \mathcal{P}} \left[\frac{1}{N} \sum_{i=1}^N (r_i + \gamma \min_{a'} Q_{\mathbf{w}}(s'_i, a') - Q_{\mathbf{w}}(s_i, a_i))^2 \right] \\ &= \mathbb{E}_{\nu, \mathcal{P}} \left[(R(s, a) + \gamma \min_{a'} Q_{\mathbf{w}}(s', a') - Q_{\mathbf{w}}(s, a))^2 \right] \\ &= \mathbb{E}_\nu \left[\mathbb{E}_{\mathcal{P}} [b(\mathbf{w})^2] \right] \\ &= \mathbb{E}_\nu \left[\text{Var}_{\mathcal{P}} [b(\mathbf{w})] + \mathbb{E}_{\mathcal{P}} [b(\mathbf{w})]^2 \right] \\ &= v(\mathbf{w}) + \|B(\mathbf{w})\|_\nu^2 \end{aligned}$$

229 where $v(\mathbf{w}) \triangleq \mathbb{E}_\nu [\text{Var}_{\mathcal{P}} [b(\mathbf{w})]]$. Thus:

$$\left| \|B(\mathbf{w})\|_\nu^2 + v(\mathbf{w}) - \|B(\mathbf{w})\|_D^2 \right| \leq 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \quad (19)$$

230 From the change of measure inequality [], we have that, for any measurable function $f(\mathbf{w})$ and any
 231 two probability measures p and q :

$$\log \mathbb{E}_p [e^{f(\mathbf{w})}] \geq \mathbb{E}_q [f(\mathbf{w})] - KL(q||p)$$

232 Thus, multiplying both sides of (19) by $\lambda^{-1}N$ and applying the change of measure inequality with
 233 $f(\mathbf{w}) = \lambda^{-1}N \left| \|B(\mathbf{w})\|_\nu^2 + v(\mathbf{w}) - \|B(\mathbf{w})\|_D^2 \right|$, we obtain:

$$\mathbb{E}_q [f(\mathbf{w})] - KL(q||p) \leq \log \mathbb{E}_p [e^{f(\mathbf{w})}] \leq 4 \frac{R_{max}^2 \lambda^{-1}N}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}}$$

Find a ref-
erence for
this

234 where the second inequality holds since the right-hand side of (19) does not depend on \mathbf{w} . Finally,
 235 we can explicitly write:

$$\mathbb{E}_q \left[\left\| B(\mathbf{w}) \right\|_\nu^2 + v(\mathbf{w}) - \left\| B(\mathbf{w}) \right\|_D^2 \right] \leq \frac{\lambda}{N} KL(q||p) + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}}$$

236 from which the lemma follows straightforwardly. \square

237 **Lemma 3.** Let p be a prior distribution over the parameter space \mathcal{W} , and ν be a probability measure
 238 over $\mathcal{S} \times \mathcal{A}$. Assume $\hat{\xi}$ is the minimizer of $ELBO(\xi) = \mathbb{E}_{q_\xi} \left[\left\| B(\mathbf{w}) \right\|_D^2 \right] + \frac{\lambda}{N} KL(q_\xi||p)$ for a
 239 dataset D of N samples. Define $v(\mathbf{w}) \triangleq \mathbb{E}_\nu [Var_{\mathcal{P}} [b(\mathbf{w})]]$. Then, for any $\delta > 0$, with probability at
 240 least $1 - \delta$:

$$\mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_\nu^2 \right] \leq \inf_{\xi \in \Xi} \left\{ \mathbb{E}_{q_\xi} \left[\left\| B(\mathbf{w}) \right\|_\nu^2 \right] + \mathbb{E}_{q_\xi} [v(\mathbf{w})] + 2 \frac{\lambda}{N} KL(q_\xi||p) \right\} + 2 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{N}}$$

241 *Proof.* Let us use Lemma 2 for the specific choice $q = q_{\hat{\xi}}$. From Eq. (17), we have:

$$\begin{aligned} \mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_\nu^2 \right] &\leq \mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_D^2 \right] - \mathbb{E}_{q_{\hat{\xi}}} [v(\mathbf{w})] + \frac{\lambda}{N} KL(q_{\hat{\xi}}||p) + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \\ &\leq \mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_D^2 \right] + \frac{\lambda}{N} KL(q_{\hat{\xi}}||p) + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \\ &= \inf_{\xi \in \Xi} \left\{ \mathbb{E}_{q_\xi} \left[\left\| B(\mathbf{w}) \right\|_D^2 \right] + \frac{\lambda}{N} KL(q_\xi||p) \right\} + 4 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{2N}} \end{aligned}$$

242 where the second inequality holds since $v(\mathbf{w}) > 0$, while the equality holds from the definition of $\hat{\xi}$.

243 We can now use Eq. (18) to bound $\mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_D^2 \right]$, thus obtaining:

$$\mathbb{E}_{q_{\hat{\xi}}} \left[\left\| B(\mathbf{w}) \right\|_\nu^2 \right] \leq \inf_{\xi \in \Xi} \left\{ \mathbb{E}_{q_\xi} \left[\left\| B(\mathbf{w}) \right\|_\nu^2 \right] + \mathbb{E}_{q_\xi} [v(\mathbf{w})] + 2 \frac{\lambda}{N} KL(q_\xi||p) \right\} + 2 \frac{R_{max}^2}{(1-\gamma)^2} \sqrt{\frac{\log \frac{2}{\delta}}{N}}$$

244 This concludes the proof. \square