

Konstanter Faktor : y = ku.


$$G = k$$
  $k = -5$   $k = 1$   $k = 10$ 



Reeller Pol :  $\dot{y} + ay = u$ .


$$G = \frac{1}{s+a} \qquad \qquad a = -1 \quad a = 0 \quad a = +1$$
 
$$\omega_g = |a| \quad \Rightarrow \qquad \qquad \angle G(j\omega_g) = -45/ - 135^\circ$$
 
$$|G(j\omega_g)| = \frac{1}{\sqrt{2}} \approx -3 \text{ dB}$$
 
$$\omega \ll \omega_g : \qquad \qquad |G| \approx \frac{1}{|a|}$$
 
$$\omega_g \ll \omega : \qquad \qquad |G| \propto -20 \text{ dB/Dek.}$$






Konjugiert komplexes Polpaar :  $\ddot{y} + 2a\dot{y} + by = u$ .


$$\frac{1}{s^2+2\zeta\omega_0s+\omega_0^2} \qquad \qquad \omega_0=1 \ \zeta=1 \ \zeta=0.5 \ \zeta=0.1$$
 
$$\angle G(j\omega_0)=-90 \ ^\circ$$
 
$$\omega\ll\omega_0: \qquad \qquad |G|\approx 1/\omega_0^2 \qquad \rightarrow \text{ Konstante Discipling}$$
 
$$\omega_0\ll\omega: \qquad |G|\propto -40 \ ^{\rm dB/Dek}.$$
 
$$M_p=e^{-\pi\zeta/\sqrt{1-\zeta^2}} \qquad \qquad {\rm relatives} \ \ddot{\rm U}{\rm berschiessen} \ \frac{\hat{h}-h_\infty}{h_\infty}$$






Reelle Nullstelle :  $y = \dot{u} + au$ .

G = s + aa = -1 a = 0 a = +1 $\angle G(j\omega_g) = +45/+135^{\circ}$  $\omega_g = |a| \quad \Rightarrow$  $|G(j\omega_g)| = \sqrt{2} \approx +3 \text{ dB}$  $\omega \ll \omega_g$ :  $|G| \approx |a|$  $|G| \propto +20^{\text{dB}}/\text{Dek}$ .  $\angle G \approx 50^{\circ}$  $\omega_g \ll \omega$ :





Konjugiert komplexes Nullstellenpaar :  $y = \ddot{u} + a\dot{u} + bu$ .

## Nor theoretisch möglich

$$s^2 + 2\zeta\omega_0 s + \omega_0^2$$

$$\omega_0 = 1 \ \zeta = 1 \ \zeta = 0.5 \ \zeta = 0.1$$

$$\angle G(j\omega_0) = +90]$$
°


 $\omega \ll \omega_0$ :


$$|G| \approx \omega_0^2$$


$$\omega_0 \ll \omega$$
:


$$|G| \propto +40 \text{ dB/Dek.}$$


## Pol-Nullstellenplan












Totzeit, Verzögerung :  $y(t) = u(t - \tau)$ .

$$G = e^{-s\tau} \quad \tau > 0$$

$$\angle G(j\omega) = -\omega\tau$$

$$|G| = 1 = 0 \text{ dB}$$

$$e^{-s\tau} = \frac{1}{\lim_{n \to \infty} (1 + s\frac{\tau}{n})^n}$$



VERKETTUNG - BEISPIEL.


$$G = \frac{10s+1}{s^3+s^2+25s} = \underbrace{10}_{G_1} \cdot \underbrace{(s+0.1)}_{G_2} \cdot \underbrace{\frac{1}{s}}_{G_3} \cdot \underbrace{\frac{1}{s^2+s+25}}_{G_4}$$


$$G_1 = 10$$

$$G_2 = (s+0.1)$$

$$G_3 = \frac{1}{s}$$

$$G_4 = \frac{1}{s^2+s+25} = \frac{1}{s^2+2\zeta\omega_0s+\omega_0^2} \quad \text{mit} \quad \omega_0 = 5 , \ \zeta = 0.1$$





Addition - Beispiel PID.

$$u = k_p e + k_i \int_0^t e(\tau) d\tau + k_d \frac{de}{dt}$$

$$u = k_p e + k_i \frac{1}{s} e + k_d s e$$

$$G = \frac{u}{e} = k_p + k_i \frac{1}{s} + k_d s = \frac{k_p s + k_i + k_d s^2}{s} = k \frac{(1 + sT_1)(1 + sT_2)}{s}$$

Graphen für  $k=1, T_1=0.1, T_2=10$  sowie  $k_p=10.1, k_i=1, k_d=1$ 

