

# 1 Introduction

- Initial Problem (graph of obs data)
- First thoughts (Hypothesis)

# 2 Method

- Which Method did I use and why?

**For this approach I used two programs: main.py and lumiradius.py.**

I had the choice between two different approaches: A montecarlo approach and a grid approach.

In my program I use the grid approach. I span a grid in distance-mass-age. The range of these three variables can be easily adjusted, as can the binning. In accordance to the observational data I try to explain, distance goes from 0 to  $r_{max} = 3kpc$ . Mass will also be in the same parameters, as the observational data: from  $M_{min} = 5M_{\odot}$  to  $M_{max} = 50M_{\odot}$ . The mass-distribution will follow the Salpeter Initial Mass Function (IMF)(see Salpeter, 1955). Because the IMF follows an inverse power law, I use a logarithmic mass-grid.

Age can be implemented in two different ways: I could either use a separate age-axis for every star going from 0 to  $t_{ms}(M)$  or I can use a single axis for all stars spanning from 0 to  $t_{ms}(M_{min})$ . Because the main sequence age ( $t_{ms}$ ) of a star is a strictly monotonic increasing function of M  $t_{ms}(M_{min}) = t_{max}$ . **I use the second approach.** With  $M_{min} = 5M_{\odot}$  this translates to  $t_{max} = 104Myr$ . This Axis will also be logarithmic to make sure massive stars with small  $t_{ms}$  are correctly represented.

The three informations I have about any given star, are its mass, its age and its distance from earth. From these three informations I need to derive its fractional main sequence age ( $\tau$ ), its apparent magnitude (V) and ultimately the probability density for all stars.

First I will implement a stellar evolution model by Hurley, Pols, & Tout (2000) which approximates the stellar evolution as a function of initial mass ( $M_{ini}$ ), fractional main sequence age ( $\tau$ ) and metallicity(Z).

To do this, first I need to find a function for  $\tau$ . Using equation 5 (Hurley et al., 2000, page 547) I know the main sequence age ( $t_{ms}$ ) and  $\tau$  then becomes:  $\tau = \frac{t}{t_{ms}}$ . Since I only include stars on the main sequence, I can safely include the condition:  $\tau < 1$  to cut down on computing time. Equation 12 and 13 from the same paper are very powerful equations to compute luminosity and radius for a star on the main sequence.

This Paper does reference Tout, Pols, Eggleton, & Han (1996) for Zero Age Mainsequence (ZAMS) Radii and Luminosities. However there is one error in this Paper I had to correct. In Equation 1  $\gamma + M^3$  has to be  $\gamma \cdot M^3$ .

**insert HRD illustrating lumiradius**

The program does allow for freely changeable metallicities. For my purposes I use  $Z=0.02$  for all stars to simulate a metallicity similar to that of our galactic neighborhood.

I now run this program for every possible mass-age tuple and save the results in a matrix. This way I don't have to call the program for every distance-mass-age triple and massively cut down on computing time.

With all this I now know distance, mass, age, fractional main sequence age, luminosity and radius for any given star. I can now use these informations to compute the apparent magnitudes.

$$M_V = V - 5 \cdot \log_{10}(\text{distance}) + 5 - \text{Red} \quad (1)$$

$$M_{bol} = M_V + BC \quad (2)$$

$$\frac{L}{L_{\odot}} = 0.4 \cdot (4.72 - M_{bol}) \quad (3)$$

Where  $M_V$  is the absolute visual magnitude, Red is the reddening as a function of distance,  $M_{bol}$  is the absolute bolometric magnitude and BC is the Bolometric Correction. Using equations 1, 2 and 3 I can now compute the apparent visual magnitude V:

$$V = 5 \cdot \log_{10}(\text{distance}) - 5 + \text{Red} + 4.72 - \frac{L}{L_{\odot} \cdot 0.4} - BC \quad (4)$$

To get a rough approximation for reddening, I use Amôres & Lépine (2005, Figure 9) and interpolate between these values 1:0.9, 2:2.25 and 3:3.273

The next thing I need to know are the probability densities for stars in space  $\left(\frac{dp}{dV}\right)$ , mass  $\left(\frac{dp}{dm}\right)$  and age  $\left(\frac{dp}{dt}\right)$ . In my simulation I assume a homogenous distribution of stars. This makes finding a probability density for space very easy. Because of the radial symmetry of my problem the distribution becomes solely dependant on the distance from earth:

$$\frac{dp}{dV} = \frac{1}{V_{tot}} = \frac{1}{\frac{4}{3} \cdot \pi \cdot \text{maxdistance}^3} \quad (5)$$

I assume a constant star formation rate, so the probability density in age would be similar to the density in distance. I do however use a logarithmic binning in age. Thus I can not simply use  $\frac{dp}{dt}$  but instead need to find  $\frac{dp}{d \log t}$  using the following:  $\frac{dp}{dt} = \frac{1}{t_{ms}} \quad \wedge \quad d \log t = \frac{dt}{t \ln 10}$

$$\frac{dp}{d \log t} = \frac{t \ln 10}{t_{ms}} \quad (6)$$

I assume, that the stars are distributed in mass following the Salpeter IMF:  $\frac{dp}{dM} = A \cdot M^{-2.35}$  (Salpeter, 1955). Similar to the density function in age, I need to convert it for my logarithmic binning using  $d \log M = \frac{dM}{M \ln 10}$

$$\frac{dp}{d \log M} = \ln 10 \cdot A \cdot M^{-1.35} \quad (7)$$

Where A is a normalization factor, that needs to be computed.

$$1 = \int_{M_{min}}^{M_{max}} A \cdot M^{-2.35} dM = [-1.35 \cdot A \cdot M^{-1.35}]_{M_{min}}^{M_{max}} \quad (8)$$

$$A = \frac{1.35}{M_{min}^{-1.35} - M_{max}^{-1.35}} \quad (9)$$

With this information I can now formulate an overarching probability density function with regard to  $\tau$

$$\frac{dp}{d\tau} = \frac{dp}{dV} dV \cdot \frac{dp}{d \log t} d \log t \cdot \frac{dp}{d \log M} d \log M \cdot \frac{1}{d\tau} \quad (10)$$

I now split  $\tau$  in 20 bins. For every possible star in my sample I will check  $\tau \leq 1$  and  $V \leq 9$ . This way I count every star on the main sequence, that falls below my magnitude cut.

The binning can be freely adjusted. To optimize runtime of my program I conducted a few tests to adjust the binnings in all three dimensions.

### 3 testing stuffs

I




Abbildung 1: Difference between computed and observed magnitudes as a function of fractional main sequence age (left) and total age (right). The theoretical magnitudes were computed using stellar radius, luminosity and distance.

Figure 1 shows the difference between computed and observed magnitudes as a function of fractional main sequence age (left) and total age (right). It shows, that for some stars there is an offset to higher  $V_{comp}$  at high  $\tau$  and low absolute ages. This implies, that this disagreement shows up in high-mass stars near TAMS. This can be explained using (Castro et al., 2014, figure 1). My code uses an evolutionary model similar to the one used at the lefthand side.  $V_{obs}$  however is obtained using a model similar to the righthand side. This subsequently means, that TAMS will shift to lower  $T_{eff}$  and higher  $L/L_{\odot}$ .




Abbildung 2: Difference between computed and observed magnitudes as a function of Mass (left) and  $\log_{10}(\text{distance})$  (right). The theoretical magnitudes were computed using stellar radius, luminosity and distance.

Figure 2




Abbildung 3: Probability density function for all stars in my sample with a magnitude cut at  $V=9$  with binnings of 100-100-100 (left) and 50-50-50 (right) in distance-mass-age.




Abbildung 4: Probability density function for all stars in my sample with a magnitude cut at  $V=9$  with binnings of 100-50-100 (left) and 100-100-50 (right) in distance-mass-age.




Abbildung 5: Probability density function for all stars in my sample with a magnitude cut at  $V=9$  with binnings of 50-100-100 (left) and 10-100-100 (right) in distance-mass-age.

## 4 ending

### Literatur

Amôres, E. B., & Lépine, J. R. D. 2005, *AJ*, 130, 659

Castro, N., Fossati, L., Langer, N., et al. 2014, *A&A*, 570, L13

Hurley, J. R., Pols, O. R., & Tout, C. A. 2000, *MNRAS*, 315, 543

Salpeter, E. E. 1955, *ApJ*, 121, 161

Tout, C. A., Pols, O. R., Eggleton, P. P., & Han, Z. 1996, *MNRAS*, 281, 257