Exerciseses chapter 4

Andreas Slåttelid

23/2/2022

Contents

Exercise $4.2 + 4.3$
Exercise 4.4 Term insurance in discrete time
Exercise 4.5 Permanent disability insurance discrete time
Exercise 4.5 i)
Exercise 4.6 (Endowment policy in discrete time)
Exercise 4.7 Discrete vs continous time endowment insurance

Exercise 4.2 + 4.3

In this exercise we are considering a disability insurance, hence $S = \{*, \diamond, \dagger\}$. The contractual information given are: as follows:

```
x <- 30  #age
T <- 35  #length of contract
D <- 20000  #disability pension
r <- 0.03  #intereset rate
premium_yearly <- 2500</pre>
```

Furthermore the transition rates are given by:

```
lambda <- function(t){
    #we do not have any t-dependency
    m12 <- 0.0279
    m13 <- 0.0229
    m11 <- -(m12+m13)
    m21 <- 0
    m23 <- 0.0229
    m22 <- -(m21+m23)
    m31 <- m32 <- m33 <- 0
    L <- matrix(c(m11,m12,m13,m21,m22,m23,m31,m32,m33), nrow=3, byrow=TRUE)
    return(L)
}</pre>
```

We also have the following policy-functions:

$$a_*(t) = \begin{cases} 0, & t < 0 \\ -\pi t, & t \in [0, T) \\ -\pi T, & t \ge T \end{cases}$$

$$a_{\diamond}(t) = \begin{cases} 0, & t < 0 \\ Dt, & t \in [0, T) \\ DT, & t \ge T \end{cases}$$

We were asked to calculate $V_i^+(t,A)$ for $j=*,\diamond$ and for t=30. We then get:

$$V_*^+(t,A) = \frac{1}{v(t)} \left[\int_t^T v(s) p_{**}(x+t,x+s) da_*(s) + \int_t^T v(s) p_{*\diamond}(x+t,x+s) da_{\diamond}(s) \right]$$
$$= \frac{1}{v(t)} \left[-\pi \int_t^T v(s) p_{**}(x+t,x+s) ds + D \int_t^T v(s) p_{*\diamond}(x+t,x+s) ds \right]$$

$$V_{\diamond}^{+}(t,A) = \frac{1}{v(t)} \left[\int_{t}^{T} v(s) p_{\diamond \diamond}(x+t,x+s) da_{\diamond}(s) \right]$$
$$= \frac{1}{v(t)} \left[D \int_{t}^{T} v(s) p_{\diamond \diamond}(x+t,x+s) ds \right]$$

We will now use an Euler-scheme to simulate what's going on, and then use a Riemann-sum to compute the integrals:

$$\int_{a}^{b} f(x)dx \approx \sum_{i=0}^{n-1} f(x_{i}^{*}) \Delta x$$

Where $\Delta x = \frac{b-a}{n}$, which in our case corresponds to: h and $x_i^* \in [x_i, x_{i+1}]$

```
field <- function(t,M){
  return(M%*%lambda(t))
#field(t0, P0)
Euler <- function(t0,P0,h,tn){</pre>
  if(t0==tn){ return(P0)}
  N \leftarrow (tn-t0)/h
  D <- \dim(PO) [1] #qives dimension of man matrix dim = c(m,n)
  #Initial condition at s
  P \leftarrow array(diag(D*(N+1)), dim=c(D,D,N+1))
  #First iteration
  P[,,1] \leftarrow P0
  for(n in 1:N){
    P[,,n+1] \leftarrow P[,,n] + h \cdot field(t0 + n \cdot h, P[,,n])
  return(P) #returns array, array(data, dim=c(3,3,2)), this stores 2 3x3 matricies
t0 <- 60
                #start age
PO <- diag(3) #initial start with P(s,s) = I
tn <- 110
               #end age
h <- 1/12
               #step size
N <- (tn-t0)/h #number of steps
sol <- Euler(t0,P0,h,tn) #contains ages transition probs from 60 to 65.
v <- function(t){</pre>
  return(exp(-r*t))
#v(s), but s in [30,35]
v_ages <- sapply(seq(30, 35, by = h), v) #we apply the function v to each element
```

```
\textit{\#sum1} \textit{ where we do not care about the premium until afterwards, we use Riemann-sum.}
s1 <- 0
for (i in 1:(length(v_ages)-1)){
 s1 <- s1 + v_ages[i]*sol[1,1, ][i]*h #survival i.e * -> *
\#sum2: do not care about D=20.000 until afterwards
s2 <- 0
for (i in 1:(length(v_ages)-1)){
  s2 <- s2 + v_ages[i]*sol[1,2,][i]*h #disability i.e * -> dis
#V*(30, A):
V_30_A \leftarrow (1/v(30))*((-1)*premium_yearly*s1 + D*s2)
V_30_A
## [1] -4781.495
#V_[disabeld](30,A):
s3 <- 0
for (i in 1:(length(v_ages)-1)){
 s3 <- s3 + v_ages[i]*sol[2,2,][i]*h #disability i.e dis -> dis
#V_[disabeld](30,A):
V_30_{dis} \leftarrow (1/v(30))*D*s3
V_30_dis
```

[1] 88057.1

Exercise 4.4 Term insurance in discrete time

In this exercise we are dealing with a term insurance in discrete time, hence: $S = \{*, \dagger\}$, the contractual information provided:

x <- 50 T <- 10 B <- 200000 r <- 0.025

We are asked to use the equivalence principle to determine the fair premium π , we start of by describing the policy functions:

$$a_*^{Pre}(n) = \begin{cases} -\pi & , n = 0, \dots, T - 1 \\ 0 & , else \end{cases}$$
$$a_{*\dagger}^{Post}(n) = \begin{cases} B & , n = 0, \dots, T - 1 \\ 0 & , else \end{cases}$$

Furthermore the formula for directe reserves are given by:

$$V_i^+(t,A) = \sum_j \sum_{n \ge t} \frac{v(n)}{v(t)} p_{ij}^x(t,n) a_j^{Pre}(n) + \sum_{k \ne j} \sum_{n \ge t} \frac{v(n+1)}{v(t)} p_{ij}^x(t,n) p_{jk}^x(n,n+1) a_{jk}^{Post}(n)$$

In our case, this translates to:

$$V_*^+(t,A) = -\pi \sum_{n=t}^{T-1} \frac{v(n)}{v(t)} p_{**}^x(t,n) + B \sum_{n=t}^{T-1} \frac{v(n+1)}{v(t)} p_{**}^x(t,n) p_{*\dagger}^x(n,n+1)$$

The equivalence principle states that we should choose π so that $V_*^+(0,A)=0$, giving us:

$$0 = -\pi \sum_{n=0}^{T-1} v(n) p_{**}^{x}(0,n) + B \sum_{n=0}^{T-1} v(n+1) p_{**}^{x}(0,n) p_{*\dagger}^{x}(n,n+1)$$

$$\downarrow \downarrow$$

$$\pi = \frac{B \sum_{n=0}^{T-1} v(n+1) p_{**}^{x}(0,n) p_{*\dagger}^{x}(n,n+1)}{\sum_{n=0}^{T-1} v(n) p_{**}^{x}(0,n)}$$

```
#0: alive, 1:dead
mu01 <- function(u){</pre>
  a < -0.002
  b <- 0.0005
  return(a + b*(u-50))
p_surv <- function(t,s){</pre>
 f <- mu01
  integral <- integrate(f, lower = t, upper = s)$value</pre>
  return(exp((-1)*integral))
}
#discount-factor
v <- function(t){</pre>
  return(exp(-r*t))
}
s1_above <- 0
for (n in 0:(T-1)){
  s1\_above \leftarrow s1\_above + v(n+1)*p\_surv(x, x+n)*(1-p\_surv(x+n,x+n+1))
s2\_below \leftarrow 0
for (n in 0:(T-1)){
  s2_below <- s2_below + v(n)*p_surv(x,x+n)</pre>
above1 <- B*s1_above
below2 <- s2_below
prem <- above1/below2</pre>
prem
```

[1] 852.2476

Exercise 4.5 Permanent disability insurance discrete time

We have the following state space $S = \{*, \diamond, \dagger\}$. Furthermore the contractual information provided is:

```
x <- 45
T <- 20
D <- 12000
r <- 0.03

#discount factor
v <- function(t){
   return(exp(-r*t))
}</pre>
```

Exercise 4.5 i)

In this case we have the following transition rates:

$$\mu_{*\diamond}(t) = 0.0279 \ \mu_{*\dagger}(t) = 0.0229 \ \mu_{\diamond\dagger}(t) = \mu_{*\dagger}(t)$$

```
#1: alive

#2: disabled

#3: dead

m12 <- 0.0279

m13 <- 0.0229

m11 <- -(m12+m13)

m21 <- 0

m23 <- 0.0229

m22 <- -(m21+m23)

m31 <- m32 <- m33 <- 0
```

Since reactivation is not allowed, we can actually solve the transition-probabilites directly. We first start with the survival probability:

$$p_{11}(t,s) = \exp((-1) * 0.0508(s-t))$$

```
p_11 <- function(t,s){
  return(exp(-1*0.0508*(s-t)))
}</pre>
```

We describe the policy functions:

$$a_*^{Pre}(n) = \begin{cases} -\pi & , n = 0, \dots, T-1 \\ 0 & , else \end{cases} \quad a_\diamond^{Pre}(n) = \begin{cases} D & , n = 0, \dots, T-1 \\ 0 & , else \end{cases}$$

And we get the following formula for the reserve:

$$V_*^+(t,A) = \frac{1}{v(t)} \left[-\pi \sum_{n=t}^{T-1} v(n) p_{**}^x(t,n) + D \sum_{n=t}^{T-1} v(n) p_{*\diamond}^x(t,n) \right]$$

Now from the equivalence principle we should have that the premium π should be decided so that $V_*^+(0,A)=0$, this leaves us with:

$$\pi = \frac{D\sum_{n=0}^{T-1} v(n) p_{*\diamond}^{x}(0, n)}{\sum_{n=0}^{T-1} v(n) p_{**}^{x}(0, n)}$$

We can use Kolmogorov's equations to find $p_{*\dagger}$ and $p_{*\diamond}$, as reactivation is not allowed, and $\mu_{\diamond\dagger}(t) = \mu_{*\dagger}(t)$

$$\partial_s p_{13}(t,s) = p_{11}(t,s)\mu_{13}(s) + p_{12}(t,s)\mu_{23}(s)$$

Now, lets introduce the notation: $\hat{\mu}(u) = \mu_{23}(u) = \mu_{13}(u)$, we then have:

$$\partial_s p_{13}(t,s) = [p_{11}(t,s) + p_{12}(t,s)] \hat{\mu}(s), \quad (p_{11} + p_{12} + p_{13} = 1)$$

= $[1 - p_{13}(t,s)] \hat{\mu}(s)$

We can now multiply by (-1), and take the derivative of 1 (which is zero), this gives us:

```
#alive to dead
p_13 <- function(t,s){
  return(1-exp((-1)*m13*(s-t)))
}</pre>
```

Exercise 4.6 (Endowment policy in discrete time)

Required-libraries

```
list.of.packages <- c("tidyverse", "scales")
new.packages <- list.of.packages[!(list.of.packages %in% installed.packages()[,"Package"])]
if(length(new.packages)) install.packages(new.packages)
library(tidyverse)
library(scales) #nice formatting of plots</pre>
```

We are dealing with an endownment policy, in discrete time, this means that $S = \{*, \dagger\}$, the contractual information provided is:

futhermore the force of mortality is given by:

```
mu <- function(t){
  return(0.0015 + 0.0004*(t-35))
}</pre>
```

discount-factor:

```
v <- function(t){
  return(exp(-r*t))
}</pre>
```

We can also calulate the survival probability:

$$p_{**}(s,t) = \exp\left(-\int_s^t \mu_{*\dagger}(u)du\right)$$

```
p_surv <- function(s,t){
  integral <- 0.0015*(t-s) + 0.0004*((1/2)*(t^2 -s^2) + 35*(s-t))
  return(exp(-integral))
}
p_surv(35,36)</pre>
```

```
## [1] 0.9983014
```

We are asked to calculate the yearly premium π , but first we start by defining the policy functions specified in the contract.

$$a_*^{Pre}(n) = \begin{cases} -\pi &, n = 0, \dots, T-1 \\ E &, n = T \end{cases} \quad a_{*\dagger}^{Post}(n) \quad = \begin{cases} B &, n = 0, \dots, T-1 \\ 0 &, else \end{cases}$$

We can now start to calculate the reserves, but we must be aware of what happens at n = T:

$$V_*^+(t,A) = \frac{1}{v(t)} \left[\sum_{n=t}^T v(n) p_{**}(x+t,x+n) a_*^{Pre}(n) + \sum_{n=t}^{T-1} v(n+1) p_{**}(x+t,x+n) p_{*\dagger}(x+n,x+n+1) a_{*\dagger}^{Post}(n) \right]$$

$$= \frac{1}{v(t)} \left[\alpha + \beta \right]$$

Here α corresponds to the first sum, and β corresponds to the second sum:

$$\alpha = \sum_{n=t}^{T-1} v(n) p_{**}(x+t, x+n) (-\pi) + v(T) p_{**}(x+t, x+T) E$$
$$\beta = \sum_{n=t}^{T-1} v(n+1) p_{**}(x+t, x+n) p_{*\dagger}(x+n, x+n+1) B$$

We will now use the equivalence principle, which states that the fair premium π should be such that $V_*(0, A) = 0$, and using this method we end up with:

$$\pi = \frac{v(T)p_{**}(x, x+T)E + B\sum_{n=0}^{T-1} v(n+1)p_{**}(x, x+n)p_{*\dagger}(x+n, x+n+1)}{\sum_{n=0}^{T-1} v(n)p_{**}(x, x+n)}$$

```
res1 <- 0
for (n in 0:(T-1)){
    res1 <- res1 + v(n+1)*p_surv(x, x+n)*(1-p_surv(x+n, x+n+1))
}

res2 <- 0
for (n in 0:(T-1)){
    res2 <- res2 + v(n)*p_surv(x, x+n)
}

above <- B*res1 + v(T)*p_surv(x,x+T)*E #whats above in the premium expression
below <- res2  #whats below in the premium expression

premium_yearly <- above/below
premium_yearly</pre>
```

[1] 4095.413


We now also want the prospective reserves: i.e $V_*^+(t,A)$ for $t=0,\ldots,T$:

```
#Present value of reserve when including premiums.
V_reserve_1 <- function(x=35, E = 125000 , T=25){
    #x: age of insured
    #B: benefit specified in contract
    #T: length of contract

V_star_pre <- NULL
    for (t in 0:(T-1)){
        s <- 0
        for (n in t:(T-1)){
            s <- s + (1/v(t))*(-1)*premium_yearly*v(n)*p_surv(t+x, n+x)
        }
        V_star_pre[t+1] <- s + (1/v(t))*v(T)*p_surv(x+t, x+T)*E
    }
    V_star_pre[T+1] <- E #specified by contract
    return(V_star_pre)
}</pre>
```

```
V_{reserve_2} \leftarrow function(x=35, B=250000, T=25){
  #x: age of insured
  #B: benefit specified in contract
  #T: length of contract
  V_star_post <- NULL</pre>
  for (t in 0:(T-1)){
    s <- 0
    for (n in t: (T-1)){
      s \leftarrow s + (1/v(t))*B*v(n+1)*p_surv(x+t, x+n)*(1-p_surv(x+n, x+n+1))
    }
    V_star_post[t+1] <- s</pre>
  }
  V_star_post[T+1] <- 0 #specified by contract</pre>
  return(V_star_post)
}
reserve_total <- V_reserve_1() + V_reserve_2()</pre>
reserve_total
```

```
## [1] 1.091394e-11 3.823145e+03 7.692302e+03 1.161236e+04 1.558847e+04 ## [6] 1.962604e+04 2.373075e+04 2.790862e+04 3.216595e+04 3.650944e+04 ## [11] 4.094613e+04 4.548348e+04 5.012938e+04 5.489218e+04 5.978071e+04 ## [16] 6.480437e+04 6.997309e+04 7.529742e+04 8.078858e+04 8.645848e+04 ## [21] 9.231977e+04 9.838591e+04 1.046712e+05 1.111910e+05 1.179615e+05 ## [26] 1.250000e+05
```



Exercise 4.7 Discrete vs continous time endowment insurance

In this exercise we are considering an endowment insurance, hence $S = \{*, \dagger\}$, the contractual information provided is:

x <- 30 T <- 35 E <- 100000 B <- 200000 r <- 0.035

We start with discrete time, the policy functions are as follows:

$$a_*^{Pre}(n) = \begin{cases} -\pi & , n = 0, \dots, T-1 \\ E & , n = T \end{cases}$$
$$a_{*\dagger}^{Post}(n) = \begin{cases} B & , n = 0, \dots, T-1 \\ 0 & , else \end{cases}$$

Again:

$$V_i^+(t,A) = \sum_{j} \sum_{n > t} \frac{v(n)}{v(t)} p_{ij}^x(t,n) a_j^{Pre}(n) + \sum_{k \neq j} \sum_{n > t} \frac{v(n+1)}{v(t)} p_{ij}^x(t,n) p_{jk}^x(n,n+1) a_{jk}^{Post}(n)$$

Which translates to:

$$V_*^+(t,A) = -\pi \sum_{n=t}^{T-1} \frac{v(n)}{v(t)} p_{**}^x(t,n) + \frac{v(T)}{v(t)} p_{**}^x(t,T) E + B \sum_{n=t}^{T-1} \frac{v(n+1)}{v(t)} p_{**}^x(t,n) p_{*\dagger}^x(n,n+1)$$

We then use the equivalence principle, telling us that the fair premium π should be chosen so that $V_*^+(0,A)=0$:

$$\pi = \frac{v(T)p_{**}^{x}(0,T)E + B\sum_{n=0}^{T-1}v(n+1)p_{**}^{x}(0,n)p_{*\dagger}^{x}(n,n+1)}{\sum_{n=0}^{T-1}v(n)p_{**}^{x}(0,n)}$$

We now discuss the situation in continous time, we then get the following policy functions:

$$a_*(t) = \begin{cases} 0, & t < 0 \\ -\pi t, & t \in [0, T) \\ -\pi T + E, & t \ge T \end{cases} \quad a_{*\dagger}(t) = \begin{cases} B, & t \in [0, T) \\ 0, & else \end{cases}$$

We start off by stating the general formula for reserves in continuous-time:

$$V_i^+(t,A) = \sum_j \frac{v(T)}{v(t)} p_{ij}^x(t,T) \Delta a_j(T) + \sum_j \int_t^T \frac{v(s)}{v(t)} p_{ij}^x(t,s) da_j(s) + \sum_{k \neq j} \int_t^T \frac{v(s)}{v(t)} p_{ij}^x(t,s) \mu_{jk}^x(s) a_{jk}(s) ds$$

Translating this into our situation:

$$V_*^+(t,A) = \frac{v(T)}{v(t)} p_{**}^x(t,T) E - \pi \int_t^T \frac{v(s)}{v(t)} p_{**}^x(t,s) ds + B \int_t^T \frac{v(s)}{v(t)} p_{**}^x(t,s) \mu_{*\dagger}^x(s) ds$$

Now as always, the equivalence principle tells us to chose π such that $V_*^+(0,A)=0$:

$$\pi = \frac{v(T)p_{**}^{x}(0,T)E + B\int_{0}^{T}v(s)p_{**}^{x}(0,s)\mu_{*\dagger}^{x}(s)ds}{\int_{0}^{T}v(s)p_{**}^{x}(0,s)ds}$$

```
#want the map function from purrr:
library(tidyverse)
packageVersion("tidyverse")
## [1] '1.3.1'
#0: alive, 1:dead
mu01 <- function(t){
  a2 < -5*10^{-4}
  b2 <- 7.5858*10^{-5}
  c2 <- 0.087498
  ans \leftarrow a2 + b2*exp(c2*t)
  return(ans)
}
p_surv <- function(t, s){</pre>
  f <- Vectorize(mu01)</pre>
  integral <- integrate(f, lower = t, upper = s)$value</pre>
  ans <- exp(-integral)</pre>
  return(ans)
}
#discount factor
v <- function(t){</pre>
  return(exp(-r*t))
#premium discrete time:
summand_upper <- function(n){</pre>
  v(n+1)*p_surv(x, x + n)*(1- p_surv(x + n, x + n +1))
}
summand_lower <- function(n){</pre>
  v(n)*p_surv(x, x +n)
}
#map_dbl: maps the function summand_upper to the elements 0:(T-1)
sum_upper <- sum(map_dbl(0:(T-1), summand_upper))</pre>
sum_lower <- sum(map_dbl(0:(T-1), summand_lower))</pre>
above_discrete <- v(T)*p_surv(x, x + T)*E + B*sum_upper
below_discrete <- sum_lower</pre>
premium_discrete <- above_discrete/below_discrete</pre>
premium_discrete
```

[1] 2204.58

```
#premium continous time
integrand_upper <- function(s){
   v(s)*p_surv(x, x + s)*mu01(x + s)
}
integrand_lower <- function(s){
   v(s)*p_surv(x, x + s)
}

f1 <- Vectorize(integrand_upper)
f2 <- Vectorize(integrand_lower)

integral_upper <- integrate(f1, lower = 0, upper = T)$value
integral_lower <- integrate(f2, lower = 0, upper = T)$value
above_cont <- v(T)*p_surv(x, x + T)*E + B*integral_upper
below_cont <- integral_lower

premium_cont <- above_cont/below_cont
premium_cont</pre>
```

[1] 2268.052

We then recall what the premiums were in the different cases:

We then recall what the premiums were:

```
premium_discrete
```

```
## [1] 2204.58
premium_cont
```

```
## [1] 2268.052
```