


# Intro to Math and Python

Lecture 1



**girafe**  
**ai**



# Today



1. Course overview
2. Linear Algebra
  - Core objects
  - Vector spaces
3. A bit of Analytic Geometry
  - Orthogonal projections
  - Hyperplanes
  - Normals

# About this course




Image source: Mathematics for Machine Learning, p. 14  
(<https://mml-book.github.io/book/mml-book.pdf>)

# About this course



This week:

1. Linear algebra
2. Calculus
3. Probability theory

Prerequisites:

- basic knowledge of math;
- some Python.

# About me

Iurii Efimov

✉ [yuri.efimov@phystech.edu](mailto:yuri.efimov@phystech.edu)

in [efimov-iurii](https://www.linkedin.com/in/efimov-iurii)

◦ PhD researcher



◦ Lecturer



◦ Artec3D DS Team



# **Linear Algebra: the Basics**



# Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$  - a scalar *Example:  $-2$*

# Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$  - a scalar *Example:  $-2$*
- $x \in \mathbb{R}^n$  - a vector with  $n$  entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$



# Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$  - a scalar *Example:  $-2$*
- $x \in \mathbb{R}^n$  - a vector with  $n$  entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

# Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$  - a scalar *Example:  $-2$*
- $x \in \mathbb{R}^n$  - a **vector** with  $n$  entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns


$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

# What are Vectors?

- Ordered sets of numbers:  $x = [1, 2]$

# What are Vectors?


- Ordered sets of numbers:  $x = [1, 2]$
- A point with Cartesian coordinates




# What are Vectors?

- Ordered sets of numbers:  $x = [1, 2]$

- A point with Cartesian coordinates



- Direction + length



# Vector Spaces



# Vector Space: Definition

- A real-valued **vector space**  $(V, +, \cdot)$  is a set of vectors  $V$  with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

# Vector Space: Definition

- A real-valued **vector space**  $(V, +, \cdot)$  is a set of vectors  $V$  with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

that satisfy the following properties (axioms):

	Property	Meaning
1.	<b>Associativity</b> of addition	$x + (y + z) = (x + y) + z$
2.	<b>Commutativity</b> of addition	$x + y = y + x$
3.	<b>Identity element</b> of addition	$\exists 0 \in V: \forall x \in V \quad 0 + x = x$
4.	<b>Identity element</b> of scalar multiplication	$\forall x \in V \quad 1 \cdot x = x$
5.	<b>Inverse element</b> of addition	$\forall x \in V \exists -x \in V: x + (-x) = 0$
6.	<b>Compatibility</b> of scalar multiplication	$\alpha(\beta x) = (\alpha\beta)x$
7.	<b>Distributivity</b>	$(\alpha + \beta)x = \alpha x + \beta x$
8.		$\alpha(x + y) = \alpha x + \alpha y$



# Let's define vector operations!

# Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

# Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

# Operations with Vectors: Example

$x, y \in \mathbb{R}^3$ :

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

# Operations with Vectors: Example

$x, y \in \mathbb{R}^3$ :

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

# Operations with Vectors: Example

$$x, y \in \mathbb{R}^3:$$

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

Sum:


$$x + y = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$

# **Vector Operations: Geometrical Interpretation**

# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$






# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$

$$-0.5\vec{a} = [-1, -1.5]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




# Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$



# Back to Vector Spaces



# Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

**satisfy axioms (1) – (8)**  
*(check it yourself)*

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

# Vector Spaces

$(\mathbb{R}^n, +, \cdot), n \in \mathbb{N}$  - a vector space with operations

1. vector addition:

$$x + y = (x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n)$$

2. multiplication by a scalar:

$$\alpha x = \alpha(x_1, x_2, \dots, x_n) = (\alpha x_1, \alpha x_2, \dots, \alpha x_n)$$

# Inner Product



# Inner Product

- Inner product is a function  $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$  that satisfies the following properties:
  - *Symmetric*:  $\forall x, y \in V \quad \langle x, y \rangle = \langle y, x \rangle$
  - *Positive definite*:  $\forall x \in V \setminus \{0\} \quad \langle x, x \rangle > 0$  and  $\langle x, 0 \rangle = 0$ .

# Dot Product

- A particular type of inner product.

# Dot Product

- A particular type of inner product.
- For  $x = [x_1, \dots, x_n]$ ,  $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

# Dot Product

- A particular type of inner product.

- For  $x = [x_1, \dots, x_n]$ ,  $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

- Example:

$$x = [1, 2, 3, 4], \quad y = [-1, 0, 1, 2]$$

$$(x, y) = 1 \cdot (-1) + 2 \cdot 0 + 3 \cdot 1 + 4 \cdot 2 = -1 + 0 + 3 + 8 = 10$$

# Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.



# Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$  with dot product  $(\cdot, \cdot)$  is called a *Euclidian vector space*.

# Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$  with dot product  $(\cdot, \cdot)$  is called a *Euclidian vector space*.
- *Note: there're inner products different from dot product.*

# Norms



# Norm

- A **norm** on a vector space  $V$  is a function  $\| \cdot \|: V \rightarrow \mathbb{R}$  which assigns each vector  $x \in V$  its *length*  $\|x\| \in \mathbb{R}$ .

# Norm

- A **norm** on a vector space  $V$  is a function  $\| \cdot \|: V \rightarrow \mathbb{R}$  which assigns each vector  $x \in V$  its *length*  $\|x\| \in \mathbb{R}$ .
- A norm must satisfy the following properties:

# Norm


- A **norm** on a vector space  $\mathbb{V}$  is a function  $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$  which assigns each vector  $x \in \mathbb{V}$  its *length*  $\|x\| \in \mathbb{R}$ .
- A norm must satisfy the following properties:
  - *Absolutely homogeneous*:  $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$

# Norm

- A **norm** on a vector space  $\mathbb{V}$  is a function  $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$  which assigns each vector  $x \in \mathbb{V}$  its *length*  $\|x\| \in \mathbb{R}$ .
- A norm must satisfy the following properties:
  - *Absolutely homogeneous*:  $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
  - *Positive definite*:  $\forall x \in \mathbb{V} \quad \|x\| \geq 0$  and  $\|x\| = 0 \Leftrightarrow x = 0$

# Norm

- A **norm** on a vector space  $\mathbb{V}$  is a function  $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$  which assigns each vector  $x \in \mathbb{V}$  its *length*  $\|x\| \in \mathbb{R}$ .
- A norm must satisfy the following properties:
  - *Absolutely homogeneous*:  $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
  - *Positive definite*:  $\forall x \in \mathbb{V} \quad \|x\| \geq 0$  and  $\|x\| = 0 \Leftrightarrow x = 0$
  - *Triangle inequality*:  $\forall x, y \in \mathbb{V} \quad \|x + y\| \leq \|x\| + \|y\|$





# Examples of Norms

# Manhattan Norm



- A norm for  $x \in \mathbb{R}^n$ :

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$



# Manhattan Norm

- A norm for  $x \in \mathbb{R}^n$ :

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 6$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$



# Manhattan Norm

- A norm for  $x \in \mathbb{R}^n$ :


$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 5$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$





# Euclidian Norm

- A norm for  $x \in \mathbb{R}^n$ :

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



# Euclidian Norm

- A norm for  $x \in \mathbb{R}^n$ :

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



# Euclidian Norm

- A norm for  $x \in \mathbb{R}^n$ :

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$



# Euclidian Norm

- A norm for  $x \in \mathbb{R}^n$ :


$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$





# Other norms

- In general, for  $x = [x_1, \dots, x_n] \in \mathbb{R}^n$  an  $\ell_p$ -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

# Other norms

- In general, for  $x = [x_1, \dots, x_n] \in \mathbb{R}^n$  an  $\ell_p$ -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- $\ell_1$  - Manhattan norm  $\|\cdot\|_1$ ;
- $\ell_2$  - Euclidian norm  $\|\cdot\|$  (default);

# Other norms

- In general, for  $x = [x_1, \dots, x_n] \in \mathbb{R}^n$  an  $\ell_p$ -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- $\ell_1$  - Manhattan norm  $\|\cdot\|_1$ ;
- $\ell_2$  - Euclidian norm  $\|\cdot\|$  (default);
- $\ell_\infty$ :  $\|x\|_\infty = \max_i |x_i|$

*Example:*  $\|[1, 2, 3]\|_\infty = 3$ ,  $\|[1, 0]\|_\infty = 1$ ,  $\|[-1, 0.5]\|_\infty = 1$ .

# Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

# Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

# Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \|x\|_2$$

# Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \|x\|_2$$

- (!) Not every norm is induced by an inner product.  
Example: Manhattan norm.

# Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$



# Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

- For dot product and Euclidian norm:

$$|(x, y)| \leq \|x\|_2 \cdot \|y\|_2$$

# Distance between Vectors

- Distance between two vectors  $x$  and  $y$  is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

# Distance between Vectors

- Distance between two vectors  $x$  and  $y$  is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

- For dot product and Euclidian norm, we get *Euclidian distance*:

$$\begin{aligned} d(x, y) &= \|x - y\|_2 = \sqrt{(x - y, x - y)} = \\ &= \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \dots + (x_n - y_n)^2}. \end{aligned}$$

# Angles and Orthogonality




# Angle between Two Vectors

- Inner product also captures the geometry of vector space by defining the angle between two vectors.
- Remember Cauchy-Schwarz inequality:

$$|(x, y)| \leq \|x\| \cdot \|y\|$$

$$-1 \leq \frac{(x, y)}{\|x\| \cdot \|y\|} \leq 1$$

$$\omega: \cos \omega = \frac{(x, y)}{\|x\| \cdot \|y\|} - \text{angle between } x \text{ and } y.$$



# Angle between Two Vectors: Example

- What is the angle  $\omega$  between  $x = [5, 0]$  and  $y = [1, 1]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

# Angle between Two Vectors: Example

- What is the angle  $\omega$  between  $x = [5, 0]$  and  $y = [1, 1]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle  $\omega$  between  $x = [\sqrt{3}, 0, 1]$  and  $y = [1, 0, 0]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3 + 1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

# Angle between Two Vectors: Example

- What is the angle  $\omega$  between  $x = [5, 0]$  and  $y = [1, 1]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle  $\omega$  between  $x = [\sqrt{3}, 0, 1]$  and  $y = [1, 0, 0]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3+1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

- What is the angle  $\omega$  between  $x = [1, 0, 0, 0, 1]$  and  $y = [0, 1, 1, 0, 0]$ ?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{0}{\sqrt{2} \cdot \sqrt{2}} = \arccos 0 = \frac{\pi}{2}.$$



# Orthogonality

- $x$  and  $y$  are orthogonal if and only if  $\langle x, y \rangle = 0$ .

# Orthogonality

- $x$  and  $y$  are orthogonal if and only if  $\langle x, y \rangle = 0$ .
- Examples in  $\mathbb{R}^n$  with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

# Orthogonality

- $x$  and  $y$  are orthogonal if and only if  $\langle x, y \rangle = 0$ .
- Examples in  $\mathbb{R}^n$  with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

# Orthogonality

- $x$  and  $y$  are orthogonal if and only if  $\langle x, y \rangle = 0$ .
- Examples in  $\mathbb{R}^n$  with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$


$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

$$x = [1, 0], \quad y = [0, 1], \quad (x, y) = 0, \quad \|x\| = \|y\| = 1 \rightarrow \\ x \text{ and } y \text{ are } \textit{orthonormal}.$$

# Orthogonal Projection




- Suppose we have two vectors  $u$  and  $v$ .



# Orthogonal Projection




- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .



# Orthogonal Projection



- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?




# Orthogonal Projection



- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?

If  $0 \leq \omega \leq 90$

$$\begin{aligned}(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\|\end{aligned}$$






# Orthogonal Projection

- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?


If  $90 \leq \omega \leq 180$



# Orthogonal Projection

- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?


If  $90 \leq \omega \leq 180$



# Orthogonal Projection

- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?

If  $90 \leq \omega \leq 180$




# Orthogonal Projection

- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?

If  $90 \leq \omega \leq 180$

$$\begin{aligned} -(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\| \end{aligned}$$




# Orthogonal Projection



- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?

$$|(\mathbf{u}, \mathbf{v})| = \|\mathbf{u}'_v\| \|\mathbf{v}\| \iff \|\mathbf{u}'_v\| = \frac{|(\mathbf{u}, \mathbf{v})|}{\|\mathbf{v}\|}$$




# Orthogonal Projection



- Suppose we have two vectors  $u$  and  $v$ .
- $u'_v$  - *orthogonal projection* of  $u$  on  $v$ .
- What is the length of  $u'_v$ ?


$$|(u, v)| = \|u'_v\| \|v\| \leftrightarrow \|u'_v\| = \frac{|(u, v)|}{\|v\|}$$

$$u'_v = \frac{(u, v)}{(v, v)} v.$$



# Orthogonal Projection: Example

- What's projection of  $u = [1, 3, 2]$  on  $z = [0, 0, 1]$ ?




# Orthogonal Projection: Example

- What's projection of  $u = [1, 3, 2]$  on  $z = [0, 0, 1]$ ?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

*(Projection on the axis = drop other coordinates)*






# Orthogonal Projection: Example

- What's projection of  $u = [1, 3, 2]$  on  $z = [0, 0, 1]$ ?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

*(Projection on the axis = drop other coordinates)*

- What's projection of  $u = [1, 3, 2]$  on  $v = [4, 1, 3]$ ?



# Orthogonal Projection: Example


- What's projection of  $u = [1, 3, 2]$  on  $z = [0, 0, 1]$ ?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

*(Projection on the axis = drop other coordinates)*

- What's projection of  $u = [1, 3, 2]$  on  $v = [4, 1, 3]$ ?

$$u'_v = \frac{(u, v)}{(v, v)} v = \frac{4 + 3 + 6}{16 + 1 + 9} v = \frac{1}{2} v = [2, 0.5, 1.5].$$



# Hyperplanes

- A hyperplane is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one  $w_i \neq 0$ .

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$

# Hyperplanes



# Hyperplanes


- A hyperplane in  $\mathbb{R}^n$  is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one  $w_i \neq 0$ .

# Hyperplanes

A hyperplane in  $\mathbb{R}^2$  is a line



A hyperplane in  $\mathbb{R}^3$  is a plane




Image source: <https://deepai.org/machine-learning-glossary-and-terms/hyperplane>

# Hyperplanes

- A hyperplane in  $\mathbb{R}^n$  is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one  $w_i \neq 0$ .

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$



# Normal to a Hyperplane


- Consider a hyperplane  $(w, x) + b = 0$ .
- Vector  $w = (w_1, \dots, w_n)$  defines the hyperplane.






# Normal to a Hyperplane

- Consider a hyperplane  $(w, x) + b = 0$ .
- Vector  $w = (w_1, \dots, w_n)$  defines the hyperplane.
- $w$  is a *normal vector* to this hyperplane: it's orthogonal to every vector on it.




# ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.




# ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane  $(w, x) + b = 0$ 
  - objects “above” are class +1
  - objects “below” are class -1.




# ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane  $(w, x) + b = 0$ 
  - objects “above” are class +1
  - objects “below” are class -1.
- How can we formalize this?



# ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane  $(w, x) + b = 0$ 
  - objects “above” are class +1
  - objects “below” are class -1.
- How can we formalize this?
  - objects “above”:  $(w, x) + b > 0$
  - objects “below”:  $(w, x) + b < 0$



# Linear Combinations



# Linear Combinations

- We can add vectors together and multiply them by scalars.  
Let's combine these two operations!

# Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- $V$  – a vector space.



# Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.

# Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$  - some scalars.

# Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- $V$  – a vector space.
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$  – some scalars.

$$v = \lambda_1 x_1 + \lambda_2 x_2 + \dots + \lambda_k x_k = \sum_{i=1}^k \lambda_i x_i \in V -$$

a *linear combination* of  $x_1, x_2, \dots, x_k$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

# Linear Combinations: Examples

- In  $(\mathbb{R}^2, +, \cdot)$ , consider vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ .

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  is a linear combination of  $e_1$  and  $e_2$ .

# Linear independence



# Linear Combinations

- A zero vector can always be represented as a trivial linear combination of  $x_1, x_2, \dots, x_k$ :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

# Linear Combinations

- A zero vector can always be represented as a trivial linear combination of  $x_1, x_2, \dots, x_k$ :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

- We are mostly interested in *non-trivial linear combinations* of  $x_1, x_2, \dots, x_k$  where not all  $\lambda_i$  are 0.



# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.

# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

# Linear (In)dependence

- Consider a vector space  $V$ .
- $x_1, x_2, \dots, x_k \in V$  – some vectors.
- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.
- If only trivial solution exists, vectors  $x_1, x_2, \dots, x_k$  are *linearly independent*.

# Linear (In)dependence

- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

# Linear (In)dependence

- If there is a non-trivial linear combination of  $x_1, x_2, \dots, x_k$  such that  $\sum_{i=1}^k \lambda_i x_i = 0$  with at least one  $\lambda_i \neq 0$ , vectors  $x_1, x_2, \dots, x_k$  are *linearly dependent*.

$\Leftrightarrow$

- A set of vectors  $x_1, x_2, \dots, x_k$  is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \dots + \alpha_k x_k$$

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .

# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .
- Vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  are linearly independent:  
there are no  $\lambda_1, \lambda_2 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$



# Linear (In)dependence: Example 1

- Consider  $\mathbb{R}^2$ .
- Vectors  $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  and  $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$  are not linearly independent:  $u = -2v$ .
- Vectors  $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  and  $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  are linearly independent:  
there are no  $\lambda_1, \lambda_2 \in \mathbb{R}$  with at least one  $\lambda_i \neq 0$  such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$
  
(Or: you cannot represent  $e_1$  as  $\lambda e_2$  or vice versa).

# Dimension of a Linear Space

- The linear space  $V$  is called  *$n$ -dimensional* if the maximal number of linearly independent vectors in it is equal to  $n$ .

# Dimension of a Linear Space

- The linear space  $V$  is called  *$n$ -dimensional* if the maximal number of linearly independent vectors in it is equal to  $n$ .

We denote this as  $\dim(V) = n$ .

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

- $e_1, \dots, e_n$  are linearly independent  $\rightarrow \dim(\mathbb{R}^n) \geq n$ .
- Can there be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ ?

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

# Dimension: Example

- $\mathbb{R}^n$  is a  $n$ -dimensional vector space. Why?
- Consider  $n$  vectors  $e_1, \dots, e_n$ :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

- $e_1, \dots, e_n$  are linearly independent  $\rightarrow \dim(\mathbb{R}^n) \geq n$ .

# Dimension: Example

- Can there be more than  $n$  linearly independent vectors in  $\mathbb{R}^n$ ?



# Basis



# Basis

- A set of  $n$  linearly independent vectors  $e_1, e_2, \dots, e_n$  in an  $n$ -dimensional space  $V$  is called a *basis* for  $V$ .

# Basis

- A set of  $n$  linearly independent vectors  $e_1, e_2, \dots, e_n$  in an  $n$ -dimensional space  $V$  is called a *basis* for  $V$ .
- Basis is A set of vectors with which we can represent every vector in the vector space by adding them together and scaling them.

# Basis: Example

# Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^2$ .

# Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^2$ .

- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$  is a basis for  $\mathbb{R}^n$ .

# Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

# Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector  $v$  in a vector space can be represented as a linear combination of  $e_1, e_2, \dots, e_n$ :

$$v = a_1 e_1 + a_2 e_2 + \dots + a_n e_n$$



# Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector  $v$  in a vector space can be represented as a linear combination of  $e_1, e_2, \dots, e_n$ :


$$v = a_1 e_1 + a_2 e_2 + \dots + a_n e_n$$

- $a_1, a_2, \dots, a_n$  - *coordinates* of the vector  $v$  in the basis  $e_1, e_2, \dots, e_n$ .

# Coordinates: Example




- Consider  $\mathbb{R}^2$ .



# Coordinates: Example



- Consider  $\mathbb{R}^2$ .
- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$



# Coordinates: Example




- Consider  $\mathbb{R}^2$ .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Coordinates: Example




- Consider  $\mathbb{R}^2$ .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$x = 4e_1 + 6e_2$$



# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.

# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$



# Orthogonal Basis

- A basis  $e_1, \dots, e_n$  where  $\forall i, j \ e_i \perp e_j$  is an *orthogonal basis*.
- If, additionally,  $\forall i \ \|e_i\| = 1$ , then  $e_1, \dots, e_n$  is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$

# Change of Basis



# Basis

- A vector space has more than one basis.

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;
  - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$  - yet another one.

# Basis

- A vector space has more than one basis.
- Example:  $\mathbb{R}^2$ 
  - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$  - canonical basis;
  - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  - another basis;
  - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$  - yet another one.
- Different basis = different coordinates.  
How exactly do they change?

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

- What are the coordinates in the new basis?

$$x_{new} = ?$$

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .
- New basis:  $e'_1, e'_2, \dots, e'_n$ .

# Coordinate Change: General Case

- Consider a vector space  $V$  with basis  $e_1, e_2, \dots, e_n$ .
- Imagine vector  $x = [x_1, x_2, \dots, x_n] \in V$   
 $x_1, x_2, \dots, x_n$  - coordinates in basis  $e_1, e_2, \dots, e_n$ .
- New basis:  $e'_1, e'_2, \dots, e'_n$ .
- What are the coordinates of  $x$  in this new basis?  
$$x'_1, x'_2, \dots, x'_n = ?$$

# Coordinate Change

- Old basis:  $e_1, e_2, \dots, e_n$   
New basis:  $e'_1, e'_2, \dots, e'_n$
- $x_{old} = [x_1, x_2, \dots, x_n]$ ,  $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:



# Coordinate Change

- Old basis:  $e_1, e_2, \dots, e_n$   
New basis:  $e'_1, e'_2, \dots, e'_n$
- $x_{old} = [x_1, x_2, \dots, x_n]$ ,  $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

$$e'_1 = \alpha_{11}e_1 + \alpha_{21}e_2 + \dots + \alpha_{n1}e_n$$

$$e'_2 = \alpha_{12}e_1 + \alpha_{22}e_2 + \dots + \alpha_{n2}e_n$$

$$\vdots$$

$$e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \dots + \alpha_{ni}e_n$$

$$\vdots$$

$$e'_n = \alpha_{1n}e_1 + \alpha_{2n}e_2 + \dots + \alpha_{nn}e_n$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 \mathbf{e}'_1 + x'_2 \mathbf{e}'_2 + \cdots x'_n \mathbf{e}'_n =$$

# Coordinate Change

$$x = x_1e_1 + x_2e_2 + \cdots + x_ne_n = x'_1\mathbf{e}'_1 + x'_2\mathbf{e}'_2 + \cdots x'_n\mathbf{e}'_n =$$

Remember:  $e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} = x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

$e_1, \dots, e_n$  linearly independent  $\rightarrow$  coefficients in front of them  
should be the same on the both sides of the equality:

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

$e_1, \dots, e_n$  linearly independent  $\rightarrow$  coefficients in front of them should be the same on the both sides of the equality:

$$x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n}$$

$$x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n}$$

$$\vdots$$

$$x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn}$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember:  $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} = x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

$x_{old}$

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{aligned}$$

# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

$$\text{Remember: } e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

$x_{old}$   $x_{new}$



# Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

$$\text{Remember: } e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

The diagram illustrates the coordinate change process. It shows a vector  $x_{old}$  (represented by a red box) being expressed as a linear combination of new basis vectors  $e'_i$  (represented by a blue box). The coefficients are grouped into a matrix  $x_{new}$  (represented by a green box). The matrix  $x_{new}$  is shown as a collection of columns, each corresponding to a new basis vector  $e'_i$ . The columns are labeled  $x'_1, x'_i, x'_n$  and the rows are labeled  $\alpha_{11}, \alpha_{21}, \dots, \alpha_{ni}, \alpha_{nn}$ . The matrix  $x_{new}$  is shown as a collection of columns, each corresponding to a new basis vector  $e'_i$ . The columns are labeled  $x'_1, x'_i, x'_n$  and the rows are labeled  $\alpha_{11}, \alpha_{21}, \dots, \alpha_{ni}, \alpha_{nn}$ .

$$\begin{matrix} x_{old} \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} \end{matrix} = \begin{matrix} x'_1 & x'_i & x'_n \\ \alpha_{11} & \alpha_{1i} & \alpha_{1n} \\ \alpha_{21} & \alpha_{2i} & \alpha_{2n} \\ \vdots & \vdots & \vdots \\ \alpha_{n1} & \alpha_{ni} & \alpha_{nn} \end{matrix} \begin{matrix} + \cdots + \\ + \cdots + \\ + \cdots + \end{matrix}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ x_{old} & \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} & = & \begin{bmatrix} x'_1 \\ x'_2 \\ \vdots \\ x'_n \end{bmatrix} & \begin{bmatrix} \alpha_{11} & \dots & \alpha_{1i} & \dots & \alpha_{1n} \\ \alpha_{21} & \dots & \alpha_{2i} & \dots & \alpha_{2n} \\ \vdots & & \vdots & & \vdots \\ \alpha_{n1} & \dots & \alpha_{ni} & \dots & \alpha_{nn} \end{bmatrix} \end{matrix}$$

$x_{new}$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} & = & \begin{matrix} x'_1 \\ x'_1 \\ \vdots \\ x'_1 \end{matrix} \begin{matrix} \alpha_{11} \\ \alpha_{21} \\ \vdots \\ \alpha_{n1} \end{matrix} + \cdots + \begin{matrix} x'_i \\ x'_i \\ \vdots \\ x'_i \end{matrix} \begin{matrix} \alpha_{1i} \\ \alpha_{2i} \\ \vdots \\ \alpha_{ni} \end{matrix} + \cdots + \begin{matrix} x'_n \\ x'_n \\ \vdots \\ x'_n \end{matrix} \begin{matrix} \alpha_{1n} \\ \alpha_{2n} \\ \vdots \\ \alpha_{nn} \end{matrix} \end{matrix}$$

$x_{old}$   $x_{new}$

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} & = & \begin{matrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix} \end{matrix}$$

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned} \Leftrightarrow \begin{aligned} x'_1 &= 3 \\ x'_2 &= 4 \end{aligned}$$

# Coordinate Change: Example

- Consider  $\mathbb{R}^2$  with basis  $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis:  $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ ,  $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} x_{old} \\ x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix}$$

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned} \Leftrightarrow \begin{aligned} x'_1 &= 3 \\ x'_2 &= 4 \end{aligned} \Leftrightarrow x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

# Coordinate Change

- Going from one basis to the other:

The diagram shows the transformation of a vector  $x_{old}$  from an old basis to a new basis  $x_{new}$ . The vector  $x_{old}$  is represented by a red box containing the components  $x_1, x_2, \dots, x_n$ . The new basis vectors  $e'_i$  are shown at the top, with lines connecting them to the corresponding terms in the expansion of  $x_{old}$ . The expansion is shown as a sum of products of new basis components  $x'_i$  and old basis components  $\alpha_{ij}$ . The new basis components  $x'_i$  are highlighted in green boxes, and the old basis components  $\alpha_{ij}$  are highlighted in blue boxes. The new basis components  $x'_i$  are shown in a green box, and the old basis components  $\alpha_{ij}$  are shown in a blue box. The new basis components  $x'_i$  are shown in a green box, and the old basis components  $\alpha_{ij}$  are shown in a blue box. The new basis components  $x'_i$  are shown in a green box, and the old basis components  $\alpha_{ij}$  are shown in a blue box.

$$\begin{matrix} x_{old} \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} \end{matrix} = \begin{matrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix}$$

$x_{new}$

# Coordinate Change

- Going from one basis to the other:

The diagram illustrates the coordinate change from an old basis to a new basis. It shows a vector  $x_{old}$  (represented by a red box) being expressed as a linear combination of new basis vectors  $e'_i$  (represented by blue boxes). The coefficients of this linear combination are the components of the new vector  $x_{new}$  (represented by a green box). The equations are:

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{aligned}$$

Blue boxes highlight the terms  $x'_i \alpha_{1i}$ ,  $x'_i \alpha_{2i}$ , and  $x'_i \alpha_{ni}$  in each equation, which are connected by a green line to the  $x'_i$  term in the  $x_{new}$  label below. A blue line connects the  $e'_i$  label to the corresponding  $\alpha_{ji}$  terms in the equations.

- There is a more compact way of writing this down using [matrices](#).



# Matrices



# A Matrix

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

# A Matrix

- $A \in \mathbb{R}^{m \times n}$  - a matrix with  $m$  rows and  $n$  columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- *Examples:*

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$   $(a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j)$

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$   $(a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j)$
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$   $(a_{ii} = 1, a_{ij} = 0 \forall i \neq j)$

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )
- Symmetric matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ij} = a_{ji}$ )

# Special Matrices

- Diagonal matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$ )
- Identity matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  ( $a_{ii} = 1, a_{ij} = 0 \forall i \neq j$ )
- Symmetric matrix:  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$  ( $a_{ij} = a_{ji}$ )
- Triangular matrix:  $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$  ( $a_{ij} = 0 \forall i > j \text{ or } \forall i < j$ )

# Vectors vs Matrices

- An  $n$ -dimensional vector can be considered a  $n \times 1$  matrix:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n \times 1}$$



# **Operations with Matrices**



# Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

# Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Transpose = writing columns as rows:

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \cdots & \cdots & \cdots & \cdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}, \quad \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}^T = [x_1, \dots, x_n]$$

# Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$

# Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$
- Transposing a symmetrical matrix = no changes:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}$$

# Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

# Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

- Example:

$$5 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

# Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$



# Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 0 & 4 \end{bmatrix}$$

# Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$  - a vector space.  
“Vectors” = matrices.

# Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$  - a vector space.  
“Vectors” = matrices.
- You can check yourself that the necessary axioms hold.

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.
- Each element  $c_{ij}$  of  $C$  is a dot product of  $i$ -th row of  $A$  and  $j$ -th column of  $B$ :  $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$

# Matrix Multiplication

- Consider two matrices  $A = \{a_{ij}\}_{m \times n}$  and  $b = \{b_{ij}\}_{n \times p}$ .
- $C = AB$  – product of two matrices.
- Each element  $c_{ij}$  of  $C$  is a dot product of  $i$ -th row of  $A$  and  $j$ -th column of  $B$ :  $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$
- Example  $\mathbb{R}^{2 \times 2}$ : 
$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$



# Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

$$= \begin{bmatrix} 16 & 7 & 26 \\ 43 & 22 & 61 \end{bmatrix}$$

# To sum up

- Vectors
  - Vector spaces
  - Inner products
  - Lengths
  - Distances
  - Angles
- Analytic Geometry
  - Projections
  - Hyperplanes
  - Normal vector
- Vector spaces
  - Linear (in)dependence
  - Basis
- Matrices
  - Matrix operations

# Next Time

- Calculus recap
- Probability theory recap