


Intro to Math and Python

Lecture 1



girafe
ai



Today

Abstract geometric shapes consisting of several rounded, irregular polygons outlined in black, located in the bottom-left corner of the slide.

1. Course overview
2. Linear Algebra
 - Core objects
 - Vector spaces
3. A bit of Analytic Geometry
 - Orthogonal projections
 - Hyperplanes
 - Normals

About this course




Image source: Mathematics for Machine Learning, p. 14
(<https://mml-book.github.io/book/mml-book.pdf>)

About this course



This week:

1. Linear algebra
2. Calculus
3. Probability theory

Prerequisites:

- basic knowledge of math;
- some Python.

About me

Iurii Efimov

✉ yuri.efimov@phystech.edu

in [efimov-iurii](https://www.linkedin.com/in/efimov-iurii)

◦ PhD researcher



◦ Lecturer



◦ Artec3D DS Team



Linear Algebra: the Basics



Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a vector with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a vector with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

Linear Algebra: Core Objects

- $\alpha \in \mathbb{R}$ - a scalar *Example: -2*
- $x \in \mathbb{R}^n$ - a **vector** with n entries

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3, \quad \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix} \in \mathbb{R}^5$$

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns


$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \quad \text{Example: } \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

What are Vectors?

- Ordered sets of numbers: $x = [1, 2]$

What are Vectors?


- Ordered sets of numbers: $x = [1, 2]$
- A point with Cartesian coordinates




What are Vectors?

- Ordered sets of numbers: $x = [1, 2]$

- A point with Cartesian coordinates



- Direction + length



Vector Spaces



Vector Space: Definition

- A real-valued **vector space** $(V, +, \cdot)$ is a set of vectors V with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

Vector Space: Definition

- A real-valued vector space $(V, +, \cdot)$ is a set of vectors V with two operations

$$(1) +: V \times V \rightarrow V, \quad (2) \cdot: \mathbb{R} \times V \rightarrow V$$

that satisfy the following properties (axioms):

	Property	Meaning
1.	Associativity of addition	$x + (y + z) = (x + y) + z$
2.	Commutativity of addition	$x + y = y + x$
3.	Identity element of addition	$\exists 0 \in V: \forall x \in V \quad 0 + x = x$
4.	Identity element of scalar multiplication	$\forall x \in V \quad 1 \cdot x = x$
5.	Inverse element of addition	$\forall x \in V \exists -x \in V: x + (-x) = 0$
6.	Compatibility of scalar multiplication	$\alpha(\beta x) = (\alpha\beta)x$
7.	Distributivity	$(\alpha + \beta)x = \alpha x + \beta x$
8.		$\alpha(x + y) = \alpha x + \alpha y$

Let's define vector operations!

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \in \mathbb{R}^n, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

Operations with Vectors: Example

$x, y \in \mathbb{R}^3$:

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Operations with Vectors: Example

$x, y \in \mathbb{R}^3$:

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

Operations with Vectors: Example

$x, y \in \mathbb{R}^3$:

$$x = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad y = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Multiplying by a scalar:

$$10x = \begin{bmatrix} 10 \\ 0 \\ 10 \end{bmatrix}$$

Sum:


$$x + y = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$

Vector Operations: Geometrical Interpretation

Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 3]$$

$$2\vec{a} = [4, 6]$$

$$-0.5\vec{a} = [-1, -1.5]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$




Vectors: Geometrical Interpretation



$$\vec{a} = [2, 5]$$

$$\vec{b} = [4, 1]$$

$$\vec{a} + \vec{b} = [6, 6]$$



Back to Vector Spaces

Operations with Vectors

1. Sum of two vectors:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, \quad x + y = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_2 \end{bmatrix} \in \mathbb{R}^n$$

2. Multiplying by a scalar:

satisfy axioms (1) – (8)
(check it yourself)

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \quad \alpha \in \mathbb{R}, \quad \alpha x = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix} \in \mathbb{R}^n$$

Vector Spaces

$(\mathbb{R}^n, +, \cdot), n \in \mathbb{N}$ - a vector space with operations

1. vector addition:

$$x + y = (x_1, x_2, \dots, x_n) + (y_1, y_2, \dots, y_n) = (x_1 + y_1, \dots, x_n + y_n)$$

2. multiplication by a scalar:

$$\alpha x = \alpha(x_1, x_2, \dots, x_n) = (\alpha x_1, \alpha x_2, \dots, \alpha x_n)$$

Inner Product



Inner Product

- Inner product is a function $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{R}$ that satisfies the following properties:
 - *Symmetric*: $\forall x, y \in V \quad \langle x, y \rangle = \langle y, x \rangle$
 - *Positive definite*: $\forall x \in V \setminus \{0\} \quad \langle x, x \rangle > 0$ and $\langle x, 0 \rangle = 0$.

Dot Product

- A particular type of inner product.

Dot Product

- A particular type of inner product.
- For $x = [x_1, \dots, x_n]$, $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

Dot Product

- A particular type of inner product.

- For $x = [x_1, \dots, x_n]$, $y = [y_1, \dots, y_n] \in \mathbb{R}^n$

$$(x, y) = x_1y_1 + x_2y_2 + \dots + x_ny_n$$

- Example:

$$x = [1, 2, 3, 4], \quad y = [-1, 0, 1, 2]$$

$$(x, y) = 1 \cdot (-1) + 2 \cdot 0 + 3 \cdot 1 + 4 \cdot 2 = -1 + 0 + 3 + 8 = 10$$

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$ with dot product (\cdot, \cdot) is called a *Euclidian vector space*.

Euclidian Vector Space

- A vector space with inner product is called an *inner product space*.
- $(\mathbb{R}^n, +, \cdot)$ with dot product (\cdot, \cdot) is called a *Euclidian vector space*.
- *Note: there're inner products different from dot product.*

Norms



Norm

- A **norm** on a vector space V is a function $\| \cdot \|: V \rightarrow \mathbb{R}$ which assigns each vector $x \in V$ its *length* $\|x\| \in \mathbb{R}$.

Norm

- A **norm** on a vector space \mathbb{V} is a function $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:

Norm


- A **norm** on a vector space \mathbb{V} is a function $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$

Norm

- A **norm** on a vector space \mathbb{V} is a function $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
 - *Positive definite*: $\forall x \in \mathbb{V} \quad \|x\| \geq 0$ and $\|x\| = 0 \Leftrightarrow x = 0$

Norm

- A **norm** on a vector space \mathbb{V} is a function $\| \cdot \|: \mathbb{V} \rightarrow \mathbb{R}$ which assigns each vector $x \in \mathbb{V}$ its *length* $\|x\| \in \mathbb{R}$.
- A norm must satisfy the following properties:
 - *Absolutely homogeneous*: $\forall x \in \mathbb{V}, \forall \lambda \in \mathbb{R} \quad \|\lambda x\| = |\lambda| \|x\|$
 - *Positive definite*: $\forall x \in \mathbb{V} \quad \|x\| \geq 0$ and $\|x\| = 0 \Leftrightarrow x = 0$
 - *Triangle inequality*: $\forall x, y \in \mathbb{V} \quad \|x + y\| \leq \|x\| + \|y\|$



Examples of Norms

Manhattan Norm



- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$



Manhattan Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 6$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$



Manhattan Norm

- A norm for $x \in \mathbb{R}^n$:


$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

- Examples:

$$\|[1, 2, 3]\|_1 = 1 + 2 + 3 = 5$$

$$\|[1, 0]\|_1 = 1 + 0 = 1$$

$$\|[-1, 0]\|_1 = |-1| + 0 = 1$$





Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$



Euclidian Norm

- A norm for $x \in \mathbb{R}^n$:


$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$$

- Computes *Euclidian distance* from the origin.
- Examples:

$$\|[1, 2, 3]\|_2 = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}$$

$$\|[1, 0]\|_2 = \sqrt{1^2 + 0} = 1$$

$$\|[-1, 0]\|_2 = \sqrt{(-1)^2 + 0} = 1$$



Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- ℓ_1 - Manhattan norm $\|\cdot\|_1$;
- ℓ_2 - Euclidian norm $\|\cdot\|$ (default);

Other norms

- In general, for $x = [x_1, \dots, x_n] \in \mathbb{R}^n$ an ℓ_p -norm is defined as follows:

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}.$$

- ℓ_1 - Manhattan norm $\|\cdot\|_1$;
- ℓ_2 - Euclidian norm $\|\cdot\|$ (default);
- ℓ_∞ : $\|x\|_\infty = \max_i |x_i|$

Example: $\|[1, 2, 3]\|_\infty = 3$, $\|[1, 0]\|_\infty = 1$, $\|[-1, 0.5]\|_\infty = 1$.

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \|x\|_2$$

Inner Product and Norms

- Any inner product induces a norm:

$$\|x\| := \sqrt{\langle x, x \rangle}$$

Check yourself that this is indeed a norm.

- Example: dot product induces Euclidian norm:

$$\sqrt{(x, x)} = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} = \|x\|_2$$

- (!) Not every norm is induced by an inner product.
Example: Manhattan norm.

Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

Cauchy-Schwarz Inequality

- For an inner product vector space, the induced norm satisfies the inequality:

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

- For dot product and Euclidian norm:

$$|(x, y)| \leq \|x\|_2 \cdot \|y\|_2$$

Distance between Vectors

- Distance between two vectors x and y is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

Distance between Vectors

- Distance between two vectors x and y is defined as

$$d(x, y) := \|x - y\| = \sqrt{\langle x - y, x - y \rangle}.$$

- For dot product and Euclidian norm, we get *Euclidian distance*:

$$\begin{aligned} d(x, y) &= \|x - y\|_2 = \sqrt{(x - y, x - y)} = \\ &= \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \dots + (x_n - y_n)^2}. \end{aligned}$$

Angles and Orthogonality




Angle between Two Vectors

- Inner product also captures the geometry of vector space by defining the angle between two vectors.
- Remember Cauchy-Schwarz inequality:

$$|(x, y)| \leq \|x\| \cdot \|y\|$$

$$-1 \leq \frac{(x, y)}{\|x\| \cdot \|y\|} \leq 1$$

$$\omega: \cos \omega = \frac{(x, y)}{\|x\| \cdot \|y\|} \text{ - angle between } x \text{ and } y.$$



Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle ω between $x = [\sqrt{3}, 0, 1]$ and $y = [1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3 + 1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

Angle between Two Vectors: Example

- What is the angle ω between $x = [5, 0]$ and $y = [1, 1]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{5 \cdot 1 + 0 \cdot 1}{\sqrt{5^2 + 0^2} \cdot \sqrt{1^2 + 1^2}} = \arccos \frac{5}{5\sqrt{2}} = \arccos \frac{\sqrt{2}}{4} = \frac{\pi}{4}.$$

- What is the angle ω between $x = [\sqrt{3}, 0, 1]$ and $y = [1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{\sqrt{3}}{\sqrt{3+1} \cdot \sqrt{1}} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}.$$

- What is the angle ω between $x = [1, 0, 0, 0, 1]$ and $y = [0, 1, 1, 0, 0]$?

$$\omega = \arccos \frac{(x, y)}{\|x\| \|y\|} = \arccos \frac{0}{\sqrt{2} \cdot \sqrt{2}} = \arccos 0 = \frac{\pi}{2}.$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$

$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

Orthogonality

- x and y are orthogonal if and only if $\langle x, y \rangle = 0$.
- Examples in \mathbb{R}^n with dot product:

$$x = [2, 3], \quad y = [1, 0], \quad (x, y) = 2 \neq 0 \rightarrow \\ x \text{ and } y \text{ are not orthogonal.}$$


$$x = [1, 2, 3], \quad y = [-2, 1, 0], \quad (x, y) = -2 + 2 + 0 = 0 \rightarrow \\ x \text{ and } y \text{ are orthogonal.}$$

$$x = [1, 0], \quad y = [0, 1], \quad (x, y) = 0, \quad \|x\| = \|y\| = 1 \rightarrow \\ x \text{ and } y \text{ are } \textit{orthonormal}.$$

Orthogonal Projection




- Suppose we have two vectors u and v .



Orthogonal Projection




- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .



Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $0 \leq \omega \leq 90$


$$\begin{aligned}(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\|\end{aligned}$$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


If $90 \leq \omega \leq 180$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


If $90 \leq \omega \leq 180$



Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $90 \leq \omega \leq 180$




Orthogonal Projection

- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

If $90 \leq \omega \leq 180$

$$\begin{aligned} -(u, v) &= \|u\| \|v\| \cos \omega = \|u\| \|v\| \frac{\|u'_v\|}{\|u\|} = \\ &= \|u'_v\| \|v\| \end{aligned}$$




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?

$$|(\mathbf{u}, \mathbf{v})| = \|\mathbf{u}'_v\| \|\mathbf{v}\| \iff \|\mathbf{u}'_v\| = \frac{|(\mathbf{u}, \mathbf{v})|}{\|\mathbf{v}\|}$$




Orthogonal Projection



- Suppose we have two vectors u and v .
- u'_v - *orthogonal projection* of u on v .
- What is the length of u'_v ?


$$|(u, v)| = \|u'_v\| \|v\| \leftrightarrow \|u'_v\| = \frac{|(u, v)|}{\|v\|}$$

$$u'_v = \frac{(u, v)}{(v, v)} v.$$



Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?




Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

(Projection on the axis = drop other coordinates)




Orthogonal Projection: Example

- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

(Projection on the axis = drop other coordinates)

- What's projection of $u = [1, 3, 2]$ on $v = [4, 1, 3]$?



Orthogonal Projection: Example


- What's projection of $u = [1, 3, 2]$ on $z = [0, 0, 1]$?

$$u'_z = \frac{(u, z)}{(z, z)} z = 3z = [0, 0, 3]$$

(Projection on the axis = drop other coordinates)

- What's projection of $u = [1, 3, 2]$ on $v = [4, 1, 3]$?

$$u'_v = \frac{(u, v)}{(v, v)} v = \frac{4 + 3 + 6}{16 + 1 + 9} v = \frac{1}{2} v = [2, 0.5, 1.5].$$



Hyperplanes

- A hyperplane is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$

Hyperplanes



Hyperplanes


- A hyperplane in \mathbb{R}^n is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

Hyperplanes

A hyperplane in \mathbb{R}^2 is a line



A hyperplane in \mathbb{R}^3 is a plane




Image source: <https://deepai.org/machine-learning-glossary-and-terms/hyperplane>

Hyperplanes

- A hyperplane in \mathbb{R}^n is described by equation

$$w_1x_1 + w_2x_2 + \cdots + w_nx_n + b = 0$$

where at least one $w_i \neq 0$.

- A more compact notation:

$$(w, x) + b = 0, \quad w = (w_1, \dots, w_n)$$




Normal to a Hyperplane

- Consider a hyperplane $(w, x) + b = 0$.
- Vector $w = (w_1, \dots, w_n)$ defines the hyperplane.




Normal to a Hyperplane

- Consider a hyperplane $(w, x) + b = 0$.
- Vector $w = (w_1, \dots, w_n)$ defines the hyperplane.
- w is a *normal vector* to this hyperplane: it's orthogonal to every vector on it.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.




ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.
- How can we formalize this?



ML Example: Linear Classifier

- Objects = 2D vectors
- Binary classification: classes +1 and -1.
- Linear classifier: separating hyperplane $(w, x) + b = 0$
 - objects “above” are class +1
 - objects “below” are class -1.
- How can we formalize this?
 - objects “above”: $(w, x) + b > 0$
 - objects “below”: $(w, x) + b < 0$



Linear Combinations



Linear Combinations

- We can add vectors together and multiply them by scalars.
Let's combine these two operations!

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ - some scalars.

Linear Combinations

- We can add vectors together and multiply them by scalars. Let's combine these two operations!
- V – a vector space.
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- $\lambda_1, \lambda_2, \dots, \lambda_k \in \mathbb{R}$ – some scalars.

$$v = \lambda_1 x_1 + \lambda_2 x_2 + \dots + \lambda_k x_k = \sum_{i=1}^k \lambda_i x_i \in V -$$

a *linear combination* of x_1, x_2, \dots, x_k .

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Linear Combinations: Examples

- In $(\mathbb{R}^2, +, \cdot)$, consider vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

$u = 2e_1 - e_2 = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ is a linear combination of e_1 and e_2 .

Linear independence



Linear Combinations

- A zero vector can always be represented as a trivial linear combination of x_1, x_2, \dots, x_k :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

Linear Combinations

- A zero vector can always be represented as a trivial linear combination of x_1, x_2, \dots, x_k :

$$0 = \sum_{i=1}^k 0 \cdot x_i$$

- We are mostly interested in *non-trivial linear combinations* of x_1, x_2, \dots, x_k where not all λ_i are 0.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

Linear (In)dependence

- Consider a vector space V .
- $x_1, x_2, \dots, x_k \in V$ – some vectors.
- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.
- If only trivial solution exists, vectors x_1, x_2, \dots, x_k are *linearly independent*.

Linear (In)dependence

- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

Linear (In)dependence

- If there is a non-trivial linear combination of x_1, x_2, \dots, x_k such that $\sum_{i=1}^k \lambda_i x_i = 0$ with at least one $\lambda_i \neq 0$, vectors x_1, x_2, \dots, x_k are *linearly dependent*.

\Leftrightarrow

- A set of vectors x_1, x_2, \dots, x_k is linearly dependent if and only if (at least) one of the vectors is a linear combination of the others

$$x_i = \alpha_1 x_1 + \dots + \alpha_k x_k$$

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.
- Vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are linearly independent:
there are no $\lambda_1, \lambda_2 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Linear (In)dependence: Example 1

- Consider \mathbb{R}^2 .
- Vectors $u = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$ and $v = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$ are not linearly independent: $u = -2v$.
- Vectors $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ are linearly independent:
there are no $\lambda_1, \lambda_2 \in \mathbb{R}$ with at least one $\lambda_i \neq 0$ such that
$$\lambda_1 e_1 + \lambda_2 e_2 = \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

(Or: you cannot represent e_1 as λe_2 or vice versa).

Basis



Basis

- A set of n linearly independent vectors e_1, e_2, \dots, e_n in an n -dimensional space V is called a *basis* for V .

Basis

- A set of n linearly independent vectors e_1, e_2, \dots, e_n in an n -dimensional space V is called a *basis* for V .
- Basis is A set of vectors with which we can represent every vector in the vector space by adding them together and scaling them.

Basis: Example

Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .

Basis: Example

- $e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .

- $e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \dots, e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^n .

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector v in a vector space can be represented as a linear combination of e_1, e_2, \dots, e_n :

$$v = a_1 e_1 + a_2 e_2 + \dots + a_n e_n$$

Coordinates

- Let's fix the order of the vectors in the basis:

$$e_1, e_2, \dots, e_n$$

- Every vector v in a vector space can be represented as a linear combination of e_1, e_2, \dots, e_n :


$$v = a_1 e_1 + a_2 e_2 + \dots + a_n e_n$$

- a_1, a_2, \dots, a_n - *coordinates* of the vector v in the basis e_1, e_2, \dots, e_n .

Coordinates: Example




- Consider \mathbb{R}^2 .



Coordinates: Example



- Consider \mathbb{R}^2 .
- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$



Coordinates: Example




- Consider \mathbb{R}^2 .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Coordinates: Example




- Consider \mathbb{R}^2 .

- $x = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$

- Canonical basis:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$x = 4e_1 + 6e_2$$



Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$

Orthogonal Basis

- A basis e_1, \dots, e_n where $\forall i, j \ e_i \perp e_j$ is an *orthogonal basis*.
- If, additionally, $\forall i \ \|e_i\| = 1$, then e_1, \dots, e_n is an *orthonormal basis*.

Example:

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \ e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \ \dots, \ e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \text{ is an orthonormal basis of } \mathbb{R}^n.$$

Change of Basis



Basis

- A vector space has more than one basis.

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;
 - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$ - yet another one.

Basis

- A vector space has more than one basis.
- Example: \mathbb{R}^2
 - $e = \left\{ e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$ - canonical basis;
 - $a = \left\{ a_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, a_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ - another basis;
 - $b = \left\{ b_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \right\}$ - yet another one.
- Different basis = different coordinates.
How exactly do they change?

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

Coordinate Change: Example

- Consider \mathbb{R}^2 with canonical basis

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- New basis:

$$e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- $x_{old} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

- What are the coordinates in the new basis?

$$x_{new} = ?$$

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .
- New basis: e'_1, e'_2, \dots, e'_n .

Coordinate Change: General Case

- Consider a vector space V with basis e_1, e_2, \dots, e_n .
- Imagine vector $x = [x_1, x_2, \dots, x_n] \in V$
 x_1, x_2, \dots, x_n - coordinates in basis e_1, e_2, \dots, e_n .
- New basis: e'_1, e'_2, \dots, e'_n .
- What are the coordinates of x in this new basis?
$$x'_1, x'_2, \dots, x'_n = ?$$

Coordinate Change

- Old basis: e_1, e_2, \dots, e_n
New basis: e'_1, e'_2, \dots, e'_n
- $x_{old} = [x_1, x_2, \dots, x_n]$, $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

Coordinate Change

- Old basis: e_1, e_2, \dots, e_n
New basis: e'_1, e'_2, \dots, e'_n
- $x_{old} = [x_1, x_2, \dots, x_n]$, $x_{new} = [x'_1, x'_2, \dots, x'_n] = ?$
- Coordinates of the new basis in the old one:

$$e'_1 = \alpha_{11}e_1 + \alpha_{21}e_2 + \dots + \alpha_{n1}e_n$$

$$e'_2 = \alpha_{12}e_1 + \alpha_{22}e_2 + \dots + \alpha_{n2}e_n$$

$$\vdots$$

$$e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \dots + \alpha_{ni}e_n$$

$$\vdots$$

$$e'_n = \alpha_{1n}e_1 + \alpha_{2n}e_2 + \dots + \alpha_{nn}e_n$$

Coordinate Change

$$x = x_1e_1 + x_2e_2 + \cdots + x_ne_n = x'_1\mathbf{e}'_1 + x'_2\mathbf{e}'_2 + \cdots x'_n\mathbf{e}'_n =$$

Coordinate Change

$$x = x_1e_1 + x_2e_2 + \cdots + x_ne_n = x'_1\mathbf{e}'_1 + x'_2\mathbf{e}'_2 + \cdots x'_n\mathbf{e}'_n =$$

Remember: $e'_i = \alpha_{1i}e_1 + \alpha_{2i}e_2 + \cdots + \alpha_{ni}e_n$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} = x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

e_1, \dots, e_n linearly independent \rightarrow coefficients in front of them
should be the same on the both sides of the equality:

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} &= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ &\quad + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

e_1, \dots, e_n linearly independent \rightarrow coefficients in front of them should be the same on the both sides of the equality:

$$x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n}$$

$$x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n}$$

$$\vdots$$

$$x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$\begin{aligned} = x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) = \end{aligned}$$

x_{old}

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{aligned}$$

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

Remember: $e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

x_{old}

x_{new}

Coordinate Change

$$x = x_1 e_1 + x_2 e_2 + \cdots + x_n e_n = x'_1 e'_1 + x'_2 e'_2 + \cdots + x'_n e'_n =$$

$$\text{Remember: } e'_i = \alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n$$

$$= x'_1 \cdot (\alpha_{11} e_1 + \alpha_{21} e_2 + \cdots + \alpha_{n1} e_n) + \cdots + x'_i \cdot (\alpha_{1i} e_1 + \alpha_{2i} e_2 + \cdots + \alpha_{ni} e_n) + \cdots \\ + x'_n (\alpha_{1n} e_1 + \alpha_{2n} e_2 + \cdots + \alpha_{nn} e_n) =$$

The diagram illustrates the coordinate change process. A vector x_{old} (represented by a red box) is shown as a linear combination of new basis vectors e'_i (represented by a blue box). The coefficients of this combination are grouped into a matrix x_{new} (represented by a green box). The matrix x_{new} is a $n \times n$ matrix where each row corresponds to a new basis vector e'_i and each column corresponds to an old basis vector e_j . The entries are the coefficients α_{ji} .

$$\begin{matrix} & & & e'_i & & \\ & & & \swarrow & \searrow & \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} & = & \begin{matrix} x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{matrix} \end{matrix}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ & & & & \swarrow \quad \downarrow \quad \searrow \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} = & \begin{matrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix} \end{matrix}$$

x_{old} x_{new}

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{array}{c}
 \mathbf{x}_{old} \\
 \begin{array}{l}
 x_1 = x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\
 x_2 = x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\
 \vdots \\
 x_n = x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn}
 \end{array}
 \end{array}$$

e'_i

\mathbf{x}_{new}

$$\begin{aligned}
 2 &= 2x'_1 - 1x'_2 \\
 -1 &= 1x'_1 - 1x'_2
 \end{aligned}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} & = & \begin{matrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix} \end{matrix}$$

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned} \iff \begin{aligned} x'_1 &= 3 \\ x'_2 &= 4 \end{aligned}$$

Coordinate Change: Example

- Consider \mathbb{R}^2 with basis $e_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $e_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- New basis: $e'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $e'_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$
- $x_{old} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$, $x_{new} = \begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix}$

$$\begin{matrix} & & & & e'_i \\ \begin{matrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{matrix} & = & \begin{matrix} x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ \vdots \\ x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{matrix} \end{matrix}$$

$$\begin{aligned} 2 &= 2x'_1 - 1x'_2 \\ -1 &= 1x'_1 - 1x'_2 \end{aligned} \Leftrightarrow \begin{aligned} x'_1 &= 3 \\ x'_2 &= 4 \end{aligned} \Leftrightarrow x_{new} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

Coordinate Change

- Going from one basis to the other:

The diagram shows the transformation of a vector x_{old} from an old basis to a new basis x_{new} . The vector x_{old} is represented by a red box containing the components x_1, x_2, \dots, x_n . The new basis vectors e'_i are shown at the top, with lines connecting them to the coefficients α_{ij} in the equations. The coefficients α_{ij} are grouped into green boxes, which are labeled x_{new} at the bottom. The equations are:

$$\begin{aligned} x_1 &= x'_1 \alpha_{11} + \dots + x'_i \alpha_{1i} + \dots + x'_n \alpha_{1n} \\ x_2 &= x'_1 \alpha_{21} + \dots + x'_i \alpha_{2i} + \dots + x'_n \alpha_{2n} \\ &\vdots \\ x_n &= x'_1 \alpha_{n1} + \dots + x'_i \alpha_{ni} + \dots + x'_n \alpha_{nn} \end{aligned}$$

Coordinate Change

- Going from one basis to the other:

$$\begin{array}{l} x_1 = x'_1 \alpha_{11} + \cdots + x'_i \alpha_{1i} + \cdots + x'_n \alpha_{1n} \\ x_2 = x'_1 \alpha_{21} + \cdots + x'_i \alpha_{2i} + \cdots + x'_n \alpha_{2n} \\ \vdots \\ x_n = x'_1 \alpha_{n1} + \cdots + x'_i \alpha_{ni} + \cdots + x'_n \alpha_{nn} \end{array}$$

x_{old}

x_{new}

e'_i

- There is a more compact way of writing this down using [matrices](#).

Matrices



A Matrix

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

A Matrix

- $A \in \mathbb{R}^{m \times n}$ - a matrix with m rows and n columns:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- *Examples:*

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \mathbb{R}^{2 \times 2}, \quad \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \in \mathbb{R}^{2 \times 3}$$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ $(a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j)$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ $(a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j)$
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ $(a_{ii} = 1, a_{ij} = 0 \forall i \neq j)$

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ($a_{ii} = 1, a_{ij} = 0 \forall i \neq j$)
- Symmetric matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ij} = a_{ji}$)

Special Matrices

- Diagonal matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ii} \neq 0, a_{ij} = 0 \forall i \neq j$)
- Identity matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ ($a_{ii} = 1, a_{ij} = 0 \forall i \neq j$)
- Symmetric matrix: $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$ ($a_{ij} = a_{ji}$)
- Triangular matrix: $\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$ ($a_{ij} = 0 \forall i > j \text{ or } \forall i < j$)

Vectors vs Matrices

- An n -dimensional vector can be considered a $n \times 1$ matrix:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^{n \times 1}$$

Operations with Matrices



Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

Transpose of a Matrix

- Consider a matrix

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- Transpose = writing columns as rows:

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \cdots & \cdots & \cdots & \cdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}, \quad \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}^T = [x_1, \dots, x_n]$$

Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$

Transpose of a Matrix: Example

- $\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 0 & 2 \\ 1 & 1 \\ 2 & 5 \end{bmatrix}$
- Transposing a symmetrical matrix = no changes:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}^T = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 1 & 2 \\ 3 & 2 & 1 \end{bmatrix}$$

Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

Multiplying by a Scalar

- We can multiply matrix by a scalar:

$$\lambda A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \lambda a_{11} & \lambda a_{12} & \cdots & \lambda a_{1n} \\ \lambda a_{21} & \lambda a_{22} & \cdots & \lambda a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \lambda a_{m1} & \lambda a_{m2} & \cdots & \lambda a_{mn} \end{bmatrix}$$

- Example:

$$5 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

Sum of Two Matrices

- We can sum up matrices of the same size:

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}, \quad B = \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ b_{m1} & b_{m2} & \cdots & b_{mn} \end{bmatrix},$$

$$A + B = \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \cdots & a_{1n} + b_{1n} \\ a_{21} + b_{21} & a_{22} + b_{22} & \cdots & a_{2n} + b_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

- Example:

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} -1 & 2 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 2 \\ 0 & 4 \end{bmatrix}$$

Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$ - a vector space.
“Vectors” = matrices.

Matrices Also Form a Vector Space!

- $(\mathbb{R}^{m \times n}, +, \cdot)$ - a vector space.
“Vectors” = matrices.
- You can check yourself that the necessary axioms hold.

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.
- Each element c_{ij} of C is a dot product of i -th row of A and j -th column of B : $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$

Matrix Multiplication

- Consider two matrices $A = \{a_{ij}\}_{m \times n}$ and $b = \{b_{ij}\}_{n \times p}$.
- $C = AB$ – product of two matrices.
- Each element c_{ij} of C is a dot product of i -th row of A and j -th column of B : $C = \{c_{ij}\}_{m \times p} = \{(A_i, B^j)\}_{m \times p}$
- Example $\mathbb{R}^{2 \times 2}$:
$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

Matrix Multiplication: Example

$$\begin{bmatrix} 0 & 1 & 2 \\ 2 & 1 & 5 \end{bmatrix} \begin{bmatrix} 3 & 6 & 4 \\ 2 & 5 & 8 \\ 7 & 1 & 9 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 + 2 + 14 & 0 + 5 + 2 & 0 + 8 + 18 \\ 6 + 2 + 35 & 12 + 5 + 5 & 8 + 8 + 45 \end{bmatrix} =$$

$$= \begin{bmatrix} 16 & 7 & 26 \\ 43 & 22 & 61 \end{bmatrix}$$

To sum up

- Vectors
 - Vector spaces
 - Inner products
 - Lengths
 - Distances
 - Angles
- Analytic Geometry
 - Projections
 - Hyperplanes
 - Normal vector
- Vector spaces
 - Linear (in)dependence
 - Basis
- Matrices
 - Matrix operations

Next Time

- Calculus recap
- Probability theory recap