

Basic identities of matrix/vector ops

$$(A \cdot B)^T = A^T \cdot B^T \quad (AB)^T = B^T \cdot A^T \quad (A^{-1})^T = (A^T)^{-1} \\ (AB)^{-1} = B^{-1} \cdot A^{-1}$$

For $A \in \mathbb{R}^{m \times n}$ A_{ij} is the i th **ROW** then j th **COLUMN**

$$(A^T)^T = A \quad (AB)_{ij} = \sum_k A_{ik} \cdot B_{kj}$$

$$(A \cdot x)_i = \sum_j A_{ij} \cdot x_j \quad x^T \cdot y = y^T \cdot x = x \cdot y = \sum_i x_i \cdot y_i$$

$$x^T \cdot A \cdot x = \sum_{i,j} A_{ij} x_i x_j$$

Scalar-multiplication + addition distributes over:

column-blocks \rightarrow

$$A \cdot B = [A_1 | \dots | A_n] \cdot [B_1 | \dots | B_n] = [A_1 \cdot B_1 | \dots | A_n \cdot B_n]$$

row-blocks \rightarrow

$$A \cdot B = [A_1 | \dots | A_n] \cdot [B_1 | \dots | B_n] = [A_1 \cdot B_1 | \dots | A_n \cdot B_n]$$

Matrix-multiplication distributes over:

column-blocks \rightarrow $AB \cdot AC = A \cdot (B \cdot C) = [A \cdot B_1 | \dots | A \cdot B_n]$

row-blocks \rightarrow $AB \cdot AC = A \cdot (B \cdot C) = [A \cdot B_1 | \dots | A \cdot B_n]$

outer-product sum \rightarrow

$$AB = [A_1 | \dots | A_n] \cdot [B_1 | \dots | B_n] = \sum_{i,j} A_i B_j$$

e.g. for $A = [A_1 | \dots | A_n]$ $B = [B_1 | \dots | B_n]$ $\rightarrow AB = \sum_i A_i B_i$

What is a projection

A **projection** $\Pi: V \rightarrow V$ is an **endomorphism** such that $\Pi^2 = \Pi$ i.e. it leaves its image unchanged (its idempotent).

A **square matrix** such that $P^2 = P$ is called a **projection matrix**

It is called an **orthogonal projection matrix** if $P^2 = P$ and $P^T = P$ (conjugate-transpose)

Eigenvalues of a **projection matrix** must be 0 or 1

Because $\Pi: V \rightarrow V$ is a **linear map**, its **image space** $U = \text{im}(\Pi)$ and **null space** $W = \text{ker}(\Pi)$ are **subspaces of V**

$\Pi|_U$ is the **identity operator on U**

The **linear map** $\Pi^T: V \rightarrow V$ is also a projection with $W = \text{im}(\Pi^T) = \text{ker}(\Pi) = U$ and $U = \text{im}(\Pi^T) = \text{ker}(\Pi)$ i.e. they swapped

$\Pi|_U$ is a projection **along U onto U**

$\Pi|_W$ is a projection **along U onto U**

Π^T is the **identity operator on W**

V can be decomposed as $V = U \oplus W$ meaning every vector $x \in V$ can be uniquely written as $x = u + w$

$\|x\| = \sqrt{\|u\|^2 + \|w\|^2}$ (**Cauchy-Schwartz inequality**)

$u \perp w$ and $u \cdot w = 0$

An **orthogonal projection** further satisfies $U \perp W$

i.e. the **image** and **kernel** of Π are **orthogonal subspaces**

infact they are each other's **orthogonal complements**, i.e. $U^\perp = W$, $W^\perp = U$ (because finite-dimensional vector spaces)

so we have $\Pi(x) \cdot y = \Pi(x) \cdot (y - \Pi(y)) = 0$ or equivalently, $\Pi(x) \cdot (y - \Pi(y)) = \Pi(x) \cdot (y - \Pi(y)) = 0$

Projection properties

By Cauchy-Schwartz inequality we have $\|\Pi(x)\| \leq \|x\|$

The **orthogonal projection onto the line** containing vector u is $\Pi = \frac{u u^T}{u^T u}$ which can also be written as

$$\Pi = \frac{u u^T}{u^T u}$$

so we get

$$\Pi^T = \frac{(u u^T)^T}{(u u^T)^T} = \frac{u u^T}{u^T u} = \Pi$$

A special case of $\Pi(x) = \frac{u u^T}{u^T u} x$ is $\Pi(x) = \frac{u u^T}{u^T u} x$ since $\Pi(u) = u$

If $U \subset \mathbb{R}^n$ is a **linear** (one-dimensional) subspace with **orthonormal basis (ONB)** $(u_1, \dots, u_m) \in \mathbb{R}^n$

Let $U = \text{span}\{u_1, \dots, u_m\}$ be the matrix of columns

Then **orthogonal projection onto the subspace U** is $\Pi_U = U U^T$

Can be rewritten as $\Pi_U(x) = \sum_{i=1}^m (x \cdot u_i) u_i$

If (u_1, \dots, u_m) is **not orthonormal**, then "normalizing factor" $(U^T U)^{-1}$ is added $\rightarrow \Pi_U = U (U^T U)^{-1} U^T$

For **line** subspaces $U = \text{span}\{u\}$ we have

$$(U^T U)^{-1} = (u^T u)^{-1} = 1/(u \cdot u) = 1/\|u\|^2$$

Gram-Schmidt method to generate orthonormal basis from any linearly independent vectors

Gram-Schmidt is **iterative** (if what is a projection/projection) so we use **current j**dim subspace

Assume **orthonormal basis (ONB)** $(q_1, \dots, q_{j-1}) \in \mathbb{R}^m$

for j th dim subspace $U_j \subset \mathbb{R}^m$

Let $Q_j = [q_1 | \dots | q_j] \in \mathbb{R}^{m \times j}$ be the matrix of columns

$Q_j^T Q_j = I_j$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

$P_j = I_m - Q_j Q_j^T$ is the **orthogonal projection** properties [orthogonal projection]

Orthogonal complement

Assume $q_1, \dots, q_j \in U_j \Rightarrow$ unique decomposition

$$q_{j+1} = \sum_{i=1}^j \alpha_i q_i + r$$

\rightarrow discard r !

\rightarrow we're after this!

\rightarrow we have **next ONB** (q_1, \dots, q_{j+1})

for $U_{j+1} \Rightarrow$ start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

Orthogonal complement

Assume $q_1, \dots, q_j \in U_j \Rightarrow$ unique decomposition

$$q_{j+1} = \sum_{i=1}^j \alpha_i q_i + r$$

\rightarrow discard r !

\rightarrow we're after this!

\rightarrow we have **next ONB** (q_1, \dots, q_{j+1})

for $U_{j+1} \Rightarrow$ start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

\rightarrow start next iteration

Generalised Eigenvalues
-TODD: this seems low-priority, do when have time
-gen-eigenvalues
-Jordan chains (common cases)
https://www.youtube.com/watch?v=ATh6pefJAQ&list=PLM3uG5w5index&...
-JNF, form
-some tips on how to solve common cases
-JNF decomposition and basis of generalized eigenvectors

General: visualizing transformations of matrices

-TODD: do when have time -> where standard basis-vectors map to
-TODD: rotations, reflections, scaling, shearing, etc
Cholesky Decomposition
-Consider **positive (semi-)definite** $A \in \mathbb{R}^{n \times n}$
-**Cholesky Decomposition** is $A = LL^T$ where L is lower-triangular
-For positive semi-definite => **always exists**, but **non-unique**
-For positive-definite => **always uniquely exists** s.t. diagonals of L are positive
-Finding a Cholesky Decomposition
-Compute LL^T and solve $A = LL^T$ by matching terms
-For square roots always pick positive
-If there is **exact solution** then **positive-definite**
-If there are **free variables** at the end, then **positive semi-definite**
i.e. the decomposition is a **solution-set** parameterized on **free variables**

e.g. $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix} = L^T$ where $L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & c & \sqrt{1-c^2} \end{bmatrix}, c \in [0,1]$

-If $A = (L^T)$ you can use [[Forward/backward substitution]/forward/backward substitution]] to solve equations
-For $Ax=b$ just let $y=L^T x$
-Solve $Ly=b$ by forward substitution to find y
-Solve $L^T x=y$ by backward substitution to find x
-For $n=3$ => $L = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix}$

Lines and hyperplanes in Euclidean spaces

Forward/backward substitution
-Forward substitution: for lower-triangular
 $L = \begin{bmatrix} l_{11} & & 0 \\ & \ddots & \\ l_{n,1} & & l_{n,n} \end{bmatrix}$
-For $Lx=b$ just solve the first row
 $x_1, x_1, x_1 = x_1 = \frac{b_1}{l_{1,1}}$ and substitute down
-Then solve the second row
 $l_{2,1}x_1 + l_{2,2}x_2 = b_2 \Rightarrow x_2 = \frac{b_2 - l_{2,1}x_1}{l_{2,2}}$ and substitute down
-... and so on until all x_i are solved
-Backward substitution: for upper-triangular
 $U = \begin{bmatrix} u_{1,n} & & u_{1,1} \\ & \ddots & \\ u_{n,n} & & u_{n,n} \end{bmatrix}$
-For $Ux=b$ just solve the last row
 $u_{n,n}x_n = b_n \Rightarrow x_n = \frac{b_n}{u_{n,n}}$ and substitute up
-Then solve the second last row
 $u_{n-1,n-1}x_{n-1} + u_{n-1,n}x_n = b_{n-1} \Rightarrow x_{n-1} = \frac{b_{n-1} - u_{n-1,n}x_n}{u_{n-1,n-1}}$ and substitute up
-... and so on until all x_i are solved

Thin QR Decomposition w/ Gram-Schmidt (GS)

-Consider full-rank $A = [a_1 | \dots | a_n] \in \mathbb{R}^{m \times n}$ ($m \geq n$), i.e. $a_1, \dots, a_n \in \mathbb{R}^m$ are linearly independent
-Apply [[tutorial 1]Gram-Schmidt method to generate orthonormal basis from any linearly independent vectors][GS] $q_1, \dots, q_n = GS(a_1, \dots, a_n)$ to build **ONB** $(q_1, \dots, q_n) \in \mathbb{R}^m$ for $C(A)$
-For exams: more efficient to compute as $u_1 = a_1, \dots, u_{j-1} = q_{j-1} - \text{proj}_{\text{span}\{u_1, \dots, u_{j-1}\}} a_j$
1) Gather $q_j = [q_1 | \dots | q_j] \in \mathbb{R}^{m \times j}$ **all-at-once**
2) Compute $q_j = [q_1 | \dots | q_j] \in \mathbb{R}^{m \times j}$ **all-at-once**
3) Compute $Q_j \in \mathbb{R}^m$ and subtract from a_{j+1} **all-at-once**

-Can now rewrite $a_j = \sum_{i=1}^j q_i \cdot q_i^T a_j = Q_j c_j$
-Choose $Q_n = [q_1 | \dots | q_n] \in \mathbb{R}^{m \times n}$ I notice its since [[tutorial 1]Orthogonality concepts][semi-orthogonal] since $Q^T Q = I_n$
-Notice $\Rightarrow a_j = Q_j c_j = Q_1 q_1^T a_j + \dots + Q_{j-1} q_{j-1}^T a_j + Q_j c_j = Q_j^T \cdot Q_j^T a_j$
-Let $R = [r_{11} | \dots | r_{nn}] \in \mathbb{R}^{n \times n}$
 $A = QR = Q \begin{bmatrix} r_{11} & & 0 \\ & \ddots & \\ 0 & & r_{nn}^T \end{bmatrix}$ notice its
[[tutorial 1]Properties of matrices][upper-triangular]

Full QR Decomposition
-Consider full-rank $A = [a_1 | \dots | a_n] \in \mathbb{R}^{m \times n}$ ($m \geq n$), i.e. $a_1, \dots, a_n \in \mathbb{R}^m$ are linearly independent
-Apply [[thin QR Decomposition w/ Gram-Schmidt (GS)](thin QR decomposition)] to obtain:
-ONB $(q_1, \dots, q_n) \in \mathbb{R}^m$ for $C(A)$
-Semi-orthogonal $Q_1 = [q_1 | \dots | q_n] \in \mathbb{R}^{m \times n}$ and upper-triangular $R_1 \in \mathbb{R}^{n \times n}$ where $A = Q_1 R_1$
-[[tutorial 1]FFRCS Computing orthonormal vector-set extensions][Compute basis extension]] to obtain remaining $q_{n+1}, \dots, q_m \in \mathbb{R}^m$ where (q_1, \dots, q_m) is **ONB** for \mathbb{R}^m
-Notice (q_{n+1}, \dots, q_m) is **ONB** for $C(A)^\perp = \ker(A^T)$
-Let $Q_2 = [q_{n+1} | \dots | q_m] \in \mathbb{R}^{m \times (m-n)}$ let $Q = [Q_1 | Q_2] \in \mathbb{R}^{m \times m}$ let $R = [R_1 | 0_{(n-m) \times n}] \in \mathbb{R}^{m \times n}$
-Then full **QR decomposition** is $A = QR = [Q_1 | Q_2] \begin{bmatrix} R_1 \\ 0_{m-n} \end{bmatrix} = Q_1 R_1$
- Q_1 is **orthogonal**, i.e. $Q_1^{-1} = Q_1^T$ so its a basis transformation
- $\text{proj}_{C(A)} = Q_1 Q_1^T$, $\text{proj}_{C(A)^\perp} = Q_2 Q_2^T$ are [[tutorial 1]Projection properties][orthogonal projections] **onto** $C(A)$, $C(A)^\perp = \ker(A^T)$ respectively
-Notice: $QQ^T = I_m = Q_1 Q_1^T + Q_2 Q_2^T$
-Generalizable to $A \in \mathbb{C}^{m \times n}$ by changing transpose to conjugate-transpose
-Inner product $x^H y := x^T y^*$
-Orthogonal matrix $U^{-1} = U^T$ => unitary matrix $U^{-1} = U^H$
-For orthogonal $U = [u_1 | \dots | u_n] \in \mathbb{C}^{m \times n}$ => $\text{proj}_U = UU^H$ projects onto $C(U)$
-For unitary $U = [u_1 | \dots | u_n] \in \mathbb{C}^{m \times m}$ => $\text{proj}_U = UU^H$ projects onto $C(U)$
-A U so on...

Lines and hyperplanes in Euclidean spaces
-Consider **standard Euclidean space** $E^n(\mathbb{R}^n)$
-with standard basis $(e_1, \dots, e_n) \in \mathbb{R}^n$
-with standard origin $0 \in \mathbb{R}^n$
-A **line** $L \subset \mathbb{R}^n$ is characterized by direction $n \in \mathbb{R}^n$ ($n \neq 0$) and offset from origin $c \in L$
-It is customary that:
- n is a **unit vector**, i.e. $\|n\| = \|n\| = 1$
- $c \in L$ is the **closest point to origin**, i.e. $c \perp n$
-If $c \perp n$ => L is not vector-subspace of \mathbb{R}^n
-i.e. $0 \notin L$ i.e. L doesn't go through the origin
- L is affine-subspace of \mathbb{R}^n
-If $c \perp n$, i.e. L passes through the origin
- L has $\dim(L)=1$ and orthonormal basis (ONB) (\hat{n})
-A **hyperplane** is characterized by normal $n \in \mathbb{R}^n$ ($n \neq 0$) and offset from origin $c \in P$
-It represents an $(n-1)$ -dimensional slice of the n -dimensional space
-Points are hyperplanes for $n=1$
-Lines are hyperplanes for $n=2$
-Planes are hyperplanes for $n=3$
-It is customary that:
- n is a **unit vector**, i.e. $\|n\| = \|n\| = 1$
- $c \in P$ is the **closest point to origin**, i.e. $c \perp n$
-With those $\Rightarrow P = \{x \in \mathbb{R}^n | x \cdot n = c\}$
-If $c \cdot n = 0$ => P is not vector-subspace of \mathbb{R}^n
-i.e. $0 \notin P$ i.e. P doesn't go through the origin
- P is affine-subspace of \mathbb{R}^n
-If $c \cdot n = 0$, i.e. P passes through the origin
- P has $\dim(P)=n-1$
-Notice $L = \text{lin}(P)$ and $P = (Rn)^\perp$ are orthogonal complements, so:
- $\text{proj}_P = \hat{n}\hat{n}^T$ is orthogonal projection onto L **along** $\text{proj}_P = id_n - \text{proj}_L = I_n - \hat{n}\hat{n}^T$ is orthogonal projection onto P **(along L)**
- $L = \text{lin}(\text{proj}_L) = \ker(\text{proj}_P)$ and $P = \ker(\text{proj}_L) = \text{lin}(\text{proj}_P)$
- $Rn = Rn \cdot (Rn)^\perp$ i.e. all vectors $v \in Rn$ uniquely decomposed into $v = v_L + v_P$

Thin QR Decomposition w/ Gram-Schmidt (GS)
-Consider full-rank $A = [a_1 | \dots | a_n] \in \mathbb{R}^{m \times n}$ ($m \geq n$), i.e. $a_1, \dots, a_n \in \mathbb{R}^m$ are linearly independent
-Apply [[tutorial 1]Gram-Schmidt method to generate orthonormal basis from any linearly independent vectors][GS] $q_1, \dots, q_n = GS(a_1, \dots, a_n)$ to build **ONB** $(q_1, \dots, q_n) \in \mathbb{R}^m$ for $C(A)$
-For exams: more efficient to compute as $u_1 = a_1, \dots, u_{j-1} = q_{j-1} - \text{proj}_{\text{span}\{u_1, \dots, u_{j-1}\}} a_j$
1) Gather $q_j = [q_1 | \dots | q_j] \in \mathbb{R}^{m \times j}$ **all-at-once**
2) Compute $q_j = [q_1 | \dots | q_j] \in \mathbb{R}^{m \times j}$ **all-at-once**
3) Compute $Q_j \in \mathbb{R}^m$ and subtract from a_{j+1} **all-at-once**

Reflection w.r.t. hyperplanes and Householder Maps

-Two points $x, y \in E^n$ are reflections w.r.t hyperplane $P = (Rn)^\perp + c$ if:
1) The translation $\overrightarrow{xy} = y - x$ is **parallel** to normal \hat{n} , i.e. $\overrightarrow{xy} = \lambda \hat{n}$
2) Midpoint $m = 1/2(x+y) \in P$ lies on P , i.e. $m \cdot n = c \cdot n$
-Suppose $P = (Ru)^\perp$ goes through the origin with unit normal $u \in \mathbb{R}^n$
-Householder matrix $H_u = I_n - 2uu^T$ is reflection w.r.t. hyperplane P_u
-Recall: let $L_u = Ru$
- $\text{proj}_{L_u} = uu^T$ and $\text{proj}_{P_u} = I_n - uu^T$ => $\text{proj}_{P_u} = \text{proj}_{L_u} - \text{proj}_{L_u}$
-Visualize as preserving component in P_u then flipping component in L_u
- H_u is involutory, orthogonal and symmetric, i.e. $H_u = H_u^{-1} = H_u^T$

Modified Gram-Schmidt

-Go check [[tutorial 1]Gram-Schmidt method] to generate orthonormal basis from any linearly independent vectors[Classical GM] first, as this is just an alternative computation method
-Let $P_{j-1} q_j = I_m - Q_j^T q_j$ be **projector** onto $\{[a_1 | \dots | a_{j-1}]$
5Lines and hyperplanes in Euclidean space
mathbb{b}(E^n) (n)=[-] mathbb{b}(R) (n)=[hyperplane] $(Rq_1)^\perp$ i.e. [[tutorial 5]Lines and hyperplanes in Euclidean space & mathbb{b}(E^n) (n)=[hyperplane] (n)=[orthogonal complement] of line Rq_1
-Notice: $P_{j-1} = I_m - Q_j^T q_j = \prod_{i=1}^j (I_m - q_i q_i^T) = \prod_{i=1}^j P_{q_i}$
-Overall, its an N -th order partial derivative where $N = \sum_{i=1}^n \nabla_i$
- $\nabla f = [\partial_1 f, \dots, \partial_n f]^T$ is gradient of f => $(\nabla f)_i = \frac{\partial f}{\partial x_i}$
- $\nabla^T f = (\nabla f)^T$ is transpose of ∇f i.e. $\nabla^T f$ is row vector
- $\partial_u f(x) = \lim_{\delta \rightarrow 0} \frac{f(x+\delta u) - f(x)}{\delta}$ **directional-derivative** of f
-It is rate-of-change in direction u where $u \in \mathbb{R}^n$ is unit-vector
- $\partial_u f(x) = \nabla f(x) \cdot u = [\nabla f(x) | u]$ cos(0) => $D_u f(x)$ **maximized** when cos(0)=1
-i.e. when x, u are parallel => hence $\nabla f(x)$ is direction of **max**, rate-of-change
- $H(f) = \nabla^2 f = \nabla(\nabla f)^T$ is the **Hessian** of f => $H(f)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$
-Re-state: $u_j = I_m - Q_j^T q_j^T a_{j+1}$ => $\text{proj}_{L_u} = \prod_{i=1}^j P_{q_i} a_{j+1} = (\prod_{i=1}^j P_{q_i} a_{j+1}) = P_{a_{j+1}} a_{j+1}$
-Projectors P_{a_1}, \dots, P_{a_j} are iteratively applied to a_{j+1} removing its components along q_1, \dots, q_j and so on...

Lines and hyperplanes in Euclidean spaces

-Consider **standard Euclidean space** $E^n(\mathbb{R}^n)$
-with standard basis $(e_1, \dots, e_n) \in \mathbb{R}^n$
-with standard origin $0 \in \mathbb{R}^n$
-A **line** $L \subset \mathbb{R}^n$ is characterized by direction $n \in \mathbb{R}^n$ ($n \neq 0$) and offset from origin $c \in L$
-It is customary that:
- n is a **unit vector**, i.e. $\|n\| = \|n\| = 1$
- $c \in L$ is the **closest point to origin**, i.e. $c \perp n$
-If $c \perp n$ => L is not vector-subspace of \mathbb{R}^n
-i.e. $0 \notin L$ i.e. L doesn't go through the origin
- L is affine-subspace of \mathbb{R}^n
-If $c \perp n$, i.e. L passes through the origin
- L has $\dim(L)=1$ and orthonormal basis (ONB) (\hat{n})
-A **hyperplane** is characterized by normal $n \in \mathbb{R}^n$ ($n \neq 0$) and offset from origin $c \in P$
-It represents an $(n-1)$ -dimensional slice of the n -dimensional space
-Points are hyperplanes for $n=1$
-Lines are hyperplanes for $n=2$
-Planes are hyperplanes for $n=3$
-It is customary that:
- n is a **unit vector**, i.e. $\|n\| = \|n\| = 1$
- $c \in P$ is the **closest point to origin**, i.e. $c \perp n$
-With those $\Rightarrow P = \{x \in \mathbb{R}^n | x \cdot n = c\}$
-If $c \cdot n = 0$ => P is not vector-subspace of \mathbb{R}^n
-i.e. $0 \notin P$ i.e. P doesn't go through the origin
- P is affine-subspace of \mathbb{R}^n
-If $c \cdot n = 0$, i.e. P passes through the origin
- P has $\dim(P)=n-1$
-Notice $L = \text{lin}(P)$ and $P = (Rn)^\perp$ are orthogonal complements, so:
- $\text{proj}_P = \hat{n}\hat{n}^T$ is orthogonal projection onto L **along** $\text{proj}_P = id_n - \text{proj}_L = I_n - \hat{n}\hat{n}^T$ is orthogonal projection onto P **(along L)**
- $L = \text{lin}(\text{proj}_L) = \ker(\text{proj}_P)$ and $P = \ker(\text{proj}_L) = \text{lin}(\text{proj}_P)$
- $Rn = Rn \cdot (Rn)^\perp$ i.e. all vectors $v \in Rn$ uniquely decomposed into $v = v_L + v_P$

Classical vs. Modified Gram-Schmidt (for thin QR)
-These algorithms both compute [[tutorial 5]Thin QR Decomposition w/ Gram-Schmidt (GS)](thin QR decomposition)]
-Where $A \in \mathbb{R}^{m \times n}$ is full-rank, $Q \in \mathbb{R}^{m \times m}$ is semi-orthogonal, and $R \in \mathbb{R}^{n \times n}$ is upper-triangular
Classical vs. Modified Gram-Schmidt (for thin QR)
-These algorithms both compute [[tutorial 5]Thin QR Decomposition w/ Gram-Schmidt (GS)](thin QR decomposition)]
-Where $A \in \mathbb{R}^{m \times n}$ is full-rank, $Q \in \mathbb{R}^{m \times m}$ is semi-orthogonal, and $R \in \mathbb{R}^{n \times n}$ is upper-triangular
-Computes at j th step:
-Classical GS => j th column of Q and the j th row of R
-Modified GS => j th column of Q and the j th row of R
-Both have **float (floating-point operation)** count of $O(2mn^2)$
-NOTE: Householder method has $2(mn^2 - n^3)/3$ (float count), but better numerical properties
-Recall: $Q^T Q = I_n$ => check for loss of orthogonality with $\|I_n - Q^T Q\| = \text{loss}$
-Classical GS => $\|I_n - Q^T Q\| = \text{Cond}(A)^2 \epsilon_{\text{mach}}$
-Modified GS => $\|I_n - Q^T Q\| = \text{Cond}(A) \epsilon_{\text{mach}}$
-NOTE: Householder method has $\|I_n - Q^T Q\| \leq \epsilon_{\text{mach}}$

Multivariate Calculus

-Consider $f: \mathbb{R}^n \rightarrow \mathbb{R}$ when clear write i th component in input as x_i instead of x
-Level curve w.r.t. to $c \in \mathbb{R}$ is all points s.t. $f(x) = c$
-Projecting level curves onto \mathbb{R}^n gives **contour-map** of f
- n th order partial derivative w.r.t. i_k of f ..., of i_k th order partial derivative w.r.t. i_k of f
 $\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}$
-Overall, its an N -th order partial derivative where $N = \sum_{i=1}^n \nabla_i$
- $\nabla f = [\partial_1 f, \dots, \partial_n f]^T$ is gradient of f => $(\nabla f)_i = \frac{\partial f}{\partial x_i}$
- $\nabla^T f = (\nabla f)^T$ is transpose of ∇f i.e. $\nabla^T f$ is row vector
- $\partial_u f(x) = \lim_{\delta \rightarrow 0} \frac{f(x+\delta u) - f(x)}{\delta}$ **directional-derivative** of f
-It is rate-of-change in direction u where $u \in \mathbb{R}^n$ is unit-vector
- $\partial_u f(x) = \nabla f(x) \cdot u = [\nabla f(x) | u]$ cos(0) => $D_u f(x)$ **maximized** when cos(0)=1
-i.e. when x, u are parallel => hence $\nabla f(x)$ is direction of **max**, rate-of-change
- $H(f) = \nabla^2 f = \nabla(\nabla f)^T$ is the **Hessian** of f => $H(f)_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$
-Re-state: $u_j = I_m - Q_j^T q_j^T a_{j+1}$ => $\text{proj}_{L_u} = \prod_{i=1}^j P_{q_i} a_{j+1} = (\prod_{i=1}^j P_{q_i} a_{j+1}) = P_{a_{j+1}} a_{j+1}$
-Projectors P_{a_1}, \dots, P_{a_j} are iteratively applied to a_{j+1} removing its components along q_1, \dots, q_j and so on...

Big-O meaning for numerical analysis
-In complexity analysis $f(n) = O(g(n))$ as $n \rightarrow \infty$
-But in numerical analysis $f(\epsilon) = O(g(\epsilon))$ as $\epsilon \rightarrow 0$
-i.e. $\limsup_{\epsilon \rightarrow 0} \frac{f(\epsilon)}{g(\epsilon)} < \infty$
-i.e. $\exists C, \delta > 0$ s.t. $\forall \epsilon \in [0, \delta]$ we have $0 \leq f(\epsilon) \leq C g(\epsilon)$
- $O(g)$ is set of functions $\{f : \limsup_{\epsilon \rightarrow 0} \frac{f(\epsilon)}{g(\epsilon)} < \infty\}$
-Smallest partial order $O(g_1) \subseteq O(g_2)$ defined by set-inclusion $O(g_1) \subseteq O(g_2)$
-i.e. as $\epsilon \rightarrow 0$, $g_1(\epsilon)$ goes to zero faster than $g_2(\epsilon)$
-Roughly same hierarchy as complexity analysis but flipped (some break pattern)
-e.g. $\dots, O(\epsilon^{-2}) \subset O(\epsilon^{-1}) \subset O(\epsilon) \subset O(1)$
-Maximum: $O(\max\{g_1, g_2\}) = O(g_2) \iff O(g_1) \subseteq O(g_2)$
-e.g. $O(\max\{\epsilon^2, \epsilon\}) = O(\epsilon)$
-Using functions f_1, \dots, f_n let $\tilde{f}([f_1, \dots, f_n])$ be formula defining some function
-Then $\tilde{f}(O(g_1), \dots, O(g_n))$ is the class of functions $\{\tilde{f}(f_1, \dots, f_n) : f_1 \in O(g_1), \dots, f_n \in O(g_n)\}$
-e.g. $O(1) = \{f(f) : f \in O(1)\}$
-General case:
 $\tilde{f}_1(O(f_1), \dots, O(f_m)) = \tilde{f}_2(O(g_1), \dots, O(g_n))$ means $\tilde{f}_1(O(f_1), \dots, O(f_m)) \subseteq \tilde{f}_2(O(g_1), \dots, O(g_n))$
-e.g. $O(1) \subseteq O(\epsilon^k)$ means $\{f(f) : f \in O(1)\} \subseteq O(\epsilon^k)$ not necessarily true
-Special case: $f = \tilde{f}(O(g_1), \dots, O(g_n))$ means $f \in \tilde{f}(O(g_1), \dots, O(g_n))$
-e.g. $\epsilon \in \epsilon^{-1} \epsilon^2 = \epsilon^2$ means $\epsilon \in \epsilon^{-1} \epsilon^2$
-e.g. $\epsilon \in \epsilon^{-1} \epsilon^2 = \epsilon^2$ means $\epsilon \in \epsilon^{-1} \epsilon^2$ not necessarily true
-Let $f_1 = O(g_1)$, $f_2 = O(g_2)$ and let \tilde{f} be a constant
- $\tilde{f}_1 \tilde{f}_2 = O(g_1 g_2)$ and $\tilde{f} \cdot O(g) = O(g)$
- $\tilde{f}_1 + \tilde{f}_2 = O(\max\{g_1, g_2\})$ => if $g_1 = g_2$ then $\tilde{f}_1 + \tilde{f}_2 = O(g)$
- $O(R) \cdot g = O(g)$

Floating-point numbers

-Consider **base/radix** $b \geq 2$ (typically 2) and precision ℓ (24 or 53 for IEEE single/double precisions)
-Floating-point numbers are discrete subsets $F = \{\pm 1^k (m/b^k) \mid 1 \leq m \leq b^k, s \in \{b, m, e, z\}\}$
- s is sign-bit, m/b^k is mantissa, e is exponent (8 bit for single, 11 bit for double)
-Equivalently, can restrict to $b^{-1} \leq m \leq b^{-1}$ for unique m and e
- F is not idealized (ignores overflow/underflow), so is countably infinite and self-similar (i.e. $F = bF$)
-For all $x \in \mathbb{R}$ there exists $f(x) \in F$ s.t. $|x - f(x)| \leq \epsilon_{\text{mach}} |x|$
-Equivalently $f(x) = x(1 + \delta)$, $|\delta| \leq \epsilon_{\text{mach}}$
-Machine epsilon $\epsilon_{\text{machine}} = \epsilon_{\text{mach}} = \frac{1}{2} b^{-1}$ is maximum relative gap between FPS
-Half the gap between 1 and next largest FP $2^{-24} \approx 5.96 \times 10^{-8}$ and $2^{-53} \approx 1.07 \times 10^{-16}$ for single/double
-FP arithmetic: let \tilde{x}, \tilde{y} be real and floating counterparts of arithmetic operation
-For $x, y \in F$ we have $x \tilde{y} = f(x \cdot y) = (x \cdot y)(1 + \epsilon)$, $|\epsilon| \leq \epsilon_{\text{mach}}$

Stability

-Given a problem $f: X \rightarrow Y$, an algorithm for f is $f: X \rightarrow Y$
- f is **computer implementation**, so inputs/outputs are FP
-Input $x \in X$ is first rounded to $f(x)$, i.e. $f(x) = f(f(x))$
- f cannot be continuous (for the most part)
-Absolute error => $\|f(x) - f(x)\|$ relative error => $\frac{\|f(x) - f(x)\|}{\|f(x)\|}$
- f is accurate if $\forall x \in X, \frac{\|f(x) - f(x)\|}{\|f(x)\|} = O(\epsilon_{\text{mach}})$
- f is stable if $\forall x \in X, \frac{\|f(x) - f(x)\|}{\|f(x)\|} = O(\epsilon_{\text{mach}})$ and $\frac{\|x - \tilde{x}\|}{\|x\|} = O(\epsilon_{\text{mach}})$
-i.e. nearly the right answer to nearly the right question
-outer-product is stable
-if backwards stable f and f has condition number $\kappa(x)$ then relative error $\frac{\|f(x) - f(x)\|}{\|f(x)\|} = O(\kappa(x) \epsilon_{\text{mach}})$
-Accuracy, stability, backwards stable are necessary conditions for fin-dim X, Y

Big-O meaning for numerical analysis

-In complexity analysis $f(n) = O(g(n))$ as $n \rightarrow \infty$
-But in numerical analysis $f(\epsilon) = O(g(\epsilon))$ as $\epsilon \rightarrow 0$
-i.e. $\limsup_{\epsilon \rightarrow 0} \frac{f(\epsilon)}{g(\epsilon)} < \infty$
-i.e. $\exists C, \delta > 0$ s.t. $\forall \epsilon \in [0, \delta]$ we have $0 \leq f(\epsilon) \leq C g(\epsilon)$
- $O(g)$ is set of functions $\{f : \limsup_{\epsilon \rightarrow 0} \frac{f(\epsilon)}{g(\epsilon)} < \infty\}$
-Smallest partial order $O(g_1) \subseteq O(g_2)$ defined by set-inclusion $O(g_1) \subseteq O(g_2)$
-i.e. as $\epsilon \rightarrow 0$, $g_1(\epsilon)$ goes to zero faster than $g_2(\epsilon)$
-Roughly same hierarchy as complexity analysis but flipped (some break pattern)
-e.g. $\dots, O(\epsilon^{-2}) \subset O(\epsilon^{-1}) \subset O(\epsilon) \subset O(1)$
-Maximum: $O(\max\{g_1, g_2\}) = O(g_2) \iff O(g_1) \subseteq O(g_2)$
-e.g. $O(\max\{\epsilon^2, \epsilon\}) = O(\epsilon)$
-Using functions f_1, \dots, f_n let $\tilde{f}([f_1, \dots, f_n])$ be formula defining some function
-Then $\tilde{f}(O(g_1), \dots, O(g_n))$ is the class of functions $\{\tilde{f}(f_1, \dots, f_n) : f_1 \in O(g_1), \dots, f_n \in O(g_n)\}$
-e.g. $O(1) = \{f(f) : f \in O(1)\}$
-General case:
 $\tilde{f}_1(O(f_1), \dots, O(f_m)) = \tilde{f}_2(O(g_1), \dots, O(g_n))$ means $\tilde{f}_1(O(f_1), \dots, O(f_m)) \subseteq \tilde{f}_2(O(g_1), \dots, O(g_n))$
-e.g. $O(1) \subseteq O(\epsilon^k)$ means $\{f(f) : f \in O(1)\} \subseteq O(\epsilon^k)$ not necessarily true
-Special case: $f = \tilde{f}(O(g_1), \dots, O(g_n))$ means $f \in \tilde{f}(O(g_1), \dots, O(g_n))$
-e.g. $\epsilon \in \epsilon^{-1} \epsilon^2 = \epsilon^2$ means $\epsilon \in \epsilon^{-1} \epsilon^2$
-e.g. $\epsilon \in \epsilon^{-1} \epsilon^2 = \epsilon^2$ means $\epsilon \in \epsilon^{-1} \epsilon^2$ not necessarily true
-Let $f_1 = O(g_1)$, $f_2 = O(g_2)$ and let \tilde{f} be a constant
- $\tilde{f}_1 \tilde{f}_2 = O(g_1 g_2)$ and $\tilde{f} \cdot O(g) = O(g)$
- $\tilde{f}_1 + \tilde{f}_2 = O(\max\{g_1, g_2\})$ => if $g_1 = g_2$ then $\tilde{f}_1 + \tilde{f}_2 = O(g)$
- $O(R) \cdot g = O(g)$

LU factorization w/ Gaussian elimination

-[[tutorial 1]Representing ERos/ECOs as transformation matrices][Recall that]] you can represent ERos and ECOs as transformation matrices R, C respectively
-LU factorization => finds $A = LU$ where $L,$

*Similar to to [[tutorial 1#Gram-Schmidt method to generate orthonormal basis from any linearly independent vectors|Gram-Schmidt]] (*different*

- Q is unitary, i.e. $Q^\dagger = Q^{-1}$ and upper-triangular U