

## Lesson 4

Тест, 8 вопроса

1

Баллы

1.

In which situation would we choose to use a Metropolis-Hastings (or any MCMC) sampler rather than straightforward Monte Carlo sampling?

- ☐ The data (likelihood) come from a Markov chain.
  - ☐ Monte Carlo estimation is easier than calculating the integral required to obtain the mean of the target distribution.
  - ☒ There is no easy way to simulate independent draws from the target distribution.
  - ☐ The target distribution follows a Markov chain.
- 

1

Баллы


2.

# Lesson 4


## Metropolis-Hastings algorithm to sample the target distribution whose PDF is shown below?

Тест, 8 вопроса


**Note:** In independent Metropolis-Hastings, the candidate-generating distribution  $q$  does not depend on the previous iteration of the chain.



☐  $q = N(15, 3.1^2)$




☐  $q = \text{Unif}(0.05, 25.0)$




## Lesson 4

Тест, 8 вопроса



☒  $q = \text{Gamma}(3.0, 0.27)$



1

Баллы

3.

If we employed an independent Metropolis-Hastings algorithm (in which the candidate-generating distribution  $q$  does not depend on the previous iteration of the chain), what would happen if we skipped the acceptance ratio step and always accepted candidate draws?

- ☒ The resulting sample would be a Monte Carlo simulation from  $q$  instead of from the target distribution.
- ☐ The sampler would become more efficient because we are no longer discarding draws.
- ☐ The chain would explore the posterior distribution very slowly, requiring more samples.
- ☐ Each draw could be considered as a sample from the target distribution.

## Lesson 4

Тест, 8 вопроса

4.

If the target distribution  $p(\theta) \propto g(\theta)$  is for a positive-valued random variable so that  $p(\theta)$  contains the indicator function  $I_{\theta>0}(\theta)$ , what would happen if a random walk Metropolis sampler proposed the candidate  $\theta^* = -0.3$ ?


- ☐ The candidate would be accepted with probability 0.3 because  $g(\theta^*) = |\theta^*|$ , yielding an acceptance ratio  $\alpha = 0.3$ .
  - ☒ The candidate would be rejected with probability 1 because  $g(\theta^*) = 0$ , yielding an acceptance ratio  $\alpha = 0$ .
  - ☐ The candidate would be accepted with probability 1 because  $g(\theta^*) = 0$ , yielding an acceptance ratio  $\alpha = 1$ .
  - ☐ The candidate would be accepted with probability 1 because  $g(\theta^*) = 0$ , yielding an acceptance ratio  $\alpha = \infty$ .
- 

1

Баллы

5.

Lesson 4  
 Suppose we use a random walk Metropolis sampler with normal proposals (centered on the current value of the chain) to sample from the target distribution whose PDF is shown below. The chain is currently at  $\theta_i = 15.0$ . Which of the other points, if used as a candidate  $\theta^*$  for the next step, would yield the largest acceptance ratio  $\alpha$ ?



- ☐ A)  $\theta^* = 3.1$
- ☒ B)  $\theta^* = 9.8$
- ☐ C)  $\theta^* = 20.3$
- ☐ D)  $\theta^* = 26.1$

1

Баллы

6.

Suppose you are using a random walk Metropolis sampler with normal proposals. After sampling the chain for 1000 iterations, you notice that the acceptance rate for the candidate draws is only 0.02. Which corrective action is most likely to help you approach a better acceptance rate (between 0.23 and 0.50)?

- ☐ Replace the normal proposal distribution with a uniform proposal distribution centered on the previous value and variance equal to that of the old normal proposal distribution.
- ☐ Fix the mean of the normal proposal distribution at the last accepted candidate's value. Use the new mean for all future proposals.

## Lesson 4

Тест, 8 вопроса

Increase the variance of the normal proposal distribution  $q$ .Decrease the variance of the normal proposal distribution  $q$ .

1

Баллы

7.

Suppose we use a random walk Metropolis sampler to sample from the target distribution  $p(\theta) \propto g(\theta)$  and propose candidates  $\theta^*$  using the  $\text{Unif}(\theta_{i-1} - \epsilon, \theta_{i-1} + \epsilon)$  distribution where  $\epsilon$  is some positive number and  $\theta_{i-1}$  is the previous iteration's value of the chain. What is the correct expression for calculating the acceptance ratio  $\alpha$  in this scenario?

**Hint:** Notice that the  $\text{Unif}(\theta_{i-1} - \epsilon, \theta_{i-1} + \epsilon)$  distribution is centered on the previous value and is symmetric (since the PDF is flat and extends the same distance  $\epsilon$  on either side).

☐  $\alpha = \frac{\text{Unif}(\theta_{i-1} | \theta^* - \epsilon, \theta^* + \epsilon)}{\text{Unif}(\theta^* | \theta_{i-1} - \epsilon, \theta_{i-1} + \epsilon)}$  where  $\text{Unif}(\theta | a, b)$  represents the PDF of a  $\text{Unif}(a, b)$  evaluated at  $\theta$ .

☐  $\alpha = \frac{\text{Unif}(\theta^* | \theta_{i-1} - \epsilon, \theta_{i-1} + \epsilon)}{\text{Unif}(\theta_{i-1} | \theta^* - \epsilon, \theta^* + \epsilon)}$  where  $\text{Unif}(\theta | a, b)$  represents the PDF of a  $\text{Unif}(a, b)$  evaluated at  $\theta$ .

☒  $\alpha = \frac{g(\theta^*)}{g(\theta_{i-1})}$

☐  $\alpha = \frac{g(\theta_{i-1})}{g(\theta^*)}$

1

Баллы

8.

The following code completes one iteration of an algorithm to simulate a chain whose stationary distribution is  $\pi(\theta) \propto g(\theta)$ . Which algorithm is employed here?

Тест, 8 вопроса

```
1 # draw candidate
2 theta_cand = rnorm(n=1, mean=0.0, sd=10.0)
3
4 # evaluate log of g with the candidate
5 lg_cand = lg(theta=theta_cand)
6
7 # evaluate log of g at the current value
8 lg_now = lg(theta=theta_now)
9
10 # evaluate log of q at candidate
11 lq_cand = dnorm(theta_cand, mean=0.0, sd=10.0, log=TRUE)
12
13 # evaluate log of q at the current value
14 lq_now = dnorm(theta_now, mean=0.0, sd=10.0, log=TRUE)
15
16 # calculate the acceptance ratio
17 lalpha = lg_cand + lq_now - lg_now - lq_cand
18 alpha = exp(lalpha)
19
20 # draw a uniform variable which will be less than alpha with probability min(1,
    alpha)
21 u = runif(1)
22
23 if (u < alpha) { # then accept the candidate
24   theta_now = theta_cand
25   accpt = accpt + 1 # to keep track of acceptance
26 }
```

- ☒ Independent Metropolis-Hastings ( $q$  does not condition on the previous value of the chain) with normal proposal
- ☐ Random walk Metropolis with normal proposal
- ☐ Random walk Metropolis with uniform proposal
- ☐ Independent Metropolis-Hastings ( $q$  does not condition on the previous value of the chain) with uniform proposal

Оплатить курс



# Lesson 4

Тест, 8 вопроса



# Lesson 4

Тест, 8 вопроса

# Lesson 4

Тест, 8 вопроса