

ALMOST PERIODIC SOLUTIONS OF LATTICE DYNAMICAL SYSTEMS WITH MONOTONE NONLINEARITY

David Cheban,
Andrei Sultan *

*Vladimir Andrunachievici Institute of Mathematics and Computer Science, Moldova
State University, Chişinău, Republic of Moldova
david.cean@usm.md, andrew15sultan@gmail.com*

Let \mathbb{R} (respectively, \mathbb{Z}) be the set of all real (respectively, integer) numbers and l_2 be the Hilbert space of all sequences $\xi = (\xi_i)_{i \in \mathbb{Z}}$ ($\xi_i \in \mathbb{R}$ for any $i \in \mathbb{Z}$) with the property $\sum_{i \in \mathbb{Z}} |\xi_i|^2 < \infty$ equipped with the scalar product $\langle \xi, \eta \rangle := \sum_{i \in \mathbb{Z}} \xi_i \eta_i$. Denote by $C(\mathbb{R}, l_2)$ the family of all continuous functions $\varphi : \mathbb{R} \rightarrow l_2$ equipped with the compact-open topology and $(C(\mathbb{R}, l_2), \mathbb{R}, \sigma)$ the shift dynamical system on the space $C(\mathbb{R}, l_2)$.

A subset $A \subset \mathbb{R}$ is called relatively dense if there exists a positive number l such that $[a, a + l] \cap \mathbb{R} \neq \emptyset, \forall a \in \mathbb{R}$.

Definition. A function $\varphi \in C(\mathbb{R}, l_2)$ is said to be almost periodic [1] if for any $\varepsilon > 0$ there exists a relatively dense subset $\mathcal{P}(\varepsilon)$ such that $\|\varphi(t + \tau) - \varphi(t)\|_{l_2} < \varepsilon$

In this talk we study the almost periodic solutions of the systems

$$u'_i = \nu(u_{i-1} - 2u_i + u_{i+1}) - \lambda u_i + F(u_i) + f_i(t) \quad (i \in \mathbb{Z}), \quad (1)$$

where $\lambda, \nu > 0$, $F \in C(\mathbb{R}, \mathbb{R})$ and $f \in C(\mathbb{R}, l_2)$ ($f(t) := (f_i(t))_{i \in \mathbb{Z}}$ for any $t \in \mathbb{R}$) is an almost periodic function.

Theorem 1. Suppose that the following conditions hold:

1. the function $f \in C(\mathbb{R}, l_2)$ is almost periodic;
2. the function F possesses the following properties:
 - (a) it is locally Lipschitzian;
 - (b) there exists a positive number α such that $F(s)s \leq -\alpha s^2$ for any $s \in \mathbb{R}$;
 - (c) the function F is monotone [1], i.e., there exists a positive number β such that $(F(x_1) - F(x_2))(x_1 - x_2) \leq -\beta |x_1 - x_2|^2$ for any $x_1, x_2 \in \mathbb{R}$.

Then the equation (1) has a unique almost periodic solution.

Funding. This research was supported by the State Program of the Republic of Moldova "Monotone Nonautonomous Dynamical Systems (24.80012.5007.20SE)" and partially was supported by the Institutional Research Program 011303 "SATGED", Moldova State University.

ORCID (D. Cheban): <https://orcid.org/0000-0002-2309-3823>

ORCID (A. Sultan): <https://orcid.org/0009-0003-9785-9291>

References:

1. David N. Cheban. *Monotone Nonautonomous Dynamical Systems*. Springer Nature Switzerland AG, 2024.

* *Sultan Andrei:* A. A. Sultan