JP Journal of Algebra, Number Theory and Applications
© 2016 Pushpa Publishing House, Allahabad, India
Published Online: May 2016
http://dx.doi.org/10.17654/NT038030279
Volume 38, Number 3, 2016, Pages 279-293 **ISSN: 0972-5555**

ON THE NUMBER OF TWO-POINT ANTICHAINS IN THE POWERSET OF AN *n*-ELEMENT SET ORDERED BY INCLUSION

Agustín Moreno Cañadas[1], **Veronica Cifuentes Vargas**[2] **and Andrés Felipe Gonzalez**[2]

[1]Department of Mathematics
National University of Colombia
Bogotá, Colombia
e-mail: amorenoca@unal.edu.co

[2]Department of Mathematics
District University F.J.C.
Bogotá, Colombia
e-mail: vcifuentesv@unal.edu.co
afgonzalezb@correo.udistrital.edu.co

Abstract

Tiled orders are used to establish properties of the sequence whose elements count the number of two-point antichains in the powerset of an *n*-element set ordered by inclusion. In particular, we find a partition formula for numbers in this sequence.

Received: December 23, 2015; Revised: January 5, 2016; Accepted: January 19, 2016
2010 Mathematics Subject Classification: 05A17, 11D09, 11D45, 11D85, 11E25, 11E57.
Keywords and phrases: antichain, Auslander-Reiten quiver, categorification of an integer sequence, indecomposable module, integer partition, poset representation, tiled order, $(0, m)$- order.
Communicated by K. K. Azad

1. Introduction

This work follows ideas of Fahr and Ringel who, recently, found out partition formulas of Fibonacci numbers by establishing a categorical equivalence between the category of representations of the 3-Kronecker quiver and the category of representations of its corresponding universal covering [5, 6]. This allowed them to find the sequence A132262 in the OEIS. They also introduced the term categorification of a sequence of numbers which means to consider instead of these numbers a suitable object in a category.

Soon afterwards, the first author et al. used the same technique in order to obtain a categorification of the sequence A002662 which counts the number of subsets with at least three elements of an n-element set. In such a work, a partition formula for elements of the sequence was obtained via the algorithm of differentiation with respect to a maximal point (introduced by Nazarova and Roiter to classify different posets of tame representation type) and a Gabriel's identity which establish a relationship between the number of indecomposable representations in the category of representations $\operatorname{rep}\mathcal{P}$ of a given poset $\mathcal{P}$ and the number of indecomposable representations of the corresponding derived poset $\operatorname{rep}\mathcal{P}'$ [1]. These results were also used by the first author et al. to describe some higher dimensional partitions [2].

In this paper, we use tiled orders also called semimaximal rings by Zavadskij and Kirichenko to establish that a general formula obtained recently by the first author et al. for numbers in the sequence A016269 constitutes a partition formula for those numbers. We recall that numbers in such a sequence count the number of two-point antichains in the powerset of an n-element set ordered by inclusion denoted $\mathcal{P}_n = (2^{\mathbf{n}}, \subseteq)$ with $\mathbf{n} = \{1, 2, ..., n\}$ [3, 4, 11, 12]. We also prove that numbers in sequence A016269 are not prime numbers giving in this way a positive answer to a conjecture posed by the first author et al. in [4].

This paper is organized as follows. Some of the basic definitions and notations concerning the category of modules over tiled orders are included in Section 2. In Section 3, we establish identities for the number of two-point antichains in $\mathcal{P}_n$. Answers to questions posed by the first author et al. in [4] are given in Section 3 as well.

2. Preliminaries

This section introduces some other basic definitions, and notations regarding tiled orders to be used throughout the paper [1-6, 9].

2.1. Tiled orders

A field T is said to be of *discrete norm* or *discrete valuation* if it is endowed with a surjective map

$$\nu : T \to \mathbb{Z} \cup \{\infty\},$$

which satisfies the following conditions:

(a) $\nu(x) = \infty$ if and only if $x = 0$,

(b) $\nu(xy) = \nu(x) + \nu(y)$,

(c) $\nu(x + y) \geq \min\{\nu(x), \nu(y)\}$.

We let $\mathbb{O}$ denote, the *normalization ring* of the field T, such that

$$\mathbb{O} = \{x \in T \mid \nu(x) \geq 0\}.$$

An element $\pi \in \mathbb{O}$ such that $\nu(\pi) = 1$ is a *prime element* of $\mathbb{O}$. For each $x \in \mathbb{O}$, we have that $x \in \mathbb{O}$ if and only if $x = \varepsilon\pi^m$, for some $m \geq 0$ and $\varepsilon \in \mathbb{O}^*$. Moreover, $x \in T$ if and only if $x = \varepsilon\pi^m$ for some $m \in \mathbb{Z}$ and $\varepsilon \in \mathbb{O}^*$.

Ring $\mathbb{O}$ is such that $\mathbb{O} \supset \pi\mathbb{O}$, where $\pi\mathbb{O}$ is the unique maximal ideal,

therefore, ideals of $\mathbb{O}$ generate a chain of the form

$$\mathbb{O} \supset \pi\mathbb{O} \supset \pi^2\mathbb{O} \supset \cdots \supset \pi^m\mathbb{O} \supset \cdots.$$

A *tiled order* or *semimaximal ring* Λ is a subring of the matrix algebra $T^{n\times n}$ with the form

$$\Lambda = \sum_{i,\, j=1}^{n} e_{ij}\pi^{\lambda_{ij}}\mathbb{O} = \begin{bmatrix} \mathbb{O} & \pi^{\lambda_{12}}\mathbb{O} & \cdots & \pi^{\lambda_{1n}}\mathbb{O} \\ \pi^{\lambda_{21}}\mathbb{O} & \mathbb{O} & \cdots & \pi^{\lambda_{2n}}\mathbb{O} \\ \vdots & \vdots & \vdots & \vdots \\ \pi^{\lambda_{n1}}\mathbb{O} & \pi^{\lambda_{n2}}\mathbb{O} & \cdots & \mathbb{O} \end{bmatrix}.$$

Λ consists of all matrices whose entries ij belong to $\pi^{\lambda_{ij}}\mathbb{O}$, in this case, the $e_{ij} \in T^{n\times n}$ are unit matrices such that $e_{ij}e_{kl} = \delta_{jk}e_{il}$ ($\delta_{jk} = 1$, if $j = k$, $\delta_{jk} = 0$ otherwise). Numbers λ_{ij} are integers which satisfy the following conditions:

1. $\lambda_{ii} = 0$ for each i.

2. $\lambda_{ij} + \lambda_{jk} \geq \lambda_{ik}$ for all i, j, k.

An order Λ is said to be *Morita reduced* or *reduced* if it satisfies the additional condition:

3. $\lambda_{ij} + \lambda_{ji} > 0$ for each $i \neq j$. In such a case, projective modules are pairwise non-isomorphic, that is, in the decomposition of $\Lambda = P_1 \oplus P_2 \oplus \cdots \oplus P_n$ via projective modules (i.e., the rows of Λ) all indecomposable projective summands are pairwise not isomorphic, i.e., $P_i \ncong P_j$ if $i \neq j$.

In this paper, we assume that tiled orders are reduced.

We denote $\Lambda = (\lambda_{ij})_{i,\, j=1\cdots n}$, note that $\Lambda \subset T^{n\times n} = Q = \Lambda \otimes_{\mathbb{O}} T$, where Q is the rational hull of Λ, $\operatorname{Rad} Q = 0$ and Λ has a unique right

simple T-module (up to isomorphism) denoted $S_R = (T, T, ..., T) = \sum_{i=1}^{n} e_i T$, $\{e_i \mid 1 \leq i \leq n\}$ is the standard basis such that $e_i e_{jk} = \delta_{ij} e_k$. We assume the notation $S_L = (T, T, ..., T)^t$ for left modules.

The *main problem* in this case consists of describing all finitely generated torsionless Λ-modules which are called *admissible modules*.

A Λ-admissible right module (not null) is said to be *irreducible* if it is a submodule of the unique simple module (up to isomorphism). For instance, any indecomposable projective module P_i is irreducible. Thus,

$$P_i = (\pi^{\lambda_{i1}}\mathbb{O}, \pi^{\lambda_{i2}}\mathbb{O}, ..., \pi^{\lambda_{in}}\mathbb{O})$$

is a finitely generated irreducible Λ-module without $\mathbb{O}$-torsion.

Any irreducible right Λ-module A has the form

$$A = (\pi^{\alpha_1}\mathbb{O}, \pi^{\alpha_2}\mathbb{O}, ..., \pi^{\alpha_n}\mathbb{O}),$$

where $\alpha_i + \lambda_{ij} \geq \alpha_j$, $\alpha_i \in \mathbb{Z}$, $1 \leq i \leq n$. If A is a left module, then we have that $\lambda_{ij} + \alpha_j \geq \alpha_i$.

Henceforth, we denote a right (left) module A in the form $A = (\alpha_1, \alpha_2, ..., \alpha_n)$ ($(\alpha_1, \alpha_2, ..., \alpha_n)^t$, respectively).

Note that, $A \simeq A'$ if and only if $\alpha_i = \alpha_i' + k$, for some $k \in \mathbb{Z}$ and any $1 \leq i \leq n$.

If $H \subseteq (\alpha_1, \alpha_2, ..., \alpha_n)$, then we let $\mathrm{Supp}(H)$ the support of the subset H. That is,

$$H = (\alpha_i \mid \alpha_i \neq 0).$$

Irreducible right modules which are contained in a Q-simple module of a Λ order constitute a lattice denoted $\mathcal{L}_R(\Lambda)$.

We denote $\mathcal{P}(\Lambda) = \mathcal{P}_R(\Lambda)$ the subposet of $\mathcal{L}_R(\Lambda)$ which consists of irreducible projective modules the projective modules P_i are called *principals* where

$$P_i = (\lambda_{i1}, \lambda_{i2}, \ldots, \lambda_{in}) = P_i^0, \quad P_i^0 \in \mathcal{P}_R(\Lambda).$$

In this poset, there are as many projective modules as infinite chains. In such a case, modules of the form

$$P_i^k = (\lambda_{i1} + k, \ldots, \lambda_{in} + k), \quad k \in \mathbb{Z}$$

are projective modules isomorphic to P_i^0. Therefore:

$$\mathcal{P}(\Lambda) = \{P_i^k \mid 1 \leq i \leq n\},$$

where

$$P_i^k \leq P_j^l \text{ if and only if } \begin{cases} k - l \geq \lambda_{ij}, & \mathcal{P}_L(\Lambda), \\ k - l \geq \lambda_{ji}, & \mathcal{P}_R(\Lambda). \end{cases}$$

Thus, the poset $\mathcal{P}(\Lambda)$ is infinite, periodic and the sum of n chains with the form $\{P_i^k \mid 1 \leq i \leq n,\ k \in \mathbb{Z}\}$, with width $w(\mathcal{P}(\Lambda)) \leq n$. For instance, the tiled order

$$\Lambda = \begin{bmatrix} 0 & 2 & 2 \\ 2 & 0 & 2 \\ 2 & 2 & 0 \end{bmatrix} = \begin{bmatrix} \mathbb{O} & \pi^2\mathbb{O} & \pi^2\mathbb{O} \\ \pi^2\mathbb{O} & \mathbb{O} & \pi^2\mathbb{O} \\ \pi^2\mathbb{O} & \pi^2\mathbb{O} & \mathbb{O} \end{bmatrix},$$

has associated the following poset of projective modules:



Figure 1

The map $\sigma : \mathcal{P}_R(\Lambda) \to \mathcal{P}_L(\Lambda)$, given by $\sigma(P_i^k) = P_i^{-k}$ is a natural poset antiisomorphism, thus the pair $\{\mathbb{O}, \mathcal{P}(\Lambda)\}$ defines the tiled order Λ up to isomorphism, in the sense that

$$\Lambda \simeq \Lambda' \text{ if and only if } \{\mathbb{O}, \mathcal{P}(\Lambda)\} \simeq \{\mathbb{O}', \mathcal{P}(\Lambda')\}.$$

That is, $\mathbb{O} \simeq \mathbb{O}'$ and $\mathcal{P}(\Lambda) \simeq \mathcal{P}(\Lambda')$ which means that there exists a poset isomorphism $\varphi : \mathcal{P}_L(\Lambda) \to \mathcal{P}_L(\Lambda')$ such that $A \simeq B$ if and only if $\varphi(A) \simeq \varphi(B)$. Λ and Λ' are Morita-equivalents provided that φ preserves isomorphisms.

The following result characterizes isomorphic orders via matrix problems [3, 7]:

Theorem 1. *Two orders* Λ *and* Λ' *are isomorphic if the corresponding exponents matrices* λ_{ij} *and* λ'_{ij} *can be turned into each other with the help of the following admissible t-transformations*:

1. *To add an integer n to each entry of a given row i and simultaneously subtract n to each entry of the column i.*

2. *To transpose simultaneously rows i and j and columns i and j.*

The following is the finite type representation type criterion for tiled orders introduced by Hazewinkel et al. [7, 11, 12].

Theorem 2. *A tiled order* Λ *is of finite representation type if and only if* $\mathcal{P}(\Lambda) \not\supset K_1, \ldots, K_5$, *where*



Figure 2

Often, posets $K_1, \ldots, K_5$ are called the *Kleiner's critical*.

For $m \geq 1$, a $(0, 1, 2, \ldots, m)$-*tiled order* is a tiled order $\Lambda = (\lambda_{ij})$, $1 \leq i, j \leq n$, where $\lambda_{ij} \in \{0, 1, 2, \ldots, m\}$. In particular, if $\Lambda = (\lambda_{ij})$ is a $(0, m)$-tiled order, then Λ has associated a finite poset $(\mathfrak{N}, \leq) = \mathfrak{N}(\Lambda) = (\{1, 2, \ldots, n\}, \leq)$, where

$$i \leq j \text{ if and only if } \lambda_{ij} = 0,$$

for instance, the following $(0, 1)$-orders are isomorphic:

$\Lambda_3 =$

0	0	0	0	0	0	0	0
1	0	**1**	**1**	0	0	**1**	0
1	1	0	1	0	1	0	0
1	1	1	0	1	0	0	0
1	1	1	1	0	**1**	**1**	0
1	1	1	1	1	0	1	0
1	1	1	1	1	1	0	0
1	1	1	1	1	1	1	0



$\Gamma_3 =$

0	1	1	1	1	1	1	1
0	**0**	1	1	1	1	1	1
0	**1**	**0**	1	1	1	1	1
0	0	0	0	1	1	1	1
0	1	1	1	0	1	1	1
0	**0**	1	1	0	**0**	1	1
0	**1**	**0**	1	0	**1**	**0**	1
0	0	0	0	0	0	0	0



$\mathfrak{N}(\Gamma_3)$

$\Lambda_4 =$

0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	**1**	**1**	**1**	0	0	0	**1**	**1**	**1**	0	0	0	**1**	0
1	1	0	1	1	0	1	1	0	0	1	0	0	1	0	0
1	1	1	0	1	1	0	1	0	1	0	0	1	0	0	0
1	1	1	1	0	1	1	0	1	0	0	1	0	0	0	0
1	1	1	1	1	0	**1**	**1**	**1**	**1**	**1**	0	0	**1**	**1**	0
1	1	1	1	1	1	0	1	1	1	1	0	1	0	1	0
1	1	1	1	1	1	1	0	1	1	1	1	0	0	1	0
1	1	1	1	1	1	1	1	0	1	1	0	1	1	0	0
1	1	1	1	1	1	1	1	1	0	1	1	0	1	1	0
1	1	1	1	1	1	1	1	1	1	0	1	1	0	1	0
1	1	1	1	1	1	1	1	1	1	1	0	**1**	**1**	**1**	0
1	1	1	1	1	1	1	1	1	1	1	1	0	1	1	0
1	1	1	1	1	1	1	1	1	1	1	1	1	0	1	0
1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	0
1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0

$\Gamma_4 =$

0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
0	**0**	1	1	1	1	1	1	1	1	1	1	1	1	1	1
0	**1**	**0**	1	1	1	1	1	1	1	1	1	1	1	1	1
0	**0**	**0**	**0**	1	1	1	1	1	1	1	1	1	1	1	1
0	**1**	1	1	**0**	0	1	1	1	1	1	1	1	1	1	1
0	**0**	1	1	**0**	**0**	1	1	1	1	1	1	1	1	1	1
0	**1**	**0**	**1**	**0**	**1**	**0**	1	1	1	1	1	1	1	1	1
0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1
0	1	1	1	1	1	1	1	0	1	1	1	1	1	1	1
0	**0**	1	1	1	1	1	1	0	**0**	1	1	1	1	1	1
0	**1**	**0**	1	1	1	1	1	0	**1**	**0**	1	1	1	1	1
0	**0**	0	**0**	1	1	1	1	0	**0**	0	**0**	1	1	1	1
0	**1**	1	1	**0**	1	1	1	0	**1**	1	1	**0**	1	1	1
0	**0**	1	1	0	**0**	1	1	0	**0**	1	1	0	**0**	1	1
0	**1**	**0**	**1**	**0**	**1**	**0**	1	0	**1**	**0**	**1**	**0**	**1**	**0**	1
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Actually, infinite families of isomorphic $(0, 1)$-tiled orders of these types Λ_n, Γ_n can be constructed:

In such a case, if we denote $\mathcal{S}_{\Lambda_n^{\nearrow}}(\mathcal{S}_{\Gamma_n^{\swarrow}})$ the size of the support of the upper diagonal (lower diagonal). Then

$$\mathrm{Supp}(\Lambda_n^{\nearrow}) = \sum_{r=0}^{n-2} B_n^r,$$

where $\{B_n^r \mid 0 \le r \le n-2\}$ is a set-partition of $\mathrm{Supp}(\Lambda_n^{\nearrow})$ such that

1. for each $n \ge 0$, $|B_n^r|$ is a sum of consecutive positive integers $\sum_{k=0}^{T_n^r} g_n^r - k$,

2.

$$g_n^r = \begin{cases} 2^{n-1} - 1, & \text{if } r = 0, \\ h_n^r = \sum_{s=1}^{n-r-1} [c(s, n-s) - c(s, n-s-r-1)], & \text{if } r \ge 1, \\ t_{n-1} = \dfrac{(n-1)n}{2}, & \text{if } r = n-1, \end{cases}$$

3.

$$T_n^h = \begin{cases} c(h+1, n), & \text{if } 0 \le h \le n-3, \\ n-1, & \text{if } h = n-2, \end{cases}$$

where $c(k, n) = \binom{n+k}{k}$.

Remark 3. For each r and n fixed, B_n^r consists of subsets ρ_k^r with $|\rho_r^k| = g_n^r - k$ defining uniquely a projective module $P_k^0(r) \subset \Lambda_n$.

3. On the Number of Antichains in $\mathcal{P}_n$

In this section, we define a partition formula for numbers in sequence A016269. Some properties of numbers in this sequence are established as well.

Tiled orders Γ_n allow concluding the following recursion formula:

$$\mathcal{S}_{\Gamma_n^{\checkmark}} = \mathcal{A}_n^2 = 3\mathcal{A}_{n-1}^2 + t_{2^{n-1}-1},$$

where $\mathcal{A}_n^2$ denotes the number of two-point antichains in $\mathfrak{N}(\Gamma_n)$.

We prove the following result by induction:

Theorem 4. *For* $n \geq 1$, $\mathfrak{N}(\Lambda_n) \simeq \mathcal{P}_n = (2^n, \subseteq)$ *and* $\mathcal{A}_n^2 = t_{2^n} - 3^n$.

Proof. First of all, we note that projective modules $P_i^0 = (\lambda_{i1}, \lambda_{i2}, ..., \lambda_{i2^n}) \in \Lambda_n$ are uniquely determined by points $x_i \in \mathcal{P}_n$ by definition. That is, $\lambda_{ij} \in \{0, 1\}$ with

$$\lambda_{ij} = 0 \text{ if and only if } x_i \subseteq x_j \text{ in } \mathcal{P}_n.$$

Note that, in this case, $x_i \subseteq x_j$ if and only if $x_i = \{i_1, i_2, ..., i_k\}$, $x_j = \{j_1, j_2, ..., j_m\}$, $k \leq m$ and $i_h \leq j_s$ for all $1 \leq h \leq k$ and $1 \leq s \leq m$, where $\leq$ is the usual order of the natural numbers.

Matrix blocks B_n^r are obtained by fixing subsets of the form $\mathcal{P}(1, 2, ..., r) = \{i_1, i_2, ..., i_r\}$ and determining the set of incomparable points of each point $x_s \in \mathcal{H}(1, 2, ..., r) \subset \mathcal{P}_n$ with $\mathcal{P}(1, 2, ..., r) = \bigcap_{x_s \in \mathcal{H}(1, 2, ..., r)} x_s$ and $|x_i| = |x_j|$ for all $x_i, x_j \in \mathcal{H}(1, 2, ..., r)$.

Secondly, we note that $\mathcal{A}_1^2 = t_2 - 3 = 0$ and $\mathcal{A}_2^2 = t_4 - 9 = 1$. Suppose

now that the second assertion is true for all j, $1 \leq j \leq k-1$, in particular,

$$\mathcal{A}_{k-1}^2 = \sum_{i=0}^{k-3} 3^i t_{2^{k-i-2}} = A_{k-1}^2 = t_{2^{k-1}} - 3^{k-1}.$$

Therefore,

$$\mathcal{A}_k^2 = \sum_{i=0}^{k-2} 3^i t_{2^{k-i-1}} = 3(t_{2^{k-1}} - 3^{k-1}) + t_{2^{k-1}-1},$$

then

$$\mathcal{A}_k^2 = 3\left(\frac{2^{k-1}(2^{k-1}+1)}{2}\right) + \left(\frac{(2^{k-1}-1)(2^{k-1})}{2}\right) - 3^k$$

$$= \frac{2^{k-1}}{2}(32^{k-1} + 3 + 2^{k-1} - 1) - 3^k.$$

Thus,

$$A_k^2 = \frac{2^{k-1}}{2}[42^{k-1} + 2] - 3^k = 2^{k-1}[2^k + 1] - 3^k = t_{2^k} - 3^k. \qquad \square$$

Isomorphisms between tiled orders Λ_n and Γ_n allow obtaining the following formula partition of A_n^2 for each $n \geq 3$:

Lemma 5. $\mathcal{A}_n^2 = \sum_{r=0}^{n-3} [t_{g_n^r} - t_{g_n^r - T_n^r}]$.

For instance:

$g_3^0 = 3$, $h_3^0 = c(1, 2) = 3$, *thus*:

$\mathcal{A}_3^2 = t_{g_3^0} + t_{g_3^0 - h_3^0} + t_2 = 6 + 3 = 9$,

$g_4^0 = 7$, $g_4^1 = 7$,

$h_4^0 = c(1, 3) = 4,\ h_4^1 = c(2, 2) = 6,$ *thus*

$$\mathcal{A}_4^2 = (t_{g_4^0} - t_{g_4^0-4}) + (t_{g_4^1} - t_{g_4^1-6}) + t_3 = (28 - 6) + (28 - 1) + 6 = 55.$$

Sequence $\mathcal{A}_n^2$ (A016269 in OEIS) grows rapidly, as an example, the following are the first ten terms of this sequence:

$$\{0, 1, 9, 55, 285, 1351, 6069, 26335, 111645, 465751\},$$

$$\mathcal{A}_{50}^2 = 633824582216127571845716253751,$$

$$\mathcal{A}_{100}^2 = 803469022128979760250249034840178665431449990319442661731375. \tag{1}$$

The following results are easy to see via Theorem 4:

Corollary 6. *For* $n \geq 1$, *sequence* $\mathcal{A}_n^2$ *satisfies the following conditions*:

1. *If* $n \equiv 0 \bmod 4$, *then* $5 \mid \mathcal{A}_2^n$ *and* $\mathcal{A}_n^2 \equiv 1 \bmod 3$.

2. *If* $n \equiv 1 \bmod 4$, *then* $3 \mid \mathcal{A}_2^n$ *and* $5 \mid \mathcal{A}_n^2$.

3. *If* $n \equiv 2 \bmod 4$, *then* $\mathcal{A}_n^2 \equiv 3 \bmod 4$, $\mathcal{A}_n^2 \equiv 1 \bmod 3$ *and* $\mathcal{A}_n^2 \equiv 1 \bmod 10$.

4. *If* $n \equiv 3 \bmod 4$, *then* $3 \mid \mathcal{A}_2^n$ *and* $\mathcal{A}_n^2 \equiv 9 \bmod 10$.

Theorem 7. *The sequence* $\mathcal{A}_n^2$ *has no prime numbers*.

Proof. According to the last corollary, it suffices to consider the case $n \equiv 2 \bmod 4$. In such a case, if $\mathcal{A}_n^2$ is a prime number p, then $\left(\frac{-3}{p}\right) = 1$ and $\left(\frac{-1}{p}\right) = -1$ which is a contradiction. □

Remark 8. Corollary 6 and Theorem 7 are answers to the questions posed by the first author et al. in [4] regarding the number of two-point antichains in $\mathcal{P}_n$. Worth noting that formula proposed in Theorem 4 was suggested by Riviere in [8].

Acknowledgment

The authors are indebted to the anonymous referee for his(her) useful comments and suggestions.

References

[1] A. M. Cañadas, H. Giraldo and P. F. F. Espinosa, Categorification of some integer sequences, Far East J. Math. Sci. (FJMS) 92(2) (2014), 125-139.

[2] A. M. Cañadas, H. Giraldo and V. C. Vargas, Categorification of some integer sequences and higher dimensional partitions, Far East J. Math. Sci. (FJMS) 93(2) (2014), 133-149.

[3] A. M. Cañadas, R.-J. Serna and C.-I. Espinosa, On the reduction of some tiled orders, JP J. Algebra Number Theory Appl. 36(2) (2015), 157-176.

[4] A. M. Cañadas, H. Giraldo and G. B. Rios, An algebraic approach to the number of some antichains in the powerset 2^n, JP J. Algebra Number Theory Appl. 38(1) (2016), 45-62.

[5] P. Fahr and C. M. Ringel, A partition formula for Fibonacci numbers, J. Integer Seq. 11(1) (2008), Article 08.1.4, 9 pp.

[6] P. Fahr and C. M. Ringel, Categorification of the Fibonacci numbers using representations of quivers, J. Integer Seq. 15(2) (2012), Article 12.2.1, 12 pp. arXiv1107.1858v2.

[7] M. Hazewinkel, N. Gubareni and V. V. Kirichenko, Algebras, Rings and Modules, Vol. 2, First Edition, Springer, 2007.

[8] N. M. Riviere, Recursive formulas on free distributive lattices, J. Combin. Theory 5 (1968), 229-234.

[9] D. Simson, Linear Representations of Partially Ordered Sets and Vector Space Categories, Gordon and Breach, London, 1992.

[10] N. J. A. Sloane, On-line Encyclopedia of Integer Sequences, The OEIS Foundation, http://oeis.org/A052558,A132262,A110122.

[11] A. G. Zavadskij and V. V. Kiričenko, Semimaximal rings of finite type, Mat. Sb. 103 (1977), 323-345 (in Russian).

[12] A. G. Zavadskij and V. V. Kirichenko, Torsion-free modules over primary rings, Journal of Soviet Mathematics 11(4) (1979), 598-612.