Relativistic Statistical Mechanics

Juan A. Guarín-Rojas
María Fernanda Carvajal
Carlos José Páez González
Antonio Calixto Gutiérrez Piñerez

July 29, 2024

Abstract Hola

1 Transformation Laws Table

The table 1 shows the results achieved for some authors [1–5] and the results of the authors of this document for the transformation laws of the thermodynamic variables. It shows the agreement of this work with [5] and the partial agreement with [3, 4].

	Einstein, Planck	**Ott, Arzelièz**	**Sutcliffe**	**Guarín-Carvajal-Páez-Gutiérrez**
Pressure	$P = P_0$		$P_m = P_0$, $P_{th} = P_0\gamma^2$	$P = P_0\gamma^2$
Volume	$V = V_0/\gamma$			
Entropy	$S = S_0$			
Temperature	$T = T_0/\gamma$	$T = T_0\gamma$		
Heat	$\delta Q = \delta Q_0/\gamma$	$\delta Q = \delta Q_0\gamma$		
Internal energy	$E = E_0\gamma + P_0V_0\gamma v^2/c^2$	$E = E_0\gamma$		
Helmholtz free energy	$F = F_0/\gamma$	$F = F_0\gamma$		

Table 1: Transformation of the thermodynamic quantities for different authors, adapted from [6]. A zero subindex indicates variables measured for a rest observer with the container of the gas. By convention $\gamma = 1/\sqrt{1 - v^2/c^2}$ is the Lorentz factor, P_m is the *mechanical pressure* and P_{th} the *thermodynamic* one (as discussed by [5]).

2 The invariance of entropy

Theorem 1. The entropy $S = k_B \ln \Omega$ for a moving inertial observer K with respect to the container of an ideal gas (N equal particles of same mass, elastic collisions between particles and

walls, and non-interacting particles) is equal to the entropy $S_0 = k_B \ln \Omega_0$ for a rest observer K_0 with the container.

Here, $\Omega = \Omega(E, V, N)$ is the number of states for K depending on the internal energy, the volume, and the number of particles respectively. Analogously, $\Omega_0 = \Omega_0(E_0, V_0, N_0)$ is the number of states for K_0 depending on its corresponding internal energy, volume and number of particles respectively. Besides, the number of particles for each frame must be the same, this is $N = N_0$.

Proof:
By definition

$$\Omega = \frac{1}{N!}\frac{1}{h^{3N}} \underbrace{\int_{\mathscr{V}} d^3r_1 \cdots d^3r_N}_{\Omega_v} \underbrace{\int_{\mathcal{R}} d^3p_1 \cdots d^3p_N}_{\Omega_p} , \tag{1}$$

where h is the planck constant, $\mathbf{r}_i$ and $\mathbf{p}_i$ are the 3−position and 3−momentum of the ith particle for K, $\mathscr{V}$ is the volume space, and $\mathcal{R}$ is the momentum space defined as

$$\mathcal{R} = \left\{ (\vec{p}_1, \ldots, \vec{p}_N) \in \mathbb{R}^{4N} \mid \vec{p}_i \cdot \vec{p}_i = -m^2 \text{ and } \sum_{i=1}^{N} \vec{p}_i = \vec{\mathscr{P}} \right\} , \tag{2}$$

where $\vec{p}_i$ is the 4−momentum of the ith particle, and $\vec{\mathscr{P}}$ is the total 4−momentum of the system. $\vec{\mathscr{P}}$ is such that for K_0 its components are $(E_0/c^2, \mathbf{0})$. Here, 4−vectors $\vec{u}$ such that $\vec{u} \cdot \vec{u} < 0$ are timelike ones, and $m\vec{u}$ is the momentum of a particle with mass m and 4−velocity $\vec{u}$, which satisfies $\vec{u} \cdot \vec{u} = -1$. These are the conventions followed in [7]. Now, by the non-interacting condition, the position $\mathbf{r}_i$ of each particle is independent one of the other. And therefore we can separate Ω_v as a product of integrals

$$\Omega_v = \left(\int_V d^3r_1 \right) \cdots \left(\int_V d^3r_N \right) = \left(\int_V d^3r_1 \right)^N = V^N . \tag{3}$$

Lemma 1. The volume V_0 for the observer K_0 changes to $V_0/\gamma = V$ by a Lorentz transformation corresponding to a Lorentz Boost in one direction. γ is the Lorentz factor.

By the previous lemma

$$\Omega_v = \frac{V^N}{\gamma^N} = \frac{\Omega_{v_0}}{\gamma^N} . \tag{4}$$

By the other side, chosen an appropriate coordinate axis we can say that the Lorentz Boost occur in the x−axis. According to [8] the transformation laws for momentum in this case are

$$p_{x_i} = \gamma \left(g_{x_i} + \frac{v}{c^2} E_{0_i} \right) , \tag{5}$$

$$p_{y_i} = g_{y_i} , \tag{6}$$

$$p_{z_i} = g_{z_i} , \tag{7}$$

where $\mathbf{p}_i$ is the momentum of the ith particle for observer K, and $\mathbf{g}_i$ is the momentum of the same particle but for K_0. Besides, v is the relative speed between the observers, $\gamma = 1/\sqrt{1 - v^2/c^2}$, c

is the speed of light, and $E_{0_i} = c\sqrt{m^2c^2 + {g_i}^2} = mc^2\sqrt{1 + \frac{{g_i}^2}{m^2c^2}}$ is the relativistic energy of the particle for K_0.

We want to implement a change of variables from p_i to g_i in the momentum integrals. The Jacobian of the transformation is

$$J(\mathbf{g}) = \frac{\partial p_{x^i}}{\partial g_{x^i}} = \begin{vmatrix} \frac{\partial p_x}{\partial g_x} & \frac{\partial p_x}{\partial g_y} & \frac{\partial p_x}{\partial g_z} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = \frac{\partial p_x}{\partial g_x} = \frac{\partial}{\partial g_x}\left[\gamma\left(g_x + \frac{v}{c^2}mc^2\sqrt{1 + \frac{g^2}{m^2c^2}}\right)\right] . \tag{8}$$

Knowing that $g^2 = {g_x}^2 + {g_y}^2 + {g_z}^2$,

$$J(\mathbf{g}) = \gamma\left(1 + \frac{v}{c^2}mc^2\frac{g_x/(m^2c^2)}{\sqrt{1 + \frac{g^2}{m^2c^2}}}\right) = \gamma\left(1 + \frac{v}{c}\frac{g_x/(mc)}{\sqrt{1 + \frac{g^2}{m^2c^2}}}\right) . \tag{9}$$

Then by implement this coordinate transformation for all the variables of the N particles one gets

$$\Omega_p = \int_{\mathcal{R}} d^3p_1 \cdots d^3p_N = \int_{\mathcal{R}} \gamma^N \prod_{i=1}^{N}\left(1 + \frac{v}{c}\frac{g_{x_i}/(mc)}{\sqrt{1 + \frac{{g_i}^2}{m^2c^2}}}\right) d^3g_1 \cdots d^3g_N \, . \tag{10}$$

On the other side, one can check that

$$\begin{aligned} \prod_{i=1}^{N}(1 + M_i) &= 1 + M_1 + M_2 + \cdots + M_N + \\ &\quad + M_1M_2 + M_1M_3 + \cdots + M_1M_N + \cdots M_{N-1}M_n + \\ &\quad + M_1M_2M_3 + \cdots M_1M_{N-1}M_N + \cdots M_{N-2}M_{N-1}M_N + \cdots , \\ &= 1 + \sum_{k=1}^{N}\sum_{1\le i_1<i_2<\cdots<i_k\le N} M_{i_1}\cdots M_{i_k} , \end{aligned} \tag{11}$$

where $\sum_{1\le i_1<i_2<\cdots<i_k\le N}$ stands for the sum over all integers $i_1, \ldots, i_k$ that satisfies $1 \le i_1 < i_2 < \cdots < i_k \le N$. Therefore

$$\Omega_p = \int_{\mathcal{R}} \gamma^N \left(1 + \sum_{k=1}^{N}\sum_{1\le i_1<i_2<\cdots<i_k\le N} \frac{v^k}{c^k}\frac{g_{x_{i_1}}/(mc)}{\sqrt{1 + \frac{{g_{i_j}}^2}{m^2c^2}}}\cdots\frac{g_{x_{i_k}}/(mc)}{\sqrt{1 + \frac{{g_{i_k}}^2}{m^2c^2}}}\right) d^3g_1 \cdots d^3g_N \, . \tag{12}$$

Now, as integration is a linear operation,

$$\Omega_p = \gamma^N \int_{\mathcal{R}} d^3g_1 \cdots d^3g_N + \gamma^N \sum_{k=1}^{N}\sum_{1\le i_1<i_2<\cdots<i_k\le N} \frac{v^k}{c^k}\int_{\mathcal{R}} \frac{g_{x_{i_1}}/(mc)}{\sqrt{1 + \frac{{g_{i_j}}^2}{m^2c^2}}}\cdots\frac{g_{x_{i_k}}/(mc)}{\sqrt{1 + \frac{{g_{i_k}}^2}{m^2c^2}}} d^3g_1 \cdots d^3g_N \, . \tag{13}$$

While integrating the second term we will find that if a point $(\mathbf{g}_1, \ldots, \mathbf{g}_k)$ is in $\mathcal{R}$, then $(-\mathbf{g}_1, \ldots, -\mathbf{g}_k)$ also will be in $\mathcal{R}$. This can be seen by viewing the momentum space $\mathcal{R}$ in the K_0 local coordinates

$$\mathcal{R} = \left\{ (\vec{p}_1, \ldots, \vec{p}_N) \in \mathbb{R}^{4N} \;\middle|\; \vec{p}_i \cdot \vec{p}_i = -m^2 \ , \ \sum_{i=1}^{N} c\sqrt{m^2c^2 + |\mathbf{g}_i|^2} = E_0 \ , \text{ and } \sum_{i=1}^{N} \mathbf{g}_i = 0 \right\} . \tag{14}$$

This shows the previous statement. And so $\mathcal{R}$ is composed by pairs of points $(\mathbf{g}_1, \ldots, \mathbf{g}_N)$ and $(-\mathbf{g}_1, \ldots, -\mathbf{g}_N)$ such that its respective contribution to the integral cancel out each other, this is

$$\int_{\mathcal{R}} \frac{g_{x_{i_1}}/(mc)}{\sqrt{1+\frac{g_{i_i}{}^2}{m^2c^2}}} \cdots \frac{g_{x_{i_k}}/(mc)}{\sqrt{1+\frac{g_{i_k}{}^2}{m^2c^2}}} d^3g_1 \cdots d^3g_N = 0\,. \tag{15}$$

Then only the first term in 13 remains

$$\Omega_p = \gamma^N \int_{\mathcal{R}} d^3g_1 \cdots d^3g_N\,. \tag{16}$$

This integral is exactly the definition of Ω_p but for K_0. Hence $\Omega_p = \gamma^N \Omega_{p_0}$, and so $\Omega = \Omega_0$. This ends the proof.

3 Appendix

In local coordinates the components of the 4−velocity $\vec{u}_i$ of a particle are $(\gamma(v_i), \gamma(v_i)\,\mathbf{v}_i/c)$, where $\mathbf{v}_i$ is the 3−velocity of the particle seen by the local frame and $\gamma(v_i) = 1/\sqrt{1 - v_i{}^2/c^2}$. Besides, the components of its 4−momentum $\vec{p}_i = m\vec{u}_i$ are $(E_i/c^2, \mathbf{p}_i/c)$, where E_i is the relativistic energy of the particle seen by the local frame and $\mathbf{p}_i$ is the corresponding 3−momentum seen by the local frame.

References

[1] A. Einstein. On the relativity principle and the conclusions drawn from it [uber das relativitat-sprinzip und die aus demselben gezogenen folgerungen]. *Jahrb. d. Radioaktivitat u. Elektronik*, 4:411–462, 1907.

[2] Max Planck. On the dynamics of moving systems [zur dynamik bewegter systeme]. *Annalen der Physik*, 331(6):1–34, 1908.

[3] H. Ott. Lorentz transformation of heat and temperature [lorentz-transformation der wärme und der temperatur]. *Zeitschrift für Physik*, 175:70–104, 1963.

[4] Henri Arzeliès. Transformation relativiste de la temperature et de quelques autres grandeurs thermodynamiques. *Il Nuovo Cimento (1955-1965)*, 35:792–804, 1965.

[5] W. G. Sutcliffe. Lorentz transformations of thermodynamic quantities. *Il Nuovo Cimento*, 39:683–686, 1965.

[6] R. Balescu. Relativistic statistical thermodynamics. *Physica*, 40(3):309–338, 1968.

[7] É. Gourgoulhon. *Special Relativity in General Frames: From Particles to Astrophysics*. Graduate Texts in Physics. Springer Berlin Heidelberg, 2013.

[8] L.D. Landau and E.M. Lifshitz. Chapter 2 - relativistic mechanics. In L.D. Landau and E.M. Lifshitz, editors, *The Classical Theory of Fields (Fourth Edition)*, volume 2 of *Course of Theoretical Physics*, pages 24–42. Pergamon, Amsterdam, fourth edition edition, 1975.