EVOCURE Candidate Screening

Pancreatic ductal adenocarcinoma, five prioritised compounds

EVOCURE-PDAC-20261027 | generated 2026-10-27 19:53:34 UTC

What this document is

A demonstration of an open source in silico screening platform. It shows what the software computed on one reproducible run: a bounded evolutionary search over molecular structures, scored by a QSAR model, toxicology heuristics and a simulated two dimensional tumour microenvironment.

What it is not: evidence that any compound here treats pancreatic cancer. No molecule in this report has been synthesised, assayed in a cell, or given to an animal or a person. These are computational hypotheses at the earliest possible stage of discovery, and the large majority of such hypotheses fail.

Result in one line

1 of 5 candidates separated from the negative control at an uncorrected p < 0.05, and none survived correction for 5 comparisons.

The screening pipeline ran end to end and produced five ranked candidates. The tissue assay was checked with a positive and a negative control before the candidates were read, so the result above is interpretable rather than merely generated. The positive control separates from the negative control, so the tissue model responds to a known active in this configuration.

Run at a glance

Structures evaluated	106 unique valid structures
Generations	4, population 24
Replicate seeds per compound	5
Compute backend	NumPy CPU float64
QSAR training set	43 curated compounds
QSAR scaffold split R2	0.323
QSAR classifier ROC AUC	0.917
Wall clock	64 seconds

Open source platform and data. Reproduce this document with the commands in the reproducibility section. Report generated directly from the run's results file; no figure in it was entered by hand.

1. Summary

A bounded evolutionary search started from four approved pancreatic cancer drugs and generated structural variants through point mutations and scaffold recombination. Every variant was validated with RDKit, scored on five objectives, and the non dominated front was carried into the next generation. The five compounds below are the highest ranked variants that were not themselves seeds.

1.1 Assay validity, checked before reading the candidates

Role	Compound	TGI %	95% CI	CI excludes 0
Positive control	Gemcitabine	67.1	60.4 to 73.8	yes
Negative control	Caffeine	-5.7	-18.3 to 6.8	no

Does the simulated assay work?
Positive versus negative control: $p < 0.001$, mean difference 72.9 percentage points, Hedges g 8.12.

The positive control separates from the negative control, so the tissue model responds to a known active in this configuration.

1.2 The five candidates

ID	Compound	pIC50	Score	TGI %	95% CI	vs control
Candidate 01	Compound-80A085	6.84	0.635	9.9	-6.1 to 25.8	not distinguishable
Candidate 02	Compound-CFB9B6	7.03	0.622	9.4	-6.8 to 25.5	not distinguishable
Candidate 03	Compound-FAFE93	5.97	0.612	1.6	-8.9 to 12.2	not distinguishable
Candidate 04	Compound-ABD9EC	6.48	0.610	10.7	-6.5 to 27.9	not distinguishable
Candidate 05	Compound-7AA5E8	5.74	0.607	9.1	0.7 to 17.5	marginal, uncorrected

TGI is tumour growth inhibition in the simulated microenvironment, relative to an untreated arm, averaged over independent seeds. Score is the weighted composite across the five search objectives. The last column reports Welch's t-test of each candidate against the negative control; it is the column that decides whether a point estimate means anything.

Scope of the positive finding
Separated at an uncorrected $p < 0.05$: Candidate 05. None of them survives Bonferroni correction for 5 comparisons ($p < 0.01$). Testing five candidates against one control arm makes a single result near $p = 0.05$ roughly what chance alone would produce, so this is a marginal signal and should be treated as one.

Either way this is a result inside a simulation whose parameters are not experimentally calibrated. It justifies further computational scrutiny and nothing beyond that. Separation here is not potency, not selectivity, not safety and not evidence about human disease.

2. Method

2.1 Search

Objective prompt	Find 5 experimental candidates for pancreatic cancer based on Gemcitabine, 5-Fluorouracil, Erlotinib and Paclitaxel. Prioritize activity, low predicted toxicity and drug-likeness.
Planner	local_rule_based_planner
Derived priorities	safety, druglikeness, activity
Seed compounds	Gemcitabine, 5-Fluorouracil, Paclitaxel, Erlotinib
Generations	4
Population size	24
Applied concentration	5.0 uM
Exposure duration	24.0 hours
Search seed	20261027

The planner converts a plain language objective into search priorities only. It does not propose molecules and it does not supply any activity, potency or safety number. All quantitative output comes from the deterministic scientific models.

2.2 Objective weights

Objective	Weight	Source of the value
Predicted PDAC activity	2.0	Calibrated QSAR regressor, sigmoidal utility at pIC50 6.0
Low toxicological risk	2.0	Ten category structural alert and rule based risk index
Drug likeness (QED)	2.0	RDKit quantitative estimate of drug likeness
Selectivity index	1.0	ADMET derived tumour versus healthy cell window
Applicability domain	1.0	Maximum Tanimoto similarity to the training set

Tumour cell kill is never maximised alone. A compound that scores well on activity but triggers toxicity alerts or falls outside the applicability domain is penalised by construction.

2.3 The QSAR model, and what it can carry

Training compounds	43
Validation	Bemis-Murcko scaffold split, no scaffold leakage
Regression RMSE	1.704 log units
Regression R2	0.323
Classifier ROC AUC	0.917
Classifier Brier score	0.163
Conformal 95% half width	4.16 log units

The potency numbers in this report are weak, by the model's own measurement

The pIC50 regressor explains little of the variance in the held out scaffolds (R2 = 0.32), and its calibrated 95% conformal interval is about 4.2 log units wide in each direction. An interval that wide spans roughly eight orders of magnitude in IC50, which is to say it does not constrain potency at all.


The active versus inactive classifier is the stronger half of the model (ROC AUC = 0.92). Reading pIC50 as a rough ordering signal is defensible. Reading any single predicted IC50 as a quantity is not. Every pIC50 in this report is printed with its conformal interval for that reason.

2.4 Tissue simulation

Model	Two dimensional agent based PDAC microenvironment
Grid	36 x 28 cells over 800 x 600 micrometres
Fields	Oxygen and nutrient reaction diffusion, drug diffusion and uptake
Features	Hypovascularity, dense collagen matrix, sensitive and resistant clones
Comparison	Treated arm against an untreated arm with the same seed
Replicates	5 independent seeds per compound

Kinetics and microenvironment parameters are exploratory and have not been fitted to experimental pancreatic tumour data. The model's drug response is driven by the predicted IC50, so two compounds with the same predicted IC50 produce the same trajectory. Simulated growth inhibition is a property of this model, not a measurement.

Candidate 01: Compound-80A085



Identity

Canonical SMILES	Nc1ccn([C@H]2O[C@@H](n3ccc(N)nc3=O)C(F)(F)[C@H]2C(=O)O)c(=O)n1
InChIKey	HKIZSQLKHYHGKJ-KHQFGBGNSA-N
Molecular formula	C13H12F2N6O5
Murcko scaffold	O=c1ncccn1C1CCC(n2cccnc2=O)O1
Origin	generation 4 variant, directed_optimization

Physicochemical profile

MW	cLogP	TPSA	HBD	HBA	QED	Lipinski
370.3	-0.97	168.3	3	8	0.615	passes

Predicted activity, with its uncertainty

Quantity	Value	95% interval	Reading
Predicted pIC50	6.845	2.68 to 11.01	ordering signal only
Implied IC50 (uM)	0.1428	spans orders of magnitude	not a quantity
Active probability	0.803	0.70 to 0.90	the stronger prediction
Applicability domain	0.62	Tanimoto, nearest neighbour	inside domain

Objective scores and risk

Activity	Selectivity	QED	Low tox	Domain	Composite	Risk index
0.700	0.000	0.615	0.916	0.620	0.635	8.4 / 100

Simulated tissue response

Metric	Value
Tumour growth inhibition, mean	9.87 % over 5 seeds
95% confidence interval	-6.06 % to 25.80 %
Interval excludes zero	no
Per seed values	17.1, -8.1, 26.2, 6.8, 7.3
Resistant clone fraction	0.336
Versus negative control	p = 0.067, Hedges g 1.22
Distinguishable, uncorrected	no
Distinguishable, Bonferroni corrected	no

Relation to known drugs


Nearest reference	Gemcitabine (Tanimoto 0.620)
Exact structural match	none in the local library
Library scope	28 curated reference drugs

The reference library is small and local. Absence of a match here does not establish that a structure is novel, and similarity to an active drug does not establish that it is active. A patent and literature search would be the next step before treating any of these as new chemical matter.

Verdict

This candidate did not separate from the negative control. Its position in the ranking reflects predicted properties and the absence of alerts, not a demonstrated simulated effect.

Candidate 02: Compound-CFB9B6



Identity

Canonical SMILES	Nc1ccn([C@@H]2O[C@H](Cc3c[nH]c(=O)[nH]c3=O)[C@@H](O)C2(F)F)c(=O)n1
InChIKey	SYMMKNWOSBOWTM-GTNGPMTGSA-N
Molecular formula	C13H13F2N5O5
Murcko scaffold	O=c1[nH]cc(CC2CCC(n3cccnc3=O)O2)c(=O)[nH]1
Origin	generation 4 variant, directed_optimization

Physicochemical profile

MW	cLogP	TPSA	HBD	HBA	QED	Lipinski
357.3	-1.66	156.1	4	7	0.512	passes

Predicted activity, with its uncertainty

Quantity	Value	95% interval	Reading
Predicted pIC50	7.030	2.87 to 11.19	ordering signal only
Implied IC50 (uM)	0.0933	spans orders of magnitude	not a quantity
Active probability	0.835	0.73 to 0.93	the stronger prediction
Applicability domain	0.64	Tanimoto, nearest neighbour	inside domain

Objective scores and risk

Activity	Selectivity	QED	Low tox	Domain	Composite	Risk index
0.737	0.000	0.512	0.916	0.643	0.622	8.4 / 100

Simulated tissue response

Metric	Value
Tumour growth inhibition, mean	9.38 % over 5 seeds
95% confidence interval	-6.75 % to 25.52 %
Interval excludes zero	no
Per seed values	17.1, -8.1, 26.2, 6.8, 4.9
Resistant clone fraction	0.345
Versus negative control	p = 0.076, Hedges g 1.17
Distinguishable, uncorrected	no
Distinguishable, Bonferroni corrected	no

Relation to known drugs


Nearest reference	Gemcitabine (Tanimoto 0.643)
Exact structural match	none in the local library
Library scope	28 curated reference drugs

The reference library is small and local. Absence of a match here does not establish that a structure is novel, and similarity to an active drug does not establish that it is active. A patent and literature search would be the next step before treating any of these as new chemical matter.

Verdict

This candidate did not separate from the negative control. Its position in the ranking reflects predicted properties and the absence of alerts, not a demonstrated simulated effect.

Candidate 03: Compound-FAFE93



Identity

Canonical SMILES	Nc1ccn([C@@H]2O[C@H](CO)[C@H](C(=O)O)C2(F)F)c(=O)n1
InChIKey	HKQIZRDZFOYESW-UEYSZJFGSA-N
Molecular formula	C10H11F2N3O5
Murcko scaffold	O=c1ncccn1C1CCCO1
Origin	generation 4 variant, directed_optimization

Physicochemical profile

MW	cLogP	TPSA	HBD	HBA	QED	Lipinski
291.2	-0.95	127.7	3	6	0.655	passes

Predicted activity, with its uncertainty

Quantity	Value	95% interval	Reading
Predicted pIC50	5.971	1.81 to 10.13	ordering signal only
Implied IC50 (uM)	1.0685	spans orders of magnitude	not a quantity
Active probability	0.740	0.64 to 0.84	the stronger prediction
Applicability domain	0.77	Tanimoto, nearest neighbour	inside domain

Objective scores and risk

Activity	Selectivity	QED	Low tox	Domain	Composite	Risk index
0.493	0.000	0.655	0.916	0.766	0.612	8.4 / 100

Simulated tissue response

Metric	Value
Tumour growth inhibition, mean	1.63 % over 5 seeds
95% confidence interval	-8.93 % to 12.19 %
Interval excludes zero	no
Per seed values	0.0, -5.4, 9.5, 11.4, -7.3
Resistant clone fraction	0.312
Versus negative control	p = 0.248, Hedges g 0.71
Distinguishable, uncorrected	no
Distinguishable, Bonferroni corrected	no

Relation to known drugs


Nearest reference	Gemcitabine (Tanimoto 0.766)
Exact structural match	none in the local library
Library scope	28 curated reference drugs

The reference library is small and local. Absence of a match here does not establish that a structure is novel, and similarity to an active drug does not establish that it is active. A patent and literature search would be the next step before treating any of these as new chemical matter.

Verdict

This candidate did not separate from the negative control. Its position in the ranking reflects predicted properties and the absence of alerts, not a demonstrated simulated effect.

Candidate 04: Compound-ABD9EC



Identity

Canonical SMILES	C#Cc1cccc(C2(F)[C@H](O)[C@@H](CO)S[C@H]2n2ccc(N)nc2=O)c1
InChIKey	VZMQZTGQSOCMBE-UBEPTHPKSA-N
Molecular formula	C17H16FN3O3S
Murcko scaffold	O=c1ncccn1C1SCCC1c1ccccc1
Origin	generation 4 variant, directed_optimization

Physicochemical profile

MW	cLogP	TPSA	HBD	HBA	QED	Lipinski
361.4	0.64	101.4	3	6	0.694	passes

Predicted activity, with its uncertainty

Quantity	Value	95% interval	Reading
Predicted pIC50	6.482	2.32 to 10.64	ordering signal only
Implied IC50 (uM)	0.3293	spans orders of magnitude	not a quantity
Active probability	0.835	0.73 to 0.93	the stronger prediction
Applicability domain	0.42	Tanimoto, nearest neighbour	inside domain

Objective scores and risk

Activity	Selectivity	QED	Low tox	Domain	Composite	Risk index
0.618	0.000	0.694	0.916	0.424	0.610	8.4 / 100

Simulated tissue response

Metric	Value
Tumour growth inhibition, mean	10.70 % over 5 seeds
95% confidence interval	-6.47 % to 27.86 %
Interval excludes zero	no
Per seed values	17.1, -10.8, 26.2, 13.6, 7.3
Resistant clone fraction	0.345
Versus negative control	p = 0.067, Hedges g 1.23
Distinguishable, uncorrected	no
Distinguishable, Bonferroni corrected	no

Relation to known drugs


Nearest reference	Gemcitabine (Tanimoto 0.424)
Exact structural match	none in the local library
Library scope	28 curated reference drugs

The reference library is small and local. Absence of a match here does not establish that a structure is novel, and similarity to an active drug does not establish that it is active. A patent and literature search would be the next step before treating any of these as new chemical matter.

Verdict

This candidate did not separate from the negative control. Its position in the ranking reflects predicted properties and the absence of alerts, not a demonstrated simulated effect.

Candidate 05: Compound-7AA5E8



Identity

Canonical SMILES	C#Cc1cccc([C@H]2O[C@@H](n3ccc(N)nc3=O)C(F)(F)[C@@H]2O)c1
InChIKey	UZLJPISFEFBHPV-MGPQQGTHSA-N
Molecular formula	C16H13F2N3O3
Murcko scaffold	O=c1ncccn1C1CCC(c2ccccc2)O1
Origin	generation 4 variant, directed_optimization

Physicochemical profile

MW	cLogP	TPSA	HBD	HBA	QED	Lipinski
333.3	1.07	90.4	2	5	0.803	passes

Predicted activity, with its uncertainty

Quantity	Value	95% interval	Reading
Predicted pIC50	5.735	1.57 to 9.90	ordering signal only
Implied IC50 (uM)	1.8429	spans orders of magnitude	not a quantity
Active probability	0.848	0.75 to 0.95	the stronger prediction
Applicability domain	0.55	Tanimoto, nearest neighbour	inside domain

Objective scores and risk

Activity	Selectivity	QED	Low tox	Domain	Composite	Risk index
0.434	0.000	0.803	0.916	0.552	0.607	8.4 / 100

Simulated tissue response

Metric	Value
Tumour growth inhibition, mean	9.11 % over 5 seeds
95% confidence interval	0.72 % to 17.51 %
Interval excludes zero	yes
Per seed values	17.1, 0.0, 14.3, 6.8, 7.3
Resistant clone fraction	0.307
Versus negative control	p = 0.029, Hedges g 1.56
Distinguishable, uncorrected	yes
Distinguishable, Bonferroni corrected	no

Relation to known drugs

Nearest reference	Gemcitabine (Tanimoto 0.552)
Exact structural match	none in the local library
Library scope	28 curated reference drugs

The reference library is small and local. Absence of a match here does not establish that a structure is novel, and similarity to an active drug does not establish that it is active. A patent and literature search would be the next step before treating any of these as new chemical matter.

Verdict

This candidate separated from the negative control at an uncorrected threshold but not after correcting for the number of candidates tested. Across five candidates, one result in this range is close to what chance produces. Treat it as a lead to re-run with more seeds, not as a demonstrated effect.

3. What this run does not show

The value of a screening report lies as much in its boundaries as in its ranking. The following are not minor caveats; they determine what the numbers above can be used for.

No experimental evidence of any kind
No compound here has been synthesised. There is no cell assay, no animal model and no clinical data. Every number is the output of a model.

The tissue model is not calibrated to real tumours
Kinetics, diffusion constants, clone behaviour and drug response parameters are plausible but have not been fitted to experimental pancreatic tumour data. Simulated growth inhibition is a property of this software.

The potency model cannot resolve potency
A scaffold split R2 of 0.32 and a conformal interval of about 4.2 log units mean the regressor orders compounds loosely and quantifies none of them.

A small, narrow training set
43 curated compounds. Applicability domain is measured against that set, so high domain confidence means close to the training data, which also means structurally unadventurous.

Virtual time is not real time
Generations and exposure hours index the simulation. They do not correspond to chemical synthesis, biological timescales or clinical development.

Structural similarity is not efficacy and not novelty
Matching a known drug does not transfer its activity. Failing to match anything in a small local library does not make a structure new.

No ADME, formulation or synthesis assessment
Nothing here addresses whether these molecules can be made at reasonable cost, are stable, are orally available, or reach a pancreatic tumour in a living system. Dense desmoplastic stroma makes that last point a central problem in this disease.

Multiple comparisons
Five candidates were tested against one control arm. The per candidate threshold is uncorrected; a Bonferroni corrected threshold would be $p < 0.01$, and the results file records both.

If you take one thing from this report
Pancreatic ductal adenocarcinoma has a five year survival rate in the low teens, and that is precisely why early computational output about it should be described carefully. This platform is a hypothesis generator. The appropriate next step for any compound listed here is scrutiny by a medicinal chemist, not optimism.

4. Reproducibility

4.1 Environment of this run

Platform	Windows-11-10.0.26200-SP0
Python	3.12.13
Compute backend	NumPy CPU float64
Device	host CPU
Field precision	float64
Search seed	20261027
Replicate seeds	20261027, 20261028, 20261029, 20261030, 20261031
Results hash	e4f7842c77e8a49a39c2e38c3896d7621df48240c59f3292

4.2 Library versions

Package	Version
numpy	2.2.6
rdkit	2026.9.1
scikit-learn	1.9.1
scipy	1.18.1
torch	2.11.0+cu128

4.3 Source file hashes, SHA-256

File	SHA-256 (first 40 characters)
data\reference\pdac_reference_dataset.json	10de36cbf66cb3f74e2f79dd40a355847247b3bd
models\scientific\pdac_qsar_v1.joblib	088c53985fe5ece80dda27fb6a073c219292c1d5
simulation\biology\diseases\pdac.py	38d708f39a7977c0d8dadd07e9c72bffe219306f
simulation\core\engine.py	4059a6fdea8603c90d4abe897475d3b48aaed4b0
simulation\discovery\pipeline.py	b2d12142a46607238489c805bfd2dfba6cb5c22a
simulation\discovery\workbench.py	a1bf63f8e409aa7e407d316beee8c21a06a20fcb
simulation\pharmacology\admet.py	064fe566b36565c07af097cb99a379543c6ddd4b

4.4 Reproducing this document

The platform is open source and so is this result. The reference dataset, the pipeline, the tissue models and the report generator are all in the repository. No model weights are required: the QSAR artifact is rebuilt from the committed dataset, and the optional language models were not used for any quantitative output.

```
python scripts/train_qsar.py

# Linux and macOS
EVOCURE_COMPUTE=cpu EVOCURE_DISABLE_LAYA_NEURAL=1 \
  python scripts/run_pdac_report_experiment.py

# Windows PowerShell
$env:EVOCURE_COMPUTE = "cpu"
$env:EVOCURE_DISABLE_LAYA_NEURAL = "1"
python scripts\run_pdac_report_experiment.py

python scripts/build_candidate_pdf.py
```

Forcing the CPU backend is what makes a run bit reproducible. GPU backends that support float64 agree to within 1e-12 on the substrate fields; float32 backends such as DirectML and Metal differ at around 1e-6, which is harmless for the conclusions and fatal for hash

equality. The backend used is recorded above and in the results file.

4.5 Licence and reuse

The platform is released under the MIT licence and the curated reference dataset under CC-BY 4.0. Results, including this report and the underlying JSON, are published so that the discoveries stay open: anyone may reproduce, check, contradict or build on them without permission.