

SHEET 1

Exercise 3

(a) Let (X, S) be a Δ -complex.

Claim 0.1. *A subset $A \subseteq X$ is closed if and only if $\sigma^{-1}(A)$ is closed for all $\sigma \in S$.*

Proof. Follows immediately from $\sigma^{-1}(A^c) = \sigma^{-1}(A)^c$ where c is the set-theoretic complement. \square

Define $X^{(n)} := \bigcup_{k \leq n} \bigcup_{\sigma \in S_k} \text{im}(\sigma)$, the union of all the images of k -simplices for $k \leq n$.

Claim 0.2. *$X^{(n)}$ is a closed subspace of X .*

Proof. If $\sigma \in S_k$, then $\sigma^{-1}(X^{(n)})$ is the union of all faces of Δ^k of dimension $\leq n$, which is always a closed subset. \square

Claim 0.3. *A subset $A \subseteq X^{(n)}$ is closed if and only if $\sigma^{-1}(A)$ is closed for all $\sigma \in S_k$ and $k \leq n$.*

Proof. Follows from $\sigma^{-1}(A) = \bigcup_{\tau} \tau^{-1}(A)$, where τ runs through the faces of σ of dimension $\leq n$, and there are only finitely many terms in the union. \square

Claim 0.4. *The filtration $X^{(0)} \subseteq X^{(1)} \subseteq \dots \subseteq X$ is a CW-structure on X .*

Proof. We will check two things:

(1) For every $n \geq 0$ there is a pushout diagram

$$\begin{array}{ccc} \bigsqcup_{\sigma \in S_n} \partial \Delta^n & \longrightarrow & \bigsqcup_{\sigma \in S_n} \Delta^n \\ \downarrow & & \downarrow \sum_{\sigma \in S_n} \sigma \\ X^{(n-1)} & \longrightarrow & X^{(n)} \end{array}$$

(for $n = 0$ this says that $X^{(0)} \cong S_0$ is a discrete space)

(2) A subset $A \subseteq X$ is closed if and only if $A \cap X^{(n)}$ is closed for every $n \geq 0$.

A commutative diagram like the above exists, because of condition (i) in the definition of a Δ -complex. So we get an induced (continuous) map

$$f: X^{(n-1)} \coprod_{\bigsqcup_{\sigma \in S_n} \partial \Delta^n} \bigsqcup_{\sigma \in S_n} \Delta^n \rightarrow X^{(n)}.$$

This map is a bijection, precisely because of condition (ii) in the definition of a Δ -complex. To see that the map is actually a homeomorphism, we check that it is closed: So let $A \subseteq X^{(n-1)} \coprod_{\sigma \in S_n} \partial \Delta^n \sqcup_{\sigma \in S_n} \Delta^n$ be a closed subset. Then $f(A)$ is closed if $\sigma^{-1}(f(A))$ is closed for all $\sigma \in S_k$ and $k \leq n$ (by Claim 0.3). But this translates precisely into the condition of A being closed in the quotient topology of $X^{(n-1)} \coprod_{\sigma \in S_n} \partial \Delta^n \sqcup_{\sigma \in S_n} \Delta^n$. By induction (starting at the trivial case $n = -1$) this proves (1).

(2) follows in one direction from Claim 0.2 and in the other direction from Claim 0.3. \square

(b) Recall the cellular chain complex:

$$\begin{array}{ccccc}
 C_n^{cell}(X) & \xlongequal{\quad} & H_n(X^{(n)}, X^{(n-1)}) & \xleftarrow[\cong]{\sum_{\sigma} \sigma_*} & \bigoplus_{\sigma \in S_n} H_n(\Delta^n, \partial \Delta^n) & \xlongequal{\quad} & C_n^{\Delta}(X) \\
 \downarrow \partial^{cell} & & \downarrow \partial & & & & \\
 & & H_{n-1}(X^{(n-1)}) & & & & \\
 & & \downarrow j_* & & & & \\
 C_{n-1}^{cell}(X) & \xlongequal{\quad} & H_{n-1}(X^{(n-1)}, X^{(n-2)}) & & & &
 \end{array}$$

A simplex $\sigma \in S_n$ induces a map σ_* as indicated; now note that $H_n(\Delta^n, \partial \Delta^n)$ has a canonical generator, namely the relative homology class represented by $id: \Delta^n \rightarrow \Delta^n$. We then obtain

$$\partial^{cell}(\sigma_*([id])) = j_*\partial([\sigma]) = j_*\left[\sum_{i=0}^n (-1)^i \sigma \circ d^i\right] = \sum_{i=0}^n (-1)^i [\sigma \circ d^i]$$

which is precisely what the differential in $C_*^{\Delta}(X)$ would do. In other words, the tautological map $\sum_{\sigma \in S_n} \sigma_*$ indicated in the diagram above is the isomorphism we are looking for.