

# Contents

1	Lecture 1	1
2	Lecture 2	3

## 1 Lecture 1

Recall that the **tangent bundle** is also a smooth manifold, with the natural footpoint projection given by

$$\pi : TM \rightarrow M. \quad (1)$$

A **section** of  $\pi$  is a (smooth) map  $X : M \rightarrow TM$  such that  $\pi \circ X = \text{id}_M$ .

### Definition 1.0.1: Riemannian Metric & Riemannian Manifold

A  $(C^k\text{-})$ **Riemannian metric**  $g$  on  $M$  is a family  $g = \{g_p\}_{p \in M}$  of inner products (positive definite, symmetric bilinear)

$$g_p : T_p M \times T_p M \rightarrow R \quad (2)$$

such that for any smooth vector fields  $X, Y$  on  $M$ , the function

$$f : M \rightarrow R, \quad p \mapsto g_p(X_p, Y_p) \quad (3)$$

is smooth ( $C^k$ -differentiable). The pair  $(M, g)$  is called a **Riemannian manifold**.

### Remarks.

1. Given a  $C^\infty$  atlas of  $M$ , then a family of inner products  $g = \{g_p\}_{p \in M}$  is a  $(C^k\text{-})$ Riemannian metric iff for any chart  $(U, x)$  in this atlas the function

$$U \rightarrow R, \quad p \mapsto g_p\left(\frac{\partial}{\partial x_j}\Big|_p, \frac{\partial}{\partial x_k}\Big|_p\right) \quad (4)$$

is smooth. (To be shown in an exercise class.)

2. Suppose we have an open subset  $U$  inside a  $(C^k\text{-})$ Riemannian manifold  $(M, g)$ . Then  $(U, g|_U)$  is also a  $(C^k\text{-})$ Riemannian manifold, where

$$g|_U := \{g_p\}_{p \in U}. \quad (5)$$

3. We sometimes write  $\langle \cdot, \cdot \rangle$  instead of  $g$  (respectively  $\langle \cdot, \cdot \rangle_p$  instead of  $g_p$ ).

4. Sometimes, also indefinite families of inner products are considered, so-called **semi-Riemannian metrics/manifolds**, e.g. in the theory of relativity where you have Lorentzian metrics, i.e. semi-Riemannian metrics of signature  $(1, n)$  like the Lorentzian metric

$$x_0 y_0 - \sum_{i=1}^n x_i y_i \text{ on } \mathbb{R}^{n+1}. \quad (6)$$

A simple example of a Riemannian manifold: any open subset  $U$  of a Euclidean vector space  $(V, \langle \cdot, \cdot \rangle)$  is a Riemannian manifold with Riemannian metric  $g_p = \langle \cdot, \cdot \rangle_p$ ,  $p \in U$ .

**Proposition 1.0.2: Riemannian Immersions**

Let  $M$  a smooth manifold,  $(N, h)$  a Riemannian manifold,  $\varphi : M \rightarrow N$  a smooth  $(C^{k+1})$ -immersion. Then  $(N, h)$  induces a Riemannian metric  $g$  on  $M$  via

$$g_p(v, w) = h_{\varphi(p)}(d\varphi_p v, d\varphi_p w); \quad p \in M; v, w \in T_p M. \quad (7)$$

This Riemannian metric is called the **pullback of  $h$  via  $\phi$** , denoted  $\varphi^*h$ . The immersion  $\varphi : (M, g) \rightarrow (N, h)$  is called a **Riemannian immersion**.

*Proof.* Since  $d\varphi_p : T_p M \rightarrow T_{\varphi(p)} N$  is linear and injective,  $\forall p \in M$ ,  $g_p$  is bilinear and positive definite (as  $h_{\varphi(p)}$  is).

Symmetry of  $h_{\varphi(p)} \Rightarrow$  symmetry of  $g_p \rightarrow g_p$  an inner product.

*Smoothness.* Let's first assume that  $\varphi$  is a (local) diffeomorphism. Then a smooth vector field  $X$  on  $M$  induces a smooth vector field on  $N$ , namely

$$(\varphi^* X) : q \mapsto d\varphi_{\varphi^{-1}(q)} X_{\varphi^{-1}(q)} \quad (8)$$

and we have that

$$p \mapsto g_p(X_p, Y_p) = h_p((\varphi^* X)_{\varphi(p)}, (\varphi^* Y)_{\varphi(p)}) \quad (9)$$

$$= (h((\varphi^* X)_., (\varphi^* Y)_.) \circ \varphi)(p) \quad (10)$$

For the general case, one can now apply the local structure for immersions and assume that

$$M = U \subset \mathbb{R}^n \text{ open}, N = V \subset \mathbb{R}^{n+k} \text{ open} \quad (11)$$

and  $\varphi = i : U \hookrightarrow V$  the inclusion. Next, extend the vector field  $X$  on  $U$  to

$$\tilde{X}(p) = (X(p), 0) \text{ on } V. \quad (12)$$

( $\rightarrow$  details next Wednesday).  $\square$

**Remarks.**

1. By Whitney's immersion theorem, any smooth manifold can be immersed (actually embedded) into some  $\mathbb{R}^n$ . So it inherits a Riemannian metric from  $\mathbb{R}^n$ .
2. By Nash's embedding, any  $(C^{k \geq 3})$ -Riemannian manifold can be isometrically immersed (embedded) into some  $\mathbb{R}^n$ .

**Note.** If  $g, h$  are Riemannian metrics on  $M$ , then also  $(g + h)_p = g_p + h_p$  defines a Riemannian metric on  $M$ .

$$f : M \rightarrow \mathbb{R} \quad (13)$$

is a positive smooth function, then

$$(f \cdot g)_p = f_p g_p \quad (14)$$

defines a Riemannian metric on  $M$ .

### Example 1.0.3: Balls and Riemannian Metric.

$B_1^n$  together with the Riemannian metric  $g$ , given by

$$g_p(v, w) = \frac{1}{(1 - \langle p, p \rangle)^2} \langle v, w \rangle_p \quad (15)$$

defines a Riemannian manifold isometric to hyperbolic space.

## 2 Lecture 2

### Definition 2.0.1: Isometry

Let  $(M, g)$  and  $(N, h)$  be Riemannian manifolds.

1. A diffeomorphism  $\varphi : M \rightarrow N$  is called an **isometry** if  $\varphi^* h = g$ , i.e.

$$g_p(v, w) = h_{\varphi(p)}(d\varphi_p v, d\varphi_p w) \quad (16)$$

$\forall p \in M; v, w \in T_p M$ .

2. A smooth map  $\varphi : M \rightarrow N$  is called a **local isometry** if every point  $p \in M$  has an open neighborhood  $U$  such that  $\varphi|_U : U \rightarrow \varphi(U)$  is an isometry.

3. The set of all isometries of  $(M, g)$

$$\text{iso}(M, g) = \{\varphi : M \rightarrow M \mid \varphi \text{ is an isometry}\} \quad (17)$$

is a group, the so called **isometry group**.

**Remarks.**

1. If  $\varphi : (M, g) \rightarrow (N, h)$  is a local isometry, then

$$d\varphi_p : T_p M \rightarrow T_p N \quad (18)$$

is a linear isometry.

#### Example 2.0.2: exp is a Local Isometry

Consider  $\mathbb{R}$  with Euclidean inner product  $g$ . Further, consider  $S^1 \subset \mathbb{R}^2 \subset \mathbb{C}^1$  with the induced Riemannian metric. Then

$$\exp : \mathbb{R} \rightarrow S^1, \quad t \mapsto e^{it} \quad (19)$$

is a local isometry. For

$$\|(d \exp)_t \left( \frac{\partial}{\partial t} \right)\| = \left| \frac{d}{dt} (e^{it}) \right| = |i \cdot e^{it}| = 1 \rightarrow d \exp_t \neq 0 \quad (20)$$

which shows that  $\exp$  is a local isometry, but not an isometry:  $\mathbb{R} \not\cong S^1$ , not even homeomorphic.

#### Definition 2.0.3: Group action, Isometric action

Let  $G$  be a group,  $M$  be a smooth manifold. A **(smooth) action** of  $G$  on  $M$  is a homomorphism

$$\phi : G \rightarrow \text{Diff}(M), \quad a \mapsto \phi_a; \quad (21)$$

i.e.,

$$\phi_{a_1 a_2}(x) = (\phi_{a_1} \circ \phi_{a_2})(x) \quad \forall a_1, a_2 \in G, x \in M \quad (22)$$

where  $\phi_e = \text{id}_M$ , and  $e \in G$  is the neutral element. We sometimes denote such an action as  $G \curvearrowright M$ , and write  $\phi_a(x)$  as  $ax$ . If  $(M, g)$  is a Riemannian manifold, and  $\phi(G) \subset \text{Iso}(M, g)$ , then we call this action **isometric**.

#### Definition 2.0.4: Free Group Action

A group action  $G \curvearrowright M$  is called **free** if for any  $g \in G$ ,  $x \in M$ ,  $gx = x \implies g = e$ .

#### Definition 2.0.5: Covering Space Action

A smooth action  $\Gamma \curvearrowright M$  of a group  $\Gamma$  on a smooth manifold  $M$  is called a **covering space action** if the following holds:

Every point  $x \in M$  has an open neighborhood  $U$  such that  $\gamma U \cap U = \emptyset \quad \forall \gamma \in \Gamma \setminus \{e\}$ .

In particular,  $\Gamma \curvearrowright M$  is free.

**Proposition 2.0.6: Quotient of Smooth Manifold by Covering Space Action**

*Let  $\Gamma \curvearrowright M$  be a smooth covering space action of a group  $\Gamma$  on a manifold  $M$ . Then  $M/\Gamma$  is a topological manifold with a unique smooth structure with respect to which the projection  $M \rightarrow M/\Gamma$  is a local diffeomorphism.*

*Proof.* See my live notes for the proof. Please examine it carefully.  $\square$

A comment is made on the universal cover of  $M$ , and examples are provided in the live notes.

Conversely, we have:

**Proposition 2.0.7: Riemannian Metric and Local Isometry**

*Let  $\Gamma \curvearrowright (M, g)$  be a covering space action by isometries on a Riemannian manifold  $(M, g)$ . Then there exists a unique Riemannian metric  $\bar{g}$  on  $M/\Gamma$  such that  $\pi : (M, g) \rightarrow (M/\Gamma, \bar{g})$  is a local isometry.*

Examples are provided, see live notes.

## Index

action, 4

covering space action, 4

free group action, 4

isometric, 4

isometry, 3

isometry group, 3

local isometry, 3

Riemannian immersion, 2

Riemannian manifold, 1

Riemannian metric, 1

section, 1

semi-Riemannian metrics/manifolds, 2

smooth action, 4

tangent bundle, 1