

Class10

```
round(183201/251600768*100, 3)
```

```
[1] 0.073
```

```
stats <- read.csv("Data Export Summary.csv", row.names=1)
head(stats)
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	158,844	11,759	12,296	197	73	32
Protein/Oligosaccharide	9,260	2,054	34	8	1	0
Protein/NA	8,307	3,667	284	7	0	0
Nucleic acid (only)	2,730	113	1,467	13	3	1
Other	164	9	32	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	183,201					
Protein/Oligosaccharide	11,357					
Protein/NA	12,265					
Nucleic acid (only)	4,327					
Other	205					
Oligosaccharide (only)	22					

Convert characters to numerics

```
string <- c("10", "100", 1)
as.numeric(string) + 1
```

```
[1] 11 101 2
```

However, this method doesn't work with numerics with commas, like 1,000.

gsub() function will replace every occurrence of the character. **sub()** will only replace the first occurrence.

#ChatGPT's way

```
remove_commas <- function(df) {  
  df[] <- lapply(df, function(x) gsub(",", "", x))  
  return(df)  
}  
stats2 <- remove_commas(stats)
```

#Barry's way

```
rm.comma <- function(x) {  
  as.numeric( gsub(",", "", x))  
}  
  
pdbstats <- apply(stats, 2, rm.comma)
```

Q1: What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy?

```
totals <- apply(pdbstats, 2, sum)  
round(totals/totals["Total"] * 100, 2)
```

X-ray	EM	NMR	Multiple.methods
84.83	8.33	6.68	0.11
Neutron	Other	Total	
0.04	0.02	100.00	

93.16%


Q2: What proportion of structures in the PDB are protein?

skipped > Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?

skipped

Using Mol* to examine HIV-Pr

Here is a shitty picture of HIV-Pr that is not very useful yet.



Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

The resolution is 2 angstroms, which is not enough to show the hydrogen atoms.

Q5: There is a critical “conserved” water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have

HOH308

Q6: Generate and save a figure clearly showing the two distinct chains of HIV-protease along with the ligand. You might also consider showing the catalytic residues ASP 25 in each chain and the critical water (we recommend “Ball & Stick” for these side-chains). Add this figure to your Quarto document.



```
# Using the bio3d package
```

```
library(bio3d)
pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
```

```
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

```
head(pdb$atom$resid)
```

```
[1] "PRO" "PRO" "PRO" "PRO" "PRO" "PRO"
```

```
pdb$atom$resid[pdb$calpha]
```

```
[1] "PRO" "GLN" "ILE" "THR" "LEU" "TRP" "GLN" "ARG" "PRO" "LEU" "VAL" "THR"
[13] "ILE" "LYS" "ILE" "GLY" "GLY" "GLN" "LEU" "LYS" "GLU" "ALA" "LEU" "LEU"
[25] "ASP" "THR" "GLY" "ALA" "ASP" "ASP" "THR" "VAL" "LEU" "GLU" "GLU" "MET"
[37] "SER" "LEU" "PRO" "GLY" "ARG" "TRP" "LYS" "PRO" "LYS" "MET" "ILE" "GLY"
[49] "GLY" "ILE" "GLY" "GLY" "PHE" "ILE" "LYS" "VAL" "ARG" "GLN" "TYR" "ASP"
[61] "GLN" "ILE" "LEU" "ILE" "GLU" "ILE" "CYS" "GLY" "HIS" "LYS" "ALA" "ILE"
[73] "GLY" "THR" "VAL" "LEU" "VAL" "GLY" "PRO" "THR" "PRO" "VAL" "ASN" "ILE"
[85] "ILE" "GLY" "ARG" "ASN" "LEU" "LEU" "THR" "GLN" "ILE" "GLY" "CYS" "THR"
[97] "LEU" "ASN" "PHE" "PRO" "GLN" "ILE" "THR" "LEU" "TRP" "GLN" "ARG" "PRO"
[109] "LEU" "VAL" "THR" "ILE" "LYS" "ILE" "GLY" "GLY" "GLN" "LEU" "LYS" "GLU"
[121] "ALA" "LEU" "LEU" "ASP" "THR" "GLY" "ALA" "ASP" "ASP" "THR" "VAL" "LEU"
[133] "GLU" "GLU" "MET" "SER" "LEU" "PRO" "GLY" "ARG" "TRP" "LYS" "PRO" "LYS"
[145] "MET" "ILE" "GLY" "GLY" "ILE" "GLY" "GLY" "PHE" "ILE" "LYS" "VAL" "ARG"
[157] "GLN" "TYR" "ASP" "GLN" "ILE" "LEU" "ILE" "GLU" "ILE" "CYS" "GLY" "HIS"
[169] "LYS" "ALA" "ILE" "GLY" "THR" "VAL" "LEU" "VAL" "GLY" "PRO" "THR" "PRO"
[181] "VAL" "ASN" "ILE" "ILE" "GLY" "ARG" "ASN" "LEU" "LEU" "THR" "GLN" "ILE"
[193] "GLY" "CYS" "THR" "LEU" "ASN" "PHE"
```

Q7: How many amino acid residues are there in this pdb object?

198

Q8: Name one of the two non-protein residues?

HOH (127), MK1 (1)

Q9: How many protein chains are in this structure?

2

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMLRAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKFNPVKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```


```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

```
modes <- nma(adk)
```

```
Building Hessian... Done in 0.014 seconds.
```

```
Diagonalizing Hessian... Done in 0.256 seconds.
```

```
plot(modes)
```



```
mktrj(modes, file="modes.pdb")
```