

Hysterical Response from Quantum Alternatives

Two- and Three-Particle Thought Experiments, Nonlocal Correlations,
and the Conjectured Macroscopic Survival of Gravity

Working conceptual formulation

September 2026

Abstract

Quantum mechanics allows physical systems to occupy superpositions of alternatives. In realistic macroscopic systems, interaction with the environment rapidly decoheres those alternatives, making them behave as effectively separate outcomes. Decoherence is extraordinarily effective, but “effectively zero” is not automatically the same statement as “exactly zero.” This motivates a simple question: could an open quantum system retain an extremely small dynamical response to alternative interaction histories that have been almost, but not perfectly, erased from its local history?

We call the hypothetical residual, history-dependent effect the **Hysterical Response**. This paper is intentionally nontechnical. It does not derive the effect from a fundamental theory. Instead, two-, three-, and four-particle thought experiments are used to explain what such a response would mean, why correlations make it intrinsically nonlocal at the level of the joint quantum state, and why a macroscopic response need not be a simple sum of independent particle contributions. We conjecture that every fundamental interaction can exhibit microscopic Hysterical Response, while only the gravitational response remains macroscopically appreciable in an appropriate many-particle limit. These conjectures are motivations, not results. The purpose of Paper 0 is to make the physical idea clear before the later papers ask whether it survives mathematics and observation.

Contents

1	The question	2
2	The Hysterical Response	2
3	Toy model I: two particles	3
4	Toy model II: three particles and the correlation effect	3
5	What we mean by nonlocal	4

6	Toy model III: four particles and higher correlations	4
7	Why should the effect be gravitational?	5
8	Then why do we not see Hysterical electromagnetism everywhere?	5
9	The gravitational survival conjecture	6
10	From a few particles to a galaxy	6
11	Where does the system end?	6
12	What has been shown, assumed, and conjectured	7
13	Why call it Hysterical?	7
14	The conceptual picture	8
15	What comes next	8
16	Conclusion	8

1 The question

Consider two particles, A and B . In ordinary Newtonian gravity, if their masses are m_A and m_B and their separation is r , the attraction has magnitude

$$F_N = \frac{Gm_A m_B}{r^2}. \quad (1)$$

Nothing unusual has happened yet.

Now suppose B is placed in a spatial superposition,

$$|B_L\rangle + |B_R\rangle. \quad (2)$$

This is the same basic idea expressed by the familiar shorthand

$$|\text{cat alive}\rangle + |\text{cat dead}\rangle. \quad (3)$$

Real objects do not normally appear as visible superpositions because they interact continuously with their surroundings. Photons scatter, air molecules collide, heat is exchanged, and information about the alternatives spreads into the environment. Schematically,

$$|B_L\rangle |E_L\rangle + |B_R\rangle |E_R\rangle. \quad (4)$$

For a strongly decohered system,

$$\langle E_R | E_L \rangle \approx 0. \quad (5)$$

The symbol $\approx$ is the point of departure. What if an open system can retain an extraordinarily small dynamical memory of alternatives that have become effectively, but not perfectly, separated?

2 The Hysterical Response

Conjectural phenomenon: Hysterical Response

An open quantum system may retain an extremely small dynamical response to alternative interaction histories that have been strongly, but not perfectly, decohered.

The name is intentional. Ordinary *hysteresis* describes systems whose present behaviour depends partly on their past. Here we use the deliberately unconventional adjective *Hysterical* for a proposed response whose present value retains information about quantum alternatives in the system's interaction history.

We denote an effective Hysterical force contribution by $\mathbf{F}_H$ and write phenomenologically

$$\boxed{\mathbf{F}_{\text{observed}} = \mathbf{F}_{\text{ordinary}} + \mathbf{F}_H}. \quad (6)$$

This is not proposed as a fundamental quantum force operator. It is the quantity that an observer using ordinary dynamics would interpret as an additional force. Paper 1 asks whether such a response can arise consistently from open-system dynamics; Paper 0 only explains the idea.

3 Toy model I: two particles

Let A be a probe and B a source. Suppose B has a near alternative B_N and a far alternative B_F , at distances $r_N < r_F$. Then

$$F_N = \frac{Gm_A m_B}{r_N^2}, \quad F_F = \frac{Gm_A m_B}{r_F^2}, \quad F_N > F_F. \quad (7)$$

Suppose the experienced configuration contains B_N . Ordinary reasoning gives F_N . The Hysterical conjecture asks whether the alternative interaction history involving F_F can leave a tiny residual response,

$$F_{\text{observed}} = F_N + F_H. \quad (8)$$

As a mnemonic only, one might imagine

$$F_H \sim \varepsilon (F_N - F_F), \quad |\varepsilon| \ll 1, \quad (9)$$

where ε represents an extremely small surviving memory. This is deliberately *not* a microscopic law.

The two-particle example gives the first intuition:

different quantum interaction histories $\implies$ possible residual dynamical response.

(10)

But it also makes the effect look more local and additive than it need be.

4 Toy model II: three particles and the correlation effect

Now let A remain the probe while B and C are sources. Each can be near or far from A . Compare

$$|\Psi_+\rangle = \frac{1}{\sqrt{2}} (|B_N C_N\rangle + |B_F C_F\rangle), \quad (11)$$

$$|\Psi_-\rangle = \frac{1}{\sqrt{2}} (|B_N C_F\rangle + |B_F C_N\rangle). \quad (12)$$

If we look at B alone, both states give

$$P(B_N) = P(B_F) = \frac{1}{2}. \quad (13)$$

The same is true for C :

$$P(C_N) = P(C_F) = \frac{1}{2}. \quad (14)$$

So the individual behaviour of B is the same in both experiments, and the individual behaviour of C is the same. What changes is their *correlation*.

Assume equal source masses and a symmetric geometry. Let one near source contribute force F_N and one far source F_F . In $|\Psi_+\rangle$ the alternatives have total forces

$$F_{NN} = 2F_N, \quad F_{FF} = 2F_F, \quad (15)$$

so their contrast is

$$\Delta F_+ = 2(F_N - F_F). \quad (16)$$

For $|\Psi_-\rangle$,

$$F_{NF} = F_N + F_F, \quad F_{FN} = F_F + F_N, \quad (17)$$

and therefore

$$\boxed{\Delta F_- = 0.} \quad (18)$$

This is the central motivating example. Two states with identical one-particle statistics can expose A to very different *joint* gravitational histories. A Hysterical Response that remembers those histories can therefore satisfy

$$\boxed{F_H^{(A)}[\Psi_+] \neq F_H^{(A)}[\Psi_-],} \quad (19)$$

even though the separate statistics of B and C have not changed.

The “oh, I get it” point

The Hysterical Response cannot in general be just “a tiny extra force from every unrealized particle position.” If it remembers alternative interaction histories, it must be capable of remembering how the alternatives of different particles are correlated.

5 What we mean by nonlocal

The three-particle example immediately makes the response nonlocal in an important but limited sense. The response at A need not be determined by independent local descriptions of B and C ; it may depend on their joint state:

$$\boxed{\mathbf{F}_H(A) = \mathcal{F}_H[\text{joint state, correlations, environmental history}].} \quad (20)$$

Changing correlations among spatially separated components can therefore change the Hysterical response at A even when their individual reduced descriptions remain unchanged.

This is *state nonlocality*. We do not claim here that it permits faster-than-light signalling, nor that relativistic causality is violated. Those are stronger questions requiring a relativistic theory that has not yet been constructed.

6 Toy model III: four particles and higher correlations

Three particles show that pair correlations can matter. A fourth raises the possibility that still higher joint structure matters.

Let A be the probe and B, C, D the source system. The response might admit a schematic hierarchy

$$F_H = F_H^{(1)} + F_H^{(2)} + F_H^{(3)} + \cdots, \quad (21)$$

where the superscripts denote information carried by one-source structure, pair correlations, irreducible three-source correlations, and so on. We do not assert that this expansion is fundamental; it simply illustrates why the many-particle response need not reduce to a sum over independent pairs.

The lesson is that the microscopic Hysterical Response may be **configuration-sensitive**, rather than merely **density-sensitive**.

7 Why should the effect be gravitational?

Nothing in the motivating argument so far knows that we chose gravity. Replace the gravitational interaction by electrostatics,

$$V_{\text{EM}}(r) = \frac{q_A q_B}{4\pi\epsilon_0 r}. \quad (22)$$

A spatial superposition again corresponds to alternative interaction energies and forces. If Hysterical Response originates from quantum alternatives and environmental memory, rather than from a special property of gravity, it is natural to ask whether electromagnetism should exhibit it too.

Conjecture 1: Interaction universality

Every fundamental interaction can, in principle, exhibit a microscopic Hysterical Response when an open quantum system contains incompletely erased alternative interaction histories.

Symbolically,

$$\boxed{\mathbf{F}_X^{\text{observed}} = \mathbf{F}_X^{\text{ordinary}} + \mathbf{F}_{H,X}}, \quad (23)$$

where X labels an interaction channel. This is a conjecture. A microscopic theory could instead show that gravity or stress-energy is special.

8 Then why do we not see Hysterical electromagnetism everywhere?

Electromagnetism is enormously stronger than gravity between elementary charged particles. If its Hysterical contribution survived straightforwardly into bulk matter, that would be an immediate problem.

But ordinary matter is approximately electrically neutral. Positive and negative charge can cancel:

$$(+q) + (-q) = 0. \quad (24)$$

Ordinary Newtonian mass does not have the analogous cancellation in normal matter: positive mass contributions add.

This gives an intuitive reason why a microscopic effect shared by several interactions might nevertheless appear gravitational on large scales. It does not prove it.

Suppose a positive and negative charge acquire Hysterical weights h_+ and h_- . Their residual contribution could contain

$$qh_+ - qh_-. \quad (25)$$

Charge neutrality says $q - q = 0$; it does not imply $qh_+ - qh_- = 0$ unless the response dynamics enforce $h_+ = h_-$. Because the weights themselves may depend on nonlocal correlations and history, neutrality alone is insufficient.

9 The gravitational survival conjecture

We therefore separate two claims: microscopic existence and macroscopic survival.

Conjecture 2: Gravitational survival

Under the appropriate macroscopic coarse-graining and many-particle limit, nongravitational Hysterical responses vanish or become macroscopically negligible, while the gravitational Hysterical Response remains macroscopically appreciable.

Very schematically,

$$N \rightarrow \infty : \quad F_{H,g} \text{ survives,} \quad F_{H,X} \rightarrow 0 \text{ macroscopically for } X \neq g. \quad (26)$$

We do not yet define the correct thermodynamic or coarse-grained limit, and we do not assume a particular N -scaling. For electromagnetism, signed charge cancellation is suggestive; for the strong interaction, confinement is suggestive; for the weak interaction, short range is suggestive. None is yet a derivation of the Hysterical many-body limit.

10 From a few particles to a galaxy

For one microscopic system, any residual response after decoherence could be fantastically small. An astronomical open system is different in scale and structure. A galaxy contains enormous numbers of particles, continually forms new structures, exchanges matter with its surroundings, emits radiation, and continually creates new environmental records.

The motivating possibility is

tiny residual response per alternative $\times$ enormous effective multiplicity $\longrightarrow$ finite macroscopic response.
--

(27)

This is qualitative. It is not a literal count of Everett branches, and it does not assume that microscopic contributions add independently. The three-particle model already tells us that correlations can change the answer.

A macroscopic response variable must therefore be allowed to summarize more than local mass density. Schematically one might eventually require

$$Q = Q[\text{matter, correlations, environment, history}]. \quad (28)$$

Later papers use an effective response variable of this kind without claiming that its microscopic form has already been derived.

11 Where does the system end?

Nonlocal correlation sensitivity creates a boundary problem. If we draw a sphere of radius R around a galaxy and call everything inside “the galaxy,” why should the Hysterical Response care about our

bookkeeping choice? Matter outside the chosen region may remain relevant through correlations or through the response history.

A physically meaningful quantity might eventually involve a large-region limit such as

$$\lim_{R \rightarrow \infty} Q(R), \quad (29)$$

if that limit exists, or a dynamically generated memory kernel that suppresses sufficiently distant contributions. Paper 0 does not choose between these possibilities. The point is simply that once correlations matter, defining the relevant system becomes a physical question.

12 What has been shown, assumed, and conjectured

To keep the logic explicit, the status of the claims in this paper is as follows.

Elementary mathematical observation. Two joint states can have identical one-particle marginals while possessing different correlations. In the symmetric three-particle example above, those correlations produce different branch-by-branch gravitational force contrasts at A .

Phenomenological postulate. A residual history-dependent force-like quantity, called the Hysterical Response, is allowed to depend on alternative interaction histories.

Conjecture. The Hysterical Response can depend on joint correlations and environmental history, making it nonlocal in the joint-state sense described above.

Conjecture. Every fundamental interaction can exhibit microscopic Hysterical Response.

Conjecture. Only gravity remains macroscopically appreciable in the appropriate $N \rightarrow \infty$ limit.

We have not derived the Hysterical Response from quantum mechanics; proved that residual environmental overlap generates an observable force; proved electromagnetic cancellation; established the many-particle limit; constructed a relativistic completion; or shown that the effect explains dark-matter phenomenology.

13 Why call it Hysterical?

The proposed phenomenon is fundamentally about memory. Its present value is conjectured not to depend solely on the instantaneous classical configuration, but also on alternative configurations through which the quantum state and its environment have evolved.


In ordinary hysteresis, history matters. Here, quantum history matters. Hence

$$\boxed{\text{Hysterical Response.}} \quad (30)$$

A corresponding macroscopic field will be called the **Hysterical Response field**, or simply the **Hysterical field** where the meaning is clear.

14 The conceptual picture

The proposal can be summarized without technical machinery:



The two-particle example says what a residual response would mean. The three-particle example explains why correlations matter. The four-particle example warns that higher-order structure may matter too. Repeating the thought experiment with Coulomb interactions motivates microscopic interaction universality. The contrasting behaviour of signed charge and positive Newtonian mass motivates, but does not prove, gravitational macroscopic survival.

15 What comes next

The simplicity of Paper 0 conceals a serious problem. One cannot simply declare that an off-diagonal part of a Newtonian force operator is “force leaking between branches.” Under natural localization assumptions that construction can vanish identically.

Paper 1 therefore asks the first technical question: can a Hysterical Response arise through legitimate open-system dynamics, with coherent mixing and dissipation converting residual alternative structure into an ordinary population/response effect?

A subsequent many-body paper asks a different question: if microscopic Hysterical Response is interaction-general, which interaction channels actually survive coarse-graining as N becomes large? In particular, can electromagnetic Hysterical Response disappear in neutral macroscopic matter without simply assuming the desired answer?

Only after those questions are separated does it make sense to investigate a surviving gravitational Hysterical field on galactic scales.

16 Conclusion

Paper 0 proposes a simple physical possibility. Decoherence makes quantum alternatives effectively separate, but an open system may not erase their dynamical influence with infinite perfection. We call any resulting residual, history-dependent effect the **Hysterical Response**.

The most important motivating example is not the two-particle superposition but the three-particle correlation experiment. Two source systems can have identical individual particle statistics while possessing different joint alternatives and therefore different interaction histories at a probe. If the Hysterical Response remembers those histories, it must be capable of depending on correlations rather than only on independent local densities.

This motivates a response that is nonlocal in the joint-state sense. We conjecture that the microscopic mechanism is interaction-general, but separately conjecture that only gravity remains macroscopically appreciable in an appropriate many-particle limit. Neither statement is assumed to be proved by the examples.

The remainder of the series asks whether this intuitive picture survives increasingly demanding mathematics and observation.