

maybe

¹ Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA; rbuta@bama.ua.edu

² Instituto de Astrofísica de Canarias, E-38200 La Laguna, Spain; jhk@iac.es

³ IBM Research Division, T.J. Watson Research Center, 1101 Kitchawan Road, Yorktown Heights, NY 10598, USA; bge@watson.ibm.com

⁴ Division of Astronomy, Department of Physical Sciences, University of Oulu, Oulu FIN-90014, Finland; hsalo@sun3.oulu.fi, eiija@sun3.oulu.fi

⁵ Vassar College, Department of Physics & Astronomy, Box 745, Poughkeepsie, NY 12604, USA; elmegreen@vassar.edu

⁶ Instituto Nacional de Astrofísica, Óptica y Electrónica, Tonantzintla, PUE 72840, Mexico; puerari@inaoep.mx

⁷ Anglo American Cosmic Dust Laboratory, School of Computational & Applied Mathematics, University of the Witwatersrand, P.O. Box 60 Wits, 2050, South Africa; David.Block@wits.ac.za

Received 2008 October 22; accepted 2009 February 28; published 2009 April 7

doi:10.1088/2041-8205/71

© 2010, The American Astronomical Society. All rights reserved. Printed in the U.S.A.

We present deep near-infrared images of a selected sample of near-infrared galaxies. The sample covers a range of Hubble types, and the images show that the dust spirals correlate with those predicted by theoretical models. Analysis of the images shows that the effects of extinction are a few times stronger than those of a few excessively strong bars. The dust is relatively weak. We find that the presence of a very strong dust pattern speed, but that this m

yes, in a
statistical
sense

H. SALO¹, E. LAURIKAINEN^{1,2}, R. BUTA³, AND J. H. KNAPEN^{4,5}

¹ Department of Physics/Astronomy Division, University of Oulu, FI-90014, Finland

² Finnish Centre for Astronomy with ESO (FINCA), University of Turku, Väisäläntie 20, FI-21500 Piikkiö, Finland

³ Department of Physics and Astronomy, University of Alabama, Box 870324, Tuscaloosa, AL 35487, USA

⁴ Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

⁵ Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain

Received 2010 February 3; accepted 2010 April 20; published 2010 May 3

ABSTRACT

Recently, Buta et al. examined the question “Do Bars Drive Spiral Density Waves?”, an idea supported by theoretical studies and also from a preliminary observational analysis. They estimated maximum bar strengths Q_b , maximum spiral strengths Q_s , and maximum $m = 2$ arm contrasts A_{2s} for 23 galaxies with deep Anglo-Australian Telescope (AAT) K_s -band images. These were combined with previously published Q_b and Q_s values for 147 galaxies from the Ohio State University Bright Spiral Galaxy Survey (OSUBSGS) sample and with the 12 galaxies from Block et al. Weak correlation between Q_b and Q_s was confirmed for the combined sample, whereas the AAT subset alone showed no significant correlations between Q_b and Q_s , nor between Q_b and A_{2s} . A similar negative result was obtained in Durbala et al. for 46 galaxies. Based on these studies, the answer to the above question remains uncertain. Here we use a novel approach, and show that although the correlation between the *maximum* bar and spiral parameters is weak, these parameters do correlate when compared *locally*. For the OSUBSGS sample, a statistically significant correlation is found between the local spiral amplitude, and the forcing due to the bar’s potential at the same distance, out to ≈ 1.6 bar radii (the typical bar perturbation is then of the order of a few percent). Also for the sample of 23 AAT galaxies of Buta et al., we find a significant correlation between local parameters out to ≈ 1.4 bar radii. Our new results confirm that, at least in a statistical sense, bars do indeed drive spiral density waves.