











Chemical-series adaptation | all assays





Acquisition | equal assay averages




One free worse-half hit; ties qualify. Every curve requires all configured assays and folds.

Relative performance on series with at least 15 compounds







Equal assay averages; NLL at five measured compounds. Descriptive comparisons, without significance claims.

Cellular clearance | prediction









Five-fold held-out predictions; pools ≥ 15 ; fixed hidden queries. Uncalibrated native mixture intervals.

Cellular clearance | acquisition




One free worse-half hit; ties qualify. Random curve is an exact expectation. No hidden labels enter model calls.

Efflux ratio | prediction









Efflux ratio | acquisition







One free worse-half hit; ties qualify. Random curve is an exact expectation. No hidden labels enter model calls.


In vivo clearance | prediction







In vivo clearance | acquisition



Microsomal clearance | prediction









Microsomal clearance | acquisition







One free worse-half hit; ties qualify. Random curve is an exact expectation. No hidden labels enter model calls.

Permeability | prediction




Permeability | acquisition




One free worse-half hit; ties qualify. Random curve is an exact expectation. No hidden labels enter model calls.

Plasma protein binding | prediction



Five-fold held-out predictions; pools ≥ 15 ; fixed hidden queries. Uncalibrated native mixture intervals.

Plasma protein binding | acquisition



One free worse-half hit; ties qualify. Random curve is an exact expectation. No hidden labels enter model calls.

Coverage and interpretation

Only complete endpoint/method fold sets enter the plotted summaries.

All-assay averages additionally require all 6 configured assays at every plotted position.

ALPaCA without offset reuses the selected offset-model hyperparameters and trains fresh weights.

R^2 is a weighted pooled statistic; Spearman is an average of within-series rankings.

NLL evaluates the native predictive mixture, while EI uses its Gaussian moments.

These folds have been explored repeatedly. The charts are descriptive, not fresh external validation.

Unavailable: in_vivo_clearance / maml (4 complete folds)

Unavailable: protein_binding / maml (1 complete folds)

Unavailable: cellular_clearance / maml (3 complete folds)