

# EOSC 213 - Quiz

---

## Instructions (20 points in total)

- Read the examination before beginning.
  - Calculators are allowed (if you don't have one, just give the expression to type in a calculator).
  - You have exactly 30 minutes for the examination.
  - Be as precise and clear as possible.
  - This is a closed book examination.
  - If you get stuck, make an assumption, state what it is and try to carry on.
- 

## Question 1: Taylor's approximation

Let us consider the exponential function 2

$$f(x) = \exp(x). \quad (1)$$

We are going to try to compute its derivative at  $x = 0$  using backward, forward and centered differences. You can use a **calculator** to compute the values of  $f(x)$ .

**Q1a** Compute the derivative of the exponential function at  $x = 0$  ( $f'(0)$ ) with a **first order** approximation **and** assess your error.

**Q1b** Compute the derivative of the exponential function at  $x = 0$  ( $f'(0)$ ) with a **second order** approximation **and** assess your error.

The Taylor series of the exponential function is:

$$\begin{aligned} f(x) &= 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + \dots \\ &= \sum_{n=0}^{\infty} \frac{x^n}{n!} \end{aligned} \quad (2)$$

**Q1c** Write the second order approximation of the exponential function and write the truncation error.

### Question 2: Diffusion

The 1D transient diffusion equation with homogeneous diffusion coefficient can be written as:

$$\frac{\partial c}{\partial t} = D \frac{\partial^2 c}{\partial x^2}. \quad (3)$$

Let us consider the physical domain  $x \in [-1; 1]$  m, with the specified boundary conditions  $x(1) = x(-1) = 0$ . Consider the following function:


$$c(x, t) = c_0 \cos\left(\frac{\pi x}{2}\right) \exp(-\alpha t) \quad (4)$$

**Q2a** Show that the function described in equation 4 solves the diffusion problem described by equation 3 [2 points]

**Q2b** Can you identify the value of  $\alpha$  [2 points]

**Q2c** Show that the function described in equation 4 satisfies the boundary conditions at all times [2 points]

**Q2d** Concentrations were measured at three different times and are represented in the graph below. The messy person who did the measurements does not remember at which time these measurements were taken. Can you help him, using your physical intuition? Give the temporal sequence of the 4 curves. [2 points]



**Q2e** What can you say about the mass evolution in the system? Is that consistent with the boundary conditions?

**Q2f** Use your physical reasoning (and equation 4) to describe the asymptotic/final solution.

**Question 3: Conservation equation** The continuity equation says:

$$\frac{\partial \text{Stuff}}{\partial t} + \text{div } \vec{j} = 0 \quad (5)$$

**Q3a** Use this equation to describe the advective movement of solutes

## Python Questions

This is a question about the following code:

---

```
class Problem_Def:
    """
    this class holds the specification for the domain,
    including the value of the porosity
    """

    def __init__(self, nx, ny, poro, wx, wy):
        self.nx = nx
        self.ny = ny
        self.poro = poro
        self.wx = wx
        self.wy = wy

def get_spacing(nx=4, ny=3, poro=0.4, wx=10, wy=20):
    the_prob = Problem_Def(nx, ny, poro, wx, wy)
    delx = the_prob.wx / the_prob.nx
    dely = the_prob.wy / the_prob.ny
    return delx, dely
```

---

**Q9** Given the code above, what does the following python statement print? [2 points]

```
print(f"{get_spacing(nx=6)}")
```

**Q10** modify the Problem\_Def class to incorporate get\_spacing as an instance method [2 points]

That is, create a version of Problem\_Def for which the following will work::

```
the_instance = Problem_Def()
delx, dely = the_instance.get_spacing()
```

where the new constructor has the signature::

```
def __init__(self, nx=4, ny=3, poro=0.4, wx=10, wy=20):
    ...
```