

Lecture 13

Properties of the Fourier transform

Preview of today's lecture

◆ Fourier transform properties

- ✦ Understand important connections between signal operations in the time and frequency domains
- ✦ Use the table of common transform pairs and the table of common properties to compute transforms without doing integration

◆ Fourier symmetry properties

- ✦ Connect signal properties like symmetric, even and odd in both time and frequency domains

Summarizing the Fourier transform and its inverse

Fourier transform (analysis)

$$X(j\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$$

Inverse Fourier transform (synthesis)

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega$$

$$x(t) \leftrightarrow X(j\omega)$$


$x(t)$
↑
domain
time

$X(j\omega)$

↑
domain
freq

Connections back to ECE 45

Lectures 2 - 3 working with signals



Lectures 4 - 7 LTI systems in the time domain

filtering *Bode plots, filtering*

Lectures 11, 17 LTI systems in the frequency domain



Lectures 8 - 10 Fourier series

Lectures 11 - 16 Fourier transform

Fourier

Essential Fourier transforms

Key points

- Know these important FT pairs

Basic Fourier transform pairs 1/3

(more pairs in the book)

	Time domain $x(t)$	Frequency domain $X(j\omega)$
Delta	$\delta(t)$	2π
Constant	$\frac{1}{2\pi}$	$\delta(\omega)$
Complex sinusoid	$\frac{e^{j\omega_0 t}}{2\pi}$	$\delta(\omega - \omega_0)$
Causal exponential	$e^{-at}u(t)$ $\text{Re}\{a\} > 0$	$\frac{1}{a + j\omega}$

RC circuit $e^{-t/RC}u(t)$

DC
0 rad/s

(more pairs in the book)

Basic Fourier transform pairs 2/3

	Time domain $x(t)$	Frequency domain $X(j\omega)$
Cosine	$\cos \omega_0 t$	$\pi \delta(\omega - \omega_0) + \pi \delta(\omega + \omega_0)$
Sine	$\sin \omega_0 t$	$\pi j (\delta(\omega + \omega_0) - \delta(\omega - \omega_0))$
Periodic signal w/ period T	$x(t)$	$2\pi \sum_{k=-\infty}^{\infty} a_k \delta(\omega - k\omega_0)$

$$\omega_0 = \frac{2\pi}{T}$$

\rightarrow Number
 FS coefficients
 domain variable

(more pairs in the book)

Basic Fourier transform pairs 3/3

	Time domain $x(t)$	Frequency domain $X(j\omega)$
Rectangle	$\text{rect}(t)$	$\frac{\sin \frac{\omega}{2}}{\frac{\omega}{2}} = \text{sinc} \left(\frac{\omega}{2\pi} \right)$
Scaled rectangle	$\text{rect} \left(\frac{t}{2T_1} \right)$	$2T_1 \frac{\sin(\omega)}{\omega} = 2T_1 \text{sinc} \left(\frac{\omega T_1}{\pi} \right)$
Sinc	$\text{sinc}(t)$	$\text{rect} \left(\frac{\omega}{2\pi} \right)$
Scaled sinc	$\frac{B}{2\pi} \text{sinc} \left(\frac{Bt}{2\pi} \right)$	$\text{rect} \left(\frac{\omega}{B} \right)$

Fourier transform properties

Key points

- Use FT properties to simplify calculation & build intuition
- Analyze problems that include FT properties

Fourier transform properties I

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega) \quad y(t) \xleftrightarrow{\mathcal{F}} Y(j\omega)$$

	Time domain	Fourier transform
Linearity	$ax(t) + by(t)$	$aX(j\omega) + bY(j\omega)$
Time shifting	$x(t - t_0)$	$e^{-j\omega t_0} X(j\omega)$
Differentiation	$\frac{dx}{dt}$	$j\omega X(j\omega)$
Integration	$\int_{-\infty}^t x(\tau) d\tau$	$\frac{1}{j\omega} X(j\omega) + \pi X(0) \delta(j\omega)$

Fourier transform properties 2

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

	Time domain	Fourier transform
Time scaling	$x(at)$	$\frac{1}{ a } X\left(\frac{j\omega}{a}\right)$
Frequency scaling	$\frac{1}{ b } x\left(\frac{t}{b}\right)$	$X(jb\omega)$
Frequency shifting	$x(t)e^{j\omega_0 t}$	$X(j(\omega - \omega_0))$
Parseval's theorem	$\int_{-\infty}^{\infty} x(t) ^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) ^2 d\omega$	

Fourier transform properties 3

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega) \quad y(t) \xleftrightarrow{\mathcal{F}} Y(j\omega)$$

$$h(t) \xleftrightarrow{\mathcal{F}} H(j\omega)$$

	Time domain	Fourier transform
Convolution in time	$y(t) = h(t) * x(t)$	$Y(j\omega) = H(j\omega)X(j\omega)$
Multiplication in time	$y(t) = h(t)x(t)$	$Y(j\omega) = \frac{1}{2\pi} \int_{-\infty}^{\infty} H(j\theta)X(j(\omega - \theta))d\theta$

scaling

convolution

These last two properties are important and feature in future lectures

Linearity

◆ If $x(t) \xleftrightarrow{\mathcal{F}} X(j\omega), \quad y(t) \xleftrightarrow{\mathcal{F}} Y(j\omega)$

◆ Then $ax(t) + by(t) \leftrightarrow aX(j\omega) + bY(j\omega)$

$$\int_{-\infty}^{\infty} (ax(t) + by(t)) e^{j\omega t} dt = \int_{-\infty}^{\infty} ax(t) e^{j\omega t} dt + \int_{-\infty}^{\infty} by(t) e^{j\omega t} dt$$

$$= aX(j\omega) + bY(j\omega)$$

Sums in **time** lead to sums in **frequency**

Linearity example

◆ Consider

$$\cos t \xleftrightarrow{\mathcal{F}} \pi[\delta(\omega - 1) + \delta(\omega + 1)]$$

$$\sin t \xleftrightarrow{\mathcal{F}} \pi j[\delta(\omega + 1) - \delta(\omega - 1)]$$

Table

◆ By linearity

$$\cos t + j \sin t \xleftrightarrow{\mathcal{F}} \underbrace{\pi\delta(\omega - 1) + \pi\delta(\omega + 1)}_{0} - \underbrace{\pi\delta(\omega + 1) + \pi\delta(\omega - 1)}_{0}$$

$$\xleftrightarrow{\mathcal{F}} 2\pi\delta(\omega - 1)$$

$$\xleftrightarrow{\mathcal{F}} \mathcal{F}\{e^{jt}\}$$

$$j \sin t \xleftrightarrow{\mathcal{F}} [\pi\delta(\omega - 1) - \pi\delta(\omega + 1)]$$

what is frequency? $\omega = 1$

e^{jt}

Time shifting

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then

$$x(t - t_0) \xleftrightarrow{\mathcal{F}} e^{-j\omega t_0} X(j\omega)$$


◆ Time shifting does not change the magnitude in the freq. domain

$$|X(j\omega)e^{-j\omega t_0}| = |X(j\omega)| |e^{-j\omega t_0}|$$

◆ Phase changes are linear with frequency ω and shift t_0

$$\angle(X(j\omega)e^{-j\omega t_0}) = \angle X(j\omega) - \omega t_0$$

Shift in time leads to linear phase shift in frequency



Very little
new work
required!


$$\text{rect}(t) \leftrightarrow \text{sinc}(\omega/2\pi)$$

$$\text{rect}(t - 1/2) \leftrightarrow e^{-j\omega/2} \text{sinc}(\omega/2\pi)$$

$$\Rightarrow e^{-j\omega/2} \text{sinc}(\omega/2\pi)$$

Time shifting example

- ◆ Find the Fourier transform of



- ◆ This signal is just a shifted rectangle function $\text{rect}(t - 1/2)$
- ◆ Using the time shifting property

$$\text{rect}(t - 1/2) \xleftrightarrow{\mathcal{F}} e^{-j\omega/2} \text{sinc}\left(\frac{\omega}{2\pi}\right)$$

Differentiation

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then

$$\frac{dx}{dt} \xleftrightarrow{\mathcal{F}} j\omega X(j\omega)$$

◆ Proof

$$\frac{d}{dt} x(t) = \frac{d}{dt} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega \right)$$

$$\underbrace{\frac{dx}{dt}}_{\text{new func}} = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) \frac{d}{dt} (e^{j\omega t}) d\omega$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} \underbrace{j\omega X(j\omega)}_{\text{new FT}} e^{j\omega t} d\omega$$

Swap
derivative
and
integral

importance of $X(j\omega)$ grows w/ ω

Differentiation example

$$\begin{aligned} x(t) &\leftrightarrow X(j\omega) \\ y(t) &\leftrightarrow Y(j\omega) \end{aligned}$$

- ◆ What is the FT of the system characterized by

$$\frac{dy}{dt} + ay(t) = x(t)$$

- ◆ Solution:

$$j\omega Y(j\omega) + aY(j\omega) = X(j\omega)$$

- ★ Take FT of both sides

$$\begin{aligned} e^{-at}u(t) \\ \text{Re}\{a\} > 0 \end{aligned}$$

$$j\omega Y(j\omega) + aY(j\omega) = X(j\omega)$$

$$(j\omega + a)Y(j\omega) = X(j\omega)$$

- ★ Therefore

Freq response
of System

$$H(j\omega) = \frac{Y(j\omega)}{X(j\omega)} = \frac{1}{j\omega + a}$$

recall
table

$$h(t) = e^{-at}u(t)$$

Integration

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then

$$\int_{-\infty}^t x(\tau) d\tau \xleftrightarrow{\mathcal{F}} \frac{1}{j\omega} X(j\omega) + \pi X(0) \delta(\omega)$$

DC component

running
"Sum"

Time scaling

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$


◆ Then

$$x(at) \xleftrightarrow{\mathcal{F}} \frac{1}{|a|} X\left(\frac{j\omega}{a}\right)$$

Handwritten notes: A blue arrow points from the a in $x(at)$ to the a in the denominator of the frequency term $\frac{j\omega}{a}$. The word "note" is written in blue above the arrow. The word "scaling" is written in blue below the $|a|$ term.

Time **expansion** $|a| < 1$ leads to frequency **compression**

Time **compression** $|a| > 1$ leads to frequency **expansion**




$$\longleftrightarrow e^{j\omega/2} \text{sinc}(\omega/2\pi)$$



$$2 \cdot 1/2 = 1$$

$$\longleftrightarrow \frac{1}{2} e^{j\omega/4} \text{sinc}(\omega/4\pi)$$



$$\longleftrightarrow 2 e^{-j\omega/2} \text{sinc}\left(\frac{\omega \cdot 2}{2\pi}\right)$$

$$\rightarrow 2 e^{j\omega} \text{sinc}(\omega/\pi)$$

Frequency scaling

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then


$$\frac{1}{|b|} x\left(\frac{t}{b}\right) \xleftrightarrow{\mathcal{F}} X(jb\omega)$$

Frequency **expansion** $|b| < 1$ leads to time **compression**


Frequency **compression** $|b| > 1$ leads to time **expansion**

Shift and scaling example

- ◆ Find the FT of




- ◆ To start, notice that




- ◆ Shifting gives our function $q(t) = \text{rect}\left(\frac{t-1}{2}\right) = \text{rect}\left(\frac{t}{2} - \frac{1}{2}\right)$

$$p(t) = \text{rect}(t - 1/2) \Leftrightarrow \text{sinc}(w/2\pi)$$



#1


$$g(t) = p(t/2) = \text{rect}(t/2 - 1/2)$$



$$= \text{rect}\left(\frac{t-1}{2}\right)$$

#2


$\text{rect}(t/2)$



$$\Leftrightarrow 2 \text{sinc}\left(\frac{w \cdot 2}{2\pi}\right)$$

$$\rightarrow 2 \text{sinc}(w/\pi)$$

Approach 2




① $p(t) \leftrightarrow e^{-j\omega t/2} \text{sinc}(\omega/2\pi)$

② $p(t/2) \leftrightarrow 2 \cdot p(j\omega \cdot 2)$

$\rightarrow 2e^{-j\omega} \text{sinc}\left(\frac{\omega \cdot 2}{2\pi}\right)$

$g(t) \leftrightarrow 2e^{-j\omega} \text{sinc}(\omega/\pi)$

Approach 2

$$g(t) = p(t/2) = \text{rect}(t/2 - 1/2) \\ = \text{rect}\left(\frac{t-1}{2}\right)$$


1)

2 2

$$\textcircled{1} \text{ rect}(t/2) \leftrightarrow 2 \text{ sinc}(2 \cdot \bar{w}/2\pi) \\ \text{rect}(t) \leftrightarrow 2 \text{ sinc}(w/\pi)$$

$$\textcircled{2} g(t) = \text{rect}\left(\frac{t-1}{2}\right) = r(t-1)$$

$$\leftrightarrow e^{-j\omega} R(j\omega)$$

$$\leftrightarrow 2 e^{-j\omega} \text{sinc}(w/\pi)$$

Shift and scaling example (continued)

- ◆ From the scaling property

$$\text{rect}\left(\frac{t}{2}\right) \xleftrightarrow{\mathcal{F}} 2\text{sinc}\left(\frac{2\omega}{2\pi}\right) = 2\text{sinc}\left(\frac{\omega}{\pi}\right)$$

- ◆ From the shift property

$$\text{rect}\left(\frac{t-1}{2}\right) \xleftrightarrow{\mathcal{F}} e^{-j\omega} 2\text{sinc}\left(\frac{\omega}{\pi}\right)$$

✖ Another scaling example

◆ What is the inverse Fourier transform of $\text{sinc}(\omega)$?

◆ We know that

★ From the rect-sinc Fourier pair

$$\text{rect}(t) \xleftrightarrow{\mathcal{F}} \text{sinc}\left(\frac{\omega}{2\pi}\right)$$

★ From the scaling law

$$x(at) \xleftrightarrow{\mathcal{F}} \frac{1}{|a|} X\left(\frac{j\omega}{a}\right) \quad a = 1/2\pi$$

◆ Using the scaling property

$$\text{rect}(t/2\pi) \xleftrightarrow{\mathcal{F}} 2\pi \text{sinc}(\omega)$$

◆ Therefore using linearity

$$\frac{1}{2\pi} \text{rect}(t/2\pi) \xleftrightarrow{\mathcal{F}} \text{sinc}(\omega)$$

$$\frac{1}{15} x(t/5) \Leftrightarrow X(j\omega 5)$$

Inversion (time reversal)

◆ Find the FT of $x(-t)$

◆ This is just a special case of time and frequency scaling

$$x(at) \xleftrightarrow{\mathcal{F}} \frac{1}{|a|} X\left(\frac{j\omega}{a}\right)$$

◆ With $a = -1$

$$x(-t) \xleftrightarrow{\mathcal{F}} X(-j\omega)$$

Example using scaling and time shift

- ◆ Determine the Fourier transform of $\text{sinc}(1 - 2t)$

①

$$x(t) = \text{sinc}(1 - 2t)$$

$$= y(2t)$$

②

$$y(t) = \text{sinc}(1 - t)$$

$$= \text{sinc}(-(t - 1))$$

$$= z(t - 1)$$

$$z(t) = \text{sinc}(-t)$$

$$= \text{sinc}(t)$$

Symmetry

$$\text{rect}(t) \xleftrightarrow{\mathcal{F}} \text{sinc}(\omega/2\pi)$$

$$\text{sinc}(t/2\pi) \xleftrightarrow{\mathcal{F}} 2\pi \text{rect}(-\omega) = 2\pi \text{rect}(\omega)$$

$$\text{sinc}(t) \xleftrightarrow{\mathcal{F}} \text{rect}(\omega/(2\pi))$$

$$Z(j\omega) = \text{rect}(\omega/(2\pi))$$

$$Y(j\omega) = e^{-j\omega} Z(j\omega)$$

$$X(j\omega) = \frac{1}{2} Y\left(j\frac{\omega}{2}\right)$$

$$= \frac{1}{2} e^{-j\omega/2} \text{rect}(\omega/(4\pi))$$

Scaled

Time sh

Scale

Frequency shifting

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then

$$x(t)e^{j\omega_0 t} \xleftrightarrow{\mathcal{F}} X(j(\omega - \omega_0))$$

this is called **modulation**

◆ Corollary

$$x(t) \cos \omega_0 t \xleftrightarrow{\mathcal{F}} \frac{1}{2} X(j(\omega - \omega_0)) + \frac{1}{2} X(j(\omega + \omega_0))$$

↑ right ↑ left

constant

Modulate in time leads to shift in frequency

Example combining shift and scaling

- ◆ Determine the inverse Fourier transform of

$$X(j\omega) = \frac{2 \sin(3(\omega - 2\pi))}{(\omega - 2\pi)}$$

Example (continued)

- ◆ Given

$$X(j\omega) = \frac{2 \sin(3(\omega - 2\pi))}{(\omega - 2\pi)}$$

- ◆ Use the following fact

$$\text{rect}(t) \xleftrightarrow{\mathcal{F}} \text{sinc}\left(\frac{\omega}{2\pi}\right) = \frac{\sin(\omega/2)}{\omega/2}$$

- ◆ For convenience let

$$R(j\omega) = \frac{\sin(\omega/2)}{\omega/2}$$

- ◆ Rewrite as another shifted function

$$X(j\omega) = Y(j(\omega - 2\pi))$$

$$Y(j\omega) = \frac{2 \sin(3\omega)}{\omega}$$


$\text{sinc}(x) = \frac{\sin(\pi x)}{\pi x}$

rewrite

Example (continued)

- ◆ Rewrite again as


$$Y(j\omega) = 3 \frac{2 \sin(6\omega/2)}{6\omega/2}$$

$$= 6R(j6\omega)$$


- ◆ Using the scaling property

$$Y(j\omega) = 6R(j6\omega) \quad \longrightarrow \quad y(t) = \text{rect}(t/6)$$


- ◆ Using the shift property

$$X(j\omega) = Y(j(\omega - 2\pi)) \quad \longrightarrow \quad x(t) = e^{j2\pi t} \text{rect}(t/6)$$


Parseval's theorem

$$\int_{-\infty}^{\infty} |x(t)|^2 dt = \left[\frac{1}{2\pi} \right] \int_{-\infty}^{\infty} |X(j\omega)|^2 d\omega$$

Energy of the signal in
the time domain

Energy of the signal in
the frequency domain

- ◆ This is a result of conservation of energy
- ◆ Scaling factor is because of radians

Example using Parseval's theorem

- ◆ If the signal $x(t)$ has the FT below $X(j\omega)$

$$x(t) = t \left(\frac{\sin t}{\pi t} \right)^2 \quad X(j\omega) = \begin{cases} \frac{j}{2\pi}, & -2 \leq \omega < 0 \\ -\frac{j}{2\pi}, & 0 \leq \omega \leq 2 \\ 0, & \text{otherwise} \end{cases}$$

- ◆ Calculate



$$A = \int_{-\infty}^{\infty} t^2 \left(\frac{\sin t}{\pi t} \right)^4 dt$$

$x^2(t) dt$

Example with Parseval's theorem (cont.)

$$\int_{-\infty}^{\infty} t^2 \left(\frac{\sin(t)}{\pi t} \right)^4 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} |X(j\omega)|^2 d\omega$$

$$= \frac{1}{2\pi^3}$$



Duality in the Fourier transform

◆ If

$$x(t) \xleftrightarrow{\mathcal{F}} X(j\omega)$$

◆ Then

$$X(jt) \xleftrightarrow{\mathcal{F}} 2\pi x(-\omega)$$

Pay attention to j

If you know one Fourier pair then you know the other Fourier pair

$$\text{rect}(t) \leftrightarrow \text{sinc}(t/2\pi)$$

$$\text{sinc}(t/2\pi) \leftrightarrow 2\pi \text{rect}(-t)$$

$$\leftrightarrow 2\pi \text{rect}(t)$$

Applications of duality

◆ Reproving frequency shift

★ Consider
$$x(t - t_0) \xleftrightarrow{\mathcal{F}} e^{-j\omega t_0} X(j\omega)$$

★ Then
$$x(t)e^{j\omega_0 t} \xleftrightarrow{\mathcal{F}} X(j(\omega - \omega_0))$$

◆ Impulse in time and frequency

★ Consider
$$\delta(t) \xleftrightarrow{\mathcal{F}} 1$$

★ Then
$$1 \xleftrightarrow{\mathcal{F}} 2\pi\delta(\omega)$$

Duality example

◆ Consider

$$\text{rect}(t) \xleftrightarrow{\mathcal{F}} \text{sinc}\left(\frac{\omega}{2\pi}\right)$$

◆ Then

$$\begin{aligned} \text{sinc}\left(\frac{t}{2\pi}\right) &\xleftrightarrow{\mathcal{F}} 2\pi \cdot \text{rect}(-\omega) \\ &= 2\pi \cdot \text{rect}(j\omega)x(t - t_0) \end{aligned}$$

Since rect is an even function

Symmetry, even, and odd

Key points

- Connect signal properties like symmetric, even and odd in both time and frequency domains

Even and odd

- ◆ Can decompose a signal into even and odd components

$$x(t) = e(t) + o(t)$$

- ◆ Even part is *(conjugate symmetric)*

$$e(t) = \frac{1}{2}(x(t) + x^*(-t))$$

- ◆ Odd part is *(conjugate asymmetric)*


$$o(t) = \frac{1}{2}(x(t) - x^*(-t))$$

- ◆ Can similarly decompose

$$X(j\omega) = E(j\omega) + O(j\omega)$$

Connecting the properties

$$x(t) = \text{Re}\{e(t)\} + j\text{Im}\{e(t)\} + \text{Re}\{o(t)\} + j\text{Im}\{o(t)\}$$



swap!

$$X(j\omega) = \text{Re}\{E(j\omega)\} + j\text{Im}\{E(j\omega)\} + \text{Re}\{O(j\omega)\} + j\text{Im}\{O(j\omega)\}$$


◆ Implications

★ Conjugate symmetry $x(t)$ is real $\rightarrow X(j\omega) = X^*(-j\omega)$

★ Real signals are even in amplitude since $|X(j\omega)| = |X(-j\omega)|$

Even / odd example

0.5c
odd



✓
A real signal $x(t)$ has the graph above

$$x(t) = \text{Re}\{e(t)\} + j\text{Im}\{e(t)\} + \text{Re}\{o(t)\} + j\text{Im}\{o(t)\}$$

real

$$X(j\omega) = \text{Re}\{E(j\omega)\} + j\text{Im}\{E(j\omega)\} + \text{Re}\{O(j\omega)\} + j\text{Im}\{O(j\omega)\}$$

Answer the following questions:

$$\text{Re}\{X(j\omega)\} = 0 \quad \text{Yes}$$


$$\text{Im}\{X(j\omega)\} = 0 \quad \text{No}$$

$$x(t) = e(t) + o(t)$$

$$= \text{Re}\{e(t)\} + j\text{Im}\{e(t)\} + \text{Re}\{o(t)\} + j\text{Im}\{o(t)\}$$

Summarizing symmetry

$$x(t) = \text{Re}\{e(t)\} + j\text{Im}\{e(t)\} + \text{Re}\{o(t)\} + j\text{Im}\{o(t)\}$$



$$X(j\omega) = \text{Re}\{E(j\omega)\} + j\text{Im}\{E(j\omega)\} + \text{Re}\{O(j\omega)\} + j\text{Im}\{O(j\omega)\}$$

- ◆ Key symmetry equation relates real, imaginary, even, and odd in the time and frequency domains
- ◆ Can determine signal characteristics in one domain by inspecting the other domain