

# Chapter 4 Problem 22

Alan Arnholt

March 27, 2014

## Problem 22

Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} F$  with corresponding pdf  $f(x) = 3x^2, 0 \leq x \leq 1$ .

- (a) Find the pdf for  $X_{\min}$ .
- (b) Find the pdf for  $X_{\max}$ .
- (c) If  $n = 10$ , find the probability that the largest value,  $X_{\max}$ , is greater than 0.92
- (d) if  $n = 10$ , find the expected value of  $X_{\max}$ .

(a) The pdf for  $X_{\min}$  is:

$$f_{\min}(x) = n(1 - F(X))^{n-1} f(x) \quad (1)$$

Since  $F(x) = \int_0^x 3t^2 dt = x^3, 0 \leq x \leq 1$ , it follows that

$$f_{\min}(x) = n(1 - x^3)^{n-1} 3x^2, 0 \leq x \leq 1$$

(b) The pdf for  $X_{\max}$  is:

$$f_{\max}(x) = nF^{n-1}(x)f(x) \quad (2)$$

Since  $F(x) = \int_0^x 3t^2 dt = x^3, 0 \leq x \leq 1$ , it follows that

$$f_{\max}(x) = n(x^3)^{n-1}3x^2, 0 \leq x \leq 1$$

$$P(X_{\max} > 0.92)$$

(c)

$$f_{\max}(x) = 10(x^3)^{10-1} 3x^2 = 30x^{29}, 0 \leq x \leq 1$$

```
f <- function(x){30*x^29}
ans <- integrate(f, 0.92, 1)$value
ans
```

```
[1] 0.918
```

$$P(X_{\max} > 0.92) = \int_{0.92}^1 30x^{29} dx = 0.918.$$

$$E[X_{\max}] = \int_0^1 x_{\max} f_{\max}(x) = \int_0^1 30x^{30}$$

```
f1 <- function(x){30*x^30}  
ans <- integrate(f1, 0, 1)$value  
ans
```

```
[1] 0.9677
```

$$E[X_{\max}] = 0.9677$$

# What does the pdf look like?

```
curve(f, 0, 1, n = 500, col = "purple")  
abline(v = ans, lty = "dashed", col = "red")
```

