

Monopoly Problems

Andrew Ye and Diva Shah

May 2024

Contents

1	Unit 1: Atomic Structure and Properties	3
2	Unit 2: Molecular and Ionic Compound Structure and Properties	4
3	Unit 3: Intermolecular Forces and Properties	6
4	Unit 4: Chemical Reactions	7
5	Unit 5: Kinetics	7
6	Answers	9
6.1	Unit 1	9
6.2	Unit 2	10
6.3	Unit 3	11
6.4	Unit 4	12

1 Unit 1: Atomic Structure and Properties

Problem 1

Calculate the number of moles in a 7.89kg sample of $\text{C}_9\text{H}_8\text{O}_4$

Problem 2

Given this graph, what is true about the element depicted



- (a) In an average sample of the element, less than 20% of the atoms have an atomic mass of $66u$.
- (b) The most abundant isotope of the element has an atomic mass of $64u$.
- (c) The element has an average atomic mass of $64u$.
- (d) The element has an average atomic mass between 66 and $68u$.

Problem 3

What is the percent composition of Carbon in $\text{C}_{13}\text{H}_{18}\text{O}_2$?

Problem 4

A compound contains 32.38% sodium, 22.65% sulfur, and 44.99% oxygen. What is the empirical formula.

Problem 5

What is the full electron configuration of mercury?

Problem 6

Below, the photoelectron spectra of the 2s electrons of Be and Mg are shown.



Is peak *X* the peak associated with Be or Mg?

Problem 7

What are the periodic trends of ionization energy, atomic radius, and electronegativity? Why?

2 Unit 2: Molecular and Ionic Compound Structure and Properties

Problem 8

Which of the following bonds is likely to have the most ionic character?

- (a) H — F
- (b) C — O
- (c) Na — F
- (d) Mg — O

Problem 9

Based on the information in the table, which of the following arranges the bonds in order of decreasing polarity?

Element	Electronegativity
H	2.2
N	3.0
F	4.0
Cl	3.2
Se	2.6
I	2.7

- (a) $\text{Se} - \text{N} > \text{H} - \text{I} > \text{Cl} - \text{F}$
 (b) $\text{H} - \text{I} > \text{Se} - \text{N} > \text{Cl} - \text{F}$
 (c) $\text{Cl} - \text{F} > \text{H} - \text{I} > \text{Se} - \text{N}$
 (d) $\text{Cl} - \text{F} > \text{Se} - \text{N} > \text{H} - \text{I}$

Problem 10

Why is the lattice energy of CsF smaller than the lattice energy of KF?

Problem 11

What type of structure do metallic elements form and through what bonds?

Problem 12

What are the two types of metallic alloys and what are their differences?

Problem 13

Draw a Lewis Diagram for Acetic Acid CH_3COOH .

Problem 14

Draw the Lewis Diagram for CO_2

Problem 15


Draw the Lewis Diagram(s) for ozone, O_3

Problem 16

Write the formal charges for all three molecules above.

Problem 17


What is the electron geometry, molecular geometry, and hybridization of the central atom in this molecule.




3 Unit 3: Intermolecular Forces and Properties

Problem 18

What are the intermolecular forces present among these molecules.



Problem 19



What phase transition is this?

Problem 20

Originally, a sample of gas is in a rigid container at $299K$ and $0.70atm$. The student increases the temperature of the $CO_2(g)$ in the container to $425K$.

- What does raising the temperature do to the motion of the molecules?
- What is the pressure at $425K$?
- In terms of Kinetic Molecular Theory, why does the pressure of gas change as it is heated?

Problem 21

A $60.3g$ of $Be(OH)_2$ is dissolved in enough water to produce $1.75L$ of solution. Calculate the concentration of OH^- ions.

Problem 22

Describe the photoelectric effect.

4 Unit 4: Chemical Reactions

Problem 23

Balance this reaction: $\text{C}_5\text{H}_{10} + \text{O}_2 \longrightarrow \text{CO}_2 + \text{H}_2\text{O}$

Problem 24

Balance this redox reaction: $\text{MnO}_4^- + \text{I}^- \longrightarrow \text{I}_2 + \text{Mn}^{2+}$

Problem 25


Aqueous FeCl_3 reacts with KOH to produce a solid precipitate of $\text{Fe}(\text{OH})_3$ and aqueous KCl . What is the balanced net ionic equation?

Problem 26

What is the difference between physical changes and chemical changes?


Problem 27

H_2O and Fe are reacted according to the reaction below. There was initially 36.0g H_2O and 67.0g Fe . What is the limiting reactant, how much of the excess reactant will remain, and how much iron oxide is produced?



Problem 28

A 56kg sample of CO and 6.0kg sample of H_2 are combined into a closed vessel.




How many moles of $\text{CH}_3\text{OH}(\text{g})$ have been produced?

5 Unit 5: Kinetics

Problem 29

For this reaction:



What would be rate be in terms of each reactant and product.

CH_4 rate =

O_2 rate =

CO_2 rate =

H_2O rate =

Problem 30

If the rate of disappearance of CH_4 equals $5.0 \frac{M}{s}$ for the above reaction, what is the rate of appearance of H_2O ?

Problem 31

For the above reaction, what is the reaction rate if O_2 decreases from $0.1M$ to $0.04M$ in $125ms$?

Problem 32

$\text{A(aq)} + 2\text{B(aq)} \longrightarrow \text{Products}$

Experiment	$[\text{A}]_0$	$[\text{B}]_0$	Initial Rate
1	$0.10M$	$0.10M$	$1.0 \times 10^{-2} \frac{M}{s}$
2	$0.3M$	$0.10M$	$9.0 \times 10^{-2} \frac{M}{s}$
3	$0.3M$	$0.15M$	$9.0 \times 10^{-2} \frac{M}{s}$

What is the rate law?

Problem 33

N_2O_5 decomposes by a 1st order reaction with $k = 4.80 \times 10^{-4} \frac{1}{s}$. What is the concentration of N_2O_5 after 825 seconds if the initial concentration is $0.0165M$? What is the half-life for this reaction?

6 Answers

6.1 Unit 1

Problem 1

The molar mass of $C_9H_8O_4$ is $1.008 * 8 + 12.01 * 9 + 16.00 * 4 = 180.2 \frac{g}{mol}$

$$7.89kg \times \frac{1g}{10^{-3}kg} \times \frac{1mol}{180.2g} = 43.8mol \quad (1)$$

Problem 2

(b), the tallest peak of the graph is the one at $64u$.

Problem 3

In one mole of $C_{13}H_{18}O_2$ is $206.31g$.

$$1mol C_{13}H_{18}O_2 \times \frac{13mol C}{1mol C_{13}H_{18}O_2} \times \frac{12.01g}{1mol C} = 156.31g \quad (2)$$

Thus, the percent composition by weight is $\frac{156.31}{206.31} = 75.764\%$

Problem 4

Take $100g$ of the substance such that there are $32.38g$ sodium, $22.65g$ sulfur, and $44.99g$ oxygen.

$$\begin{aligned} 32.38g Na \times \frac{1mol Na}{22.99g} &= 1.408mol Na \\ 22.65g S \times \frac{1mol S}{32.07g} &= 0.7063mol S \\ 44.99g O \times \frac{1mol O}{16g} &= 2.812mol O \end{aligned} \quad (3)$$

Take the ratio of each compound with the smallest quantity.

$$\begin{aligned} S : \frac{0.7063}{0.7063} &= 1 \\ Na : \frac{1.408}{0.7063} &= 2 \\ O : \frac{2.812}{0.7063} &= 4 \end{aligned} \quad (4)$$

Therefore, the empirical formula is Na_2SO_4

Problem 5

$$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 5s^2 4d^{10} 5p^6 6s^2 4f^{14} 5d^{10}$$

Problem 6

Be. The peak location of the peak on the x-axis means that there is less binding energy for the electrons in element X. Be has fewer protons and both electrons are in the same shell, so its peak must belong to Be.

Problem 7

- The electronegativity increases from left to right across a period. This is because if a valence shell of electrons is less than half full then it requires less energy to lose an electron than gain one. If the valence shell of electrons is more than half full, it is easier to pull an electron into the valence shell. The electronegativity decreases from the top to the bottom of a group. This is because there is a greater atomic radius lower on the group.
- The ionization energy increases from left to right in a period. This is because of greater valence shell stability also because of smaller atomic radius. The ionization energy also decreases from top to bottom of a group. This is because of greater electron shielding and greater atomic radius.
- Atomic radius decreases from left to right within a period. This is because there are more protons to the right of the period. Atomic radius increases from top to bottom within a group. This is because of electron shielding and there are more electron shells in the atom.

6.2 Unit 2

Problem 8

The ionic character increases the greater the electronegativity difference. In this case, Na and O had the greatest electronegativity difference.

Problem 9

(c) $\text{Cl} - \text{F} > \text{H} - \text{I} > \text{Se} - \text{N}$

Problem 10

Cs^+ has a larger atomic radius than K^+ . So the distance between the cation and anion is greater than in CsF than in KF


Problem 11

Most metallic elements form crystalline solids at room temperature. Their bonds are metallic bonds due to electrostatic attraction between metal cations and delocalized electrons.


Problem 12

- Substitutional alloys. These alloys form when one atom of a similar size to the host metal replaces an atom of the host metal. The substitute atom must be of similar size. These alloys have good thermal and electrical conductivity.
- Interstitial alloys. These alloys are formed when smaller atoms fill in the gaps between the larger host atoms. This makes the metal harder and less malleable.


Problem 13



Problem 14




Problem 15



Problem 16

All formal charges of CH_3COOH and CO_2 are zero.



Problem 17

The electron geometry is tetrahedral. The molecular geometry is trigonal pyramidal. Hybridization of N atom is sp^3 since it has tetrahedral electron geometry.

6.3 Unit 3

Problem 18

Dipole-dipole and London dispersion forces. The $\text{C} - \text{O}$ bond is polar and the molecule is asymmetrical so it is polar. There are no $\text{H} - \text{F}$, $\text{H} - \text{O}$, or $\text{H} - \text{N}$ bonds, so there is no hydrogen bonding.

Problem 19

Condensation. Both have no regular arrangement, the one on the left is separated by the one on the right is close together, so the molecules are transitioning from gas to liquid.

Problem 20

(a) As you increase the temperature, the average kinetic energy or speed increases as well.

(b) $n_1 = n_2$ and $V_1 = V_2$, so $P_1V_1 = n_1RT_1$ and $P_2V_2 = n_2RT_2$. $\frac{P_2V_2}{RT_2} = \frac{P_1V_1}{RT_1} \implies P_2 = \frac{P_1T_2}{T_1} = \frac{0.7 \times 425}{299} = 0.99 \text{ atm}$


(c) As the temperature increases, the average kinetic energy increases, so the molecules undergo more collisions with the walls of the container.

Problem 21





$$60.3g \times \frac{1 \text{ mol Ba(OH)}_2}{171.35g} \times \frac{2 \text{ mol OH}^-}{1 \text{ mol Ba(OH)}_2} * \frac{1}{1.75L} = 0.402M$$

Problem 22






The photoelectric effect occurs when light of a certain minimum frequency/energy hits the surface of a metal and electrons are ejected.

6.4 Unit 4**Problem 23****Problem 24**


The oxidation reaction:




The reduction reaction:



Add the reactions together:



Problem 25



Problem 26

Chemical processes are characterized by changes in intramolecular forces, while physical processes are characterized by changes only in intermolecular forces.

Problem 27

Find the limiting reactant:

$$\begin{aligned} 36.0 \text{ gH}_2\text{O} * \frac{1 \text{ molH}_2\text{O}}{18.02 \text{ gH}_2\text{O}} * \frac{1 \text{ molFe}_3\text{O}_4}{4 \text{ molH}_2\text{O}} &= 0.49945 \text{ molFe}_3\text{O}_4 \\ 67.0 \text{ gFe} * \frac{1 \text{ molFe}}{55.85 \text{ gFe}} * \frac{1 \text{ molFe}_3\text{O}_4}{3 \text{ molFe}} &= 0.39988 \text{ molFe}_3\text{O}_4 \end{aligned} \quad (5)$$

Therefore, the limiting reactant is Fe_3

Find how much iron oxide is produced: Use the limiting reactant

$$0.39988 \text{ molFe}_3 * \frac{231.55 \text{ gFe}_3\text{O}_4}{1 \text{ molFe}_3} = 92.6 \text{ gFe}_3\text{O}_4 \quad (6)$$

Find how much excess reactant is left over:

$$67.0 \text{ gFe} * \frac{1 \text{ molFe}}{55.85 \text{ gFe}} * \frac{4 \text{ molH}_2\text{O}}{3 \text{ molFe}} * \frac{18.02 \text{ gH}_2\text{O}}{1 \text{ molH}_2\text{O}} = 28.8 \text{ gH}_2\text{O} \quad (7)$$

28.8 grams of water is used out of 32.0 grams. So there is 7.2 grams left over of the excess reagent.

Problem 28

$$1.5 \times 10^3 \text{ mol}$$

Problem 29

$$\begin{aligned} \text{CH}_4 \quad \text{rate} &= -\frac{\Delta[\text{CH}_4]}{\Delta t} \\ \text{O}_2 \quad \text{rate} &= -\frac{1}{2} \frac{\Delta[\text{O}_2]}{\Delta t} \\ \text{CO}_2 \quad \text{rate} &= \frac{\Delta[\text{CO}_2]}{\Delta t} \\ \text{H}_2\text{O} \quad \text{rate} &= \frac{1}{2} \frac{\Delta[\text{H}_2\text{O}]}{\Delta t} \end{aligned}$$

Problem 30

$$10 \frac{M}{s}$$

Problem 31

$$\begin{aligned} \frac{(0.04-0.1)M}{125ms} \times \frac{1ms}{10^{-3}s} &= -0.48 \frac{M}{s} \\ \text{rate} &= -\frac{1}{2} \times -0.48 = 0.24 \frac{M}{s} \end{aligned}$$

Problem 32

When concentration of B is held constant and the concentration of A is tripled, the initial rate is multiplied by 9 so the order with respect to A is 2. The order with respect to B is 0 because nothing changes when the concentration is increased. Thus the rate law is

$$\text{rate} = k[A]^2[B]^0 = k[A]^2$$

$$1.0 \times 10^{-2} = k \times 0.1^2 \implies k = 1.0 \frac{1}{ms}, \text{ thus } \text{rate} = [A]^2$$

Problem 33

$$\ln[A]_t = -kt + \ln[A]_0$$

$$\ln[A]_t = -(4.8 \times 10^{-4})(825) + \ln(0.0165)$$

$$[A]_t = e^{-4.50} = 0.0111M$$

Half-life:

$$\ln\left(\frac{1}{2}[A]_0\right) - \ln[A]_0 = -kt$$

$$\ln\left(\frac{1}{2}\right) = -kt$$

$$t = \frac{\ln(2)}{k}$$