Monopoly Problems

Andrew Ye and Diva Shah

May 2024

Contents

1 Unit 1: Atomic Structure and Properties

Problem 1

Calculate the number of moles in a $7.89kg$ sample of $C_9H_8O_4$

Problem 2

Given this graph, what is true about the element depicted

(a) In an average sample of the element, less than 20% of the atoms have an atomic mass of $66u$.
(b) The most abundant isotope of the element has an atomic mass of $64u$.
(c) The element has an average atomic mass of $64u$.
(d) The element has an average atomic mass between 66 and $68u$.

Problem 3

What is the percent composition of Carbon in $C_{13}H_{18}O_2$?

Problem 4

A compound contains 32.38% sodium, 22.65% sulfur, and 44.99% oxygen. What is the emperical forumula.

Problem 5

What is the full electron configuration of mercury?

Problem 6

Below, the photoelectron spectra of the $2s$ electrons of Be and Mg are shown.

Is peak X the peak associated with Be or Mg?

Problem 7

What are the periodic trends of ionization energy, atomic radius, and electronegativity? Why?

2 Unit 2: Molecular and Ionic Compound Structure and Properties

Problem 8

Which of the following bonds is likely to have the most ionic character?
(a) H——F
(b) C——O
(c) Na——F
(d) Mg——O

Problem 9

Based on the information in the table, which of the following arranges the bonds in order of decreasing polarity?

Element	Electronegativity
H	2.2
N	3.0
F	4.0
Cl	3.2
Se	2.6
I	2.7

(a) Se——N > H——I > Cl——F
(b) H——I > Se——N > Cl——F
(c) Cl——F > H——I > Se——N
(d) Cl——F > Se——N > H——I

Problem 10

Why is the lattice energy of CsF smaller than the lattice energy of KF?

Problem 11

What type of structure do metallic elements form and through what bonds?

Problem 12

What are the two types of metallic alloys and what are there differences?

Problem 13

Draw a Lewis Diagram for Acetic Acid CH_3COOH.

## Problem 14

Draw the Lewis Diagram for CO_2

Problem 15

Draw the Lewis Diagram(s) for ozone, O_3

Problem 16

Write the formal charges for all three molecules above.

Problem 17

What is the electron geometry, molecular geometry, and hybridization of the central atom in this molecule.

3 Unit 3: Intermolecular Forces and Properties

Problem 18

What are the intermolecular forces present among these molecules.

Problem 19

What phase transition is this?

Problem 20

Originally, a sample of gas is in a rigid container at $299K$ and $0.70atm$. The student increases the temperature of the $CO_2(g)$ in the container to $425K$.
(a) What does raising the temperature do to the mostion of the molecules?
(b) What is the pressure at $425K$?
(c) In terms of Kinetic Molecular Theory, why does the pressure of gas change as it is heated?

Problem 21

A $60.3g$ of $Be(OH)_2$ is dissolved in enough water to produce $1.75L$ of solution. Calculate the concentration of OH^- ions.

Problem 22

Describe the photoelectric effect.

4 Unit 4: Chemical Reactions

Problem 23

Balance this reaction: $C_5H_{10} + O_2 \longrightarrow CO_2 + H_2O$

Problem 24

Balance this redox reaction: $MnO_4^- + I^- \longrightarrow I_2 + Mn^{2+}$

Problem 25

Aqueous $FeCl_3$ reacts with KOH to produce a solid precipitate of $Fe(OH)_3$ and aqueous KCl. What is the balanced net ionic equation?

Problem 26

What is the difference between physical changes and chemical changes?

Problem 27

H_2O and Fe are reacted according to the reaction below. There was initially $36.0g$ H_2O and $67.0g$ Fe. What is the limiting reactant, how much of the excess reactant will remain, and how much iron oxide is produced?
$3\,Fe\,(s) + 4\,H_2O\,(g) \longrightarrow Fe_3O_4\,(s)\,4\,H_2\,(g)$

Problem 28

A $56kg$ sample of CO and $6.0kg$ sample of H_2 are combined into a closed vessel.
$CO\,(g) + 2\,H_2\,(g) \longrightarrow CH_3OH\,(g)$
How many moles of $CH_3OH\,(g)$ have been produced?

5 Unit 5: Kinetics

Problem 29

For this reaction:
$CH_4\,(g) + 2\,O_2\,(g) \longrightarrow CO_2\,(g) + 2\,H_2O\,(g)$
What would be rate be in terms of each reactant and product.
CH_4 rate =
O_2 rate =
CO_2 rate =
H_2O rate =

Problem 30

If the rate of dissapearance of CH_4 equals $5.0\frac{M}{s}$ for the above reaction, what is the rate of appearance of H_2O?

Problem 31

For the above reaction, what is the reaction rate if O_2 decreases from $0.1M$ to $0.04M$ in $125ms$?

Problem 32

$A(aq) + 2B(aq) \longrightarrow Products$

Experiment	$[A]_0$	$[B]_0$	Initial Rate
1	$0.10M$	$0.10M$	$1.0 \times 10^{-2}\frac{M}{s}$
2	$0.3M$	$0.10M$	$9.0 \times 10^{-2}\frac{M}{s}$
3	$0.3M$	$0.15M$	$9.0 \times 10^{-2}\frac{M}{s}$

What is the rate law?

Problem 33

N_2O_5 decomposes by a 1st order reaction with $k = 4.80 \times 10^{-4}\frac{1}{s}$. What is the concentration of N_2O_5 after 825 seconds if the intial concentration is $0.0165M$? What is the half-life for this reaction?

## Problem 34

This problem relates to problem 35 as well
The reaction $2C_4H_6(g) \longrightarrow C_8H_{12}(g)$ is a 2nd order reaction with $k = 4.0 \times 10^{-4}\frac{1}{Ms}$. If the initial concentration of C_4H_6 is $0.100M$ what is the concentration after 6 days?

Problem 35

How long does it take for the concentration to drop to $0.085M$?

Problem 36

What is the net chemical reaction and predict the experimental rate law for a chemical reaction with this chemical mechanism.

$$H_2O_2 + I^- \longrightarrow H_2O + IO^- \quad k_1 \text{ (slow)}$$
$$IO^- + H_2O_2 \longrightarrow H_2O + O_2 + I^- \quad k_2 \text{ (fast)}$$

Also identify catalysts and intermediates.

Problem 37

Predict the experimental rate law for a chemical reaction that proceeds by the following mechanism:

$$2\,NO \rightleftharpoons N_2O_2 \quad \text{(Fast equilibrium step)}$$
$$N_2O_2 + H_2 \longrightarrow H_2O + N_2O \quad \text{(slow)}$$
$$N_2O + H_2 \longrightarrow N_2 + H_2O \quad \text{(fast)}$$

6 Unit 6: Thermodynamics

Problem 38

It takes 1.8×10^{-19} calories of energy to break an $O\text{---}H$ bond in water. How much energy does it take to break all of the $O\text{---}H$ bonds in 50.0 grams of water?

Problem 39

120. grams of an unknown metal at $100.°C$ is dropped in a styrofoam cup that contains $100.0 mL$ of water that is at $20.0 \deg C$. After some times, the final temperature of the equilibiated system is measured to be $27.3°C$. What is the specific heat capacity of the metal?