

Department of Applied Mathematics and Statistics
COLORADO SCHOOL OF MINES
MATH 455: Partial Differential Equations

Assignment #1 - Fourier Series
Due Tuesday, September 8, 2026

1. (15 points) Let f be a 2π -periodic function satisfying $f(x) = x$ for $x \in (-\pi, \pi)$.
(a) Compute the Fourier series of f and show all work.

Solution — The Fourier series of a periodic function $f(x)$ over the domain of one period $[a, b]$ is given by

$$f(x) = \frac{1}{2}a_0 + \sum_{n=1}^{\infty} a_n \cos(nx) + \sum_{n=1}^{\infty} b_n \sin(nx)$$

where

$$a_0 = \frac{2}{b-a} \int_a^b f(x) dx \quad a_n = \frac{2}{b-a} \int_a^b f(x) \cos(nx) dx$$

$$b_n = \frac{2}{b-a} \int_a^b f(x) \sin(nx) dx$$

Then for our specific 2π -periodic function $f(x) = x$, we have

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} x dx = 0 \quad (\text{by oddness})$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x \cos(nx) dx = 0 \quad (\text{also by oddness})$$

$$\begin{aligned} b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin(nx) dx \\ &= \frac{2}{\pi} \int_0^{\pi} x \sin(nx) dx \quad (\text{by evenness}) \\ &= \frac{2}{\pi} \left[\frac{-x}{n} \cos(nx) \Big|_0^{\pi} + \frac{1}{n} \int_0^{\pi} \cos(nx) dx \right] \quad (\text{by IBP}) \\ &= \frac{2}{\pi} \left[\frac{-\pi}{n} \cos(n\pi) + \frac{\sin(n\pi)}{n^2} \right] \\ &= \frac{2 \sin(n\pi)}{\pi n^2} - \frac{2}{n} \cos(n\pi) \end{aligned}$$

Thus our series expansion is

$$f(x) = \sum_{n=1}^{\infty} \left[\frac{2 \sin(n\pi)}{\pi n^2} - \frac{2}{n} \cos(nx) \right] \sin(nx)$$

Note that $\sin(n\pi) = 0$ and $\cos(n\pi) = (-1)^n$ for $n = 1, 2, 3 \dots$. Hence, we have

$$f(x) = 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1} \sin(nx)}{n}$$

- (b) Using your answer from (a), graph by hand on the same axes the three functions $f(x)$, $g(x)$, and $h(x)$ on the interval $[-\pi, \pi]$ where


$$g(x) = b_1 \sin(x)$$

and

$$h(x) = b_1 \sin(x) + b_2 \sin(2x)$$

are the one-term and two-term Fourier Series approximations of f , respectively.

We have $g(x) = 2 \sin(x)$ and $h(x) = 2 \sin(x) - \sin(2x)$



2. (5 points) Let $f(x)$ be a given function for $x \in [-\pi, \pi]$ and find $c \in \mathbb{R}$ which minimizes

$$\int_{-\pi}^{\pi} (f(x) - c)^2 dx.$$

Note that your answer should depend in some manner upon the given function f .

Let us find the c that minimizes

$$D(c) = \int_{-\pi}^{\pi} (f(x) - c)^2$$

Note that $D(c)$ appears to be some of sort of distance-squared metric measuring the accumulated distance between $f(x)$ and the line $y = c$. So intuitively, we expect c has to be something like the average value of $f(x)$ in order for all of f to be as close as possible to c —that is, to minimize this distance. We will use the first derivative test.

$$\frac{dD}{dc} = \frac{d}{dc} \int_{-\pi}^{\pi} (f(x) - c)^2 dx$$

We can bring the derivative inside the integral because it is independent of the variable of integration. Doing this will translate the derivative into a partial derivative, because the integrand is a function of both x and the parameter c .

$$\begin{aligned} &= \int_{-\pi}^{\pi} \frac{\partial}{\partial c} (f(x) - c)^2 dx \\ &= \int_{-\pi}^{\pi} 2(f(x) - c)(0 - 1) dx \\ &= \int_{-\pi}^{\pi} (c - 2f(x)) dx \end{aligned}$$

3. (10 points) Let $f(x)$ be a given function for $x \in [-\pi, \pi]$ and find $a, b, c \in \mathbb{R}$ which minimize

$$\int_{-\pi}^{\pi} (f(x) - a \cos(x) - b \sin(x) - c)^2 dx.$$

Again, note that your answer should depend in some manner upon the given function f and use the analogous hint as above for a multivariate function.