

Numerical Relativity Studies of Black Holes Beyond General Relativity and Compact
Finite-Difference Methods

Andrew James Carroll

A dissertation submitted to the faculty of
Brigham Young University
in partial fulfillment of the requirements for the degree of
Doctor of Philosophy

Eric Hirschmann, Advisors
David Neilsen
Gus Hart
Chris Verharen
Ben Frandsen

Department of Physics and Astronomy
Brigham Young University

Copyright © [2026] Andrew James Carroll

All Rights Reserved

ABSTRACT

Numerical Relativity Studies of Black Holes Beyond General Relativity and Compact Finite-Difference Methods

Andrew James Carroll
Department of Physics and Astronomy, BYU
Doctor of Philosophy

This dissertation analyzes two complementary aspects of numerical relativity. First, we generalize our numerical relativity code to evolve black holes in alternative theories of gravity. In particular, we develop a numerical framework for studying black holes in Einstein–Maxwell–Dilaton–Axion theory. Black holes in this theory can support additional scalar and pseudoscalar fields, or hair, that are not present for black holes in general relativity. We present a new method for generating approximate binary black-hole initial data in EMDA theory and use head-on collisions to demonstrate the stability of charged black holes in this setting. We then develop a modest catalog of inspiraling charged black-hole waveforms and compare these waveforms with general-relativistic reference waveforms. These comparisons show that the additional fields can produce changes in the gravitational waveform, with differences that generally increase with charge. We then further explore the broader theory space within the EMDA theory.

Second, we study compact finite-difference derivative operators for use in adaptive numerical simulations. These operators are of interest because they have the potential to improve the accuracy and efficiency of numerical relativity calculations. We construct several novel compact finite-difference operators and test their performance on a variety of problems, including the Maxwell equations, the nonlinear sigma model, and the BSSN equations. These tests allow us to compare convergence, stability, and robustness across both linear and nonlinear systems. We show that compact finite-difference operators can improve performance compared to their explicit counterparts.

Keywords: Numerical relativity, alternative theories of gravity, black holes, gravitational waves, compact finite-difference methods, adaptive mesh refinement

ACKNOWLEDGMENTS

Contents

Table of Contents	iv
List of Figures	vii
List of Tables	xiv
1 Introduction	1
1.1 Introduction	2
1.2 Background	6
1.3 Summary of Dissertation	12
1.4 Conventions	13
2 EMDA Chapter 1	15
2.1 Introduction	16
2.2 Einstein–Maxwell–Dilaton–Axion	20
2.3 Numerical methods	20
2.3.1 Code Description	21
2.3.2 Initial Data	22
2.4 Results	23
2.4.1 Initially Scalarized Black Holes	23
2.4.2 Initially Unscalarized Black Holes	26
2.4.3 Constraint and refinement tests	29
2.5 Conclusion	31
2.6 Summary of Tests Done	31
3 EMDA Chapter 2	33
3.1 Introduction	34
3.2 Einstein–Maxwell–dilaton–axion Theory	36
3.3 Numerical Methods	37
3.3.1 Numerical implementation	38
3.3.2 Controlling the Eccentricity	39
3.3.3 Waveform Mismatch and Distinguishability	44

3.4	Results	45
3.5	Conclusion	48
3.6	Simulation Catalog	50
3.7	Measured Eccentricities	50
4	EMDA Chapter 3	54
4.1	Introduction	55
4.2	Theory space	57
4.3	Numerical methods	58
4.3.1	Initial data	59
4.4	Results	60
4.4.1	Waveform comparison	60
4.4.2	Constraints	61
4.5	Conclusion	62
5	CFD Chapter 1	66
5.1	CFD	67
6	Conclusion	68
6.1	Conclusion	69
Appendix A	Mathematical Overview	72
A.1	Electricity and Magnetism	73
A.2	Spacetime and Kinematics	75
A.3	General Relativity	78
A.3.1	Reformulating the Einstein Equations	82
A.4	EMDA Equations of Motion	92
Appendix B	EMDA Black Holes	97
B.1	Schwarzschild Metric	98
B.2	Reissner–Nordström Metric	99
B.3	Kerr Metric	102
B.4	Kerr–Newman Metric	104
B.5	Einstein–Maxwell–Dilaton Black Holes	107
B.6	Kerr–Sen Metric	109
Appendix C	EMDA Initial data	111
C.1	Initial Data	112
C.1.1	TPID Solver	116

Appendix D	Computation of Ψ_4	119
D.1	Wave Extraction	120
D.1.1	Null Tetrad and Spatial Triad	120
D.1.2	Gravitational Radiation	121
D.1.3	Electromagnetic, Dilaton, and Axion Radiation	123
D.1.4	Multipolar Decomposition	123
D.1.5	Radiated Energy Fluxes	124
D.1.6	Relation to the 3 + 1 Variables	125
D.1.7	Waveform Mismatch	126
Appendix E	Dendro Notes	128
E.1	Wavelet Adaptive Mesh Refinement	129
E.2	Sobolev-Informed Refinement	131
E.3	Performance of the Code	134
E.3.1	Constraint Quantities	134
E.3.2	Inspiral Constraint Quantities	139
Appendix F	CFD Notes	142
F.0.1	Stability tests	143
Appendix G	Running the EMDA Code on MaryLou	144
G.1	Building emda-gr	144
G.2	Building the Solver	147
G.2.1	Build Outputs	148
G.3	Runs and Job Scripts	149
G.3.1	Submitting a Run with Slurm	150
G.4	Parameter-File Explanation	152
Appendix H	ParaView	169
H.1	Opening Simulation Data	169
H.2	Creating a Slice	170
H.3	Adjusting the Slice and Viewing Data	171
H.4	Using the Calculator	171
H.5	Warping by Scalar	172
H.6	Using the Color Map Editor	173
H.7	Macros, Animations, CSV Output, and Finding Data	173
Appendix I	Waveform Catalog	176
I.1	Gravitational Waveform Catalog	177
I.2	Φ_1 Waveform Catalog	177
I.3	Φ_2 Waveform Catalog	177
	Bibliography	194

List of Figures

1.1	Schematic illustration of the LIGO interferometer used to detect gravitational waves through differential arm length changes.	3
1.2	Walltime of numerical relativity simulations as a function of mass ratio.	5
1.3	Spectral resolution of explicit and compact fourth-order first-derivative operators. The blue line represents perfect resolution, and the dashed line marks the cutoff frequency used in the optimization. The compact finite-difference operator maintains better agreement with the ideal resolution curve over a wider range of wavenumbers.	13
2.1	The real portions of the $(2, 0)$ modes of the gravitational radiation Ψ_4 , electromagnetic radiation Φ_1 , and Φ_2 , extracted at $R_{\text{ex}} = 100M$ for a binary with charge-to-mass ratio $q_i/m_i = 0.1$, dimensionless spin $\chi = 0.4$, and an initial coordinate separation of $16M$	24
2.2	Plot of the radiated energy for the case $\chi = 0.4$ and $q_i/m_i = 0.1$. The curves have been aligned such that $t = 0$ corresponds to the global maximum. On this plot the axion and dilaton radiation energy remain at least 3 orders of magnitude smaller than the GW or EM contributions	26

- 2.3 Scalar fields on the $z = 0$ plane for the case $\chi = 0.4$ and $q_i/m_i = 0.1$ Kerr–Sen black holes. We show the amplitudes of the axion, κ (top four panels), and dilaton, ϕ (bottom four panels), fields at the beginning of the evolution (top left), shortly before merger (top right), just after merger (bottom left), and at a later time after merger (bottom right). 27
- 2.4 Percent deviation of the remnant black hole’s horizon mass M , electric charge Q , and dimensionless spin χ from their calculated average values: $\langle M \rangle = 0.9682$, $\langle Q \rangle = 0.2001$, and $\langle \chi \rangle = 0.1961$. These quantities are measured on the apparent horizon of the remnant black hole for an initially spinning and charged binary black hole configuration with individual charge-to-mass ratios $q_i/m_i = 0.1$, initial dimensionless spins $\chi = 0.4$, and total mass $M = 1$ 28
- 2.5 Scalar fields on the $z = 0$ plane for Kerr–Newman black holes with initial dimensionless spin $\chi = 0.4$ and charge-to-mass ratio $q_i/m_i = 0.1$. Initially, neither black hole carries dilaton or axion hair. Shown are the amplitudes of the scalar fields κ and ϕ at various stages of the evolution: the initial stage (top left), where both fields begin at zero; a short time later as the fields start to develop (top right); just before merger (bottom left); and some time after merger (bottom right). 30
- 3.1 Orbital dynamics for the $q = 1$ and $q = 2$ configurations. For each mass ratio, the left panel shows the coordinate separation as a function of time, while the right panel shows representative coordinate trajectories in the orbital plane. The charge configuration affects both the inspiral time and the orbital phasing through the modified effective attraction. 41
- 3.2 Orbital dynamics for the $q = 4$ and $q = 8$ configurations. For each mass ratio, the left panel shows the coordinate separation as a function of time, while the right panel shows representative coordinate trajectories in the orbital plane. 42

3.3	Measured orbital-frequency eccentricity and fitted eccentricity model for a binary black-hole inspiral with $q = 1$, $\chi = 0.4$, $\lambda = 0.3$, and charge configuration $++$. The fit is performed from $t_{\min} = 200M$ to $t_{\max} = 1300M$, excluding early gauge-settling effects and ending before the onset of plunge. The measured eccentricity extrapolated to $t = 0M$ is $\varepsilon_{\Omega} = (14.44 \pm 0.33) \times 10^{-3}$	44
3.4	Peak-aligned comparisons of the gravitational-wave strain component rh_+ for representative charged EMDA configurations against the corresponding neutral GR reference waveforms. In each panel, the time coordinate is shifted so that the waveform peak occurs at $t - t_{\text{peak}} = 0$. The remaining differences after peak alignment reflect changes in the accumulated phase, merger time, and waveform morphology.	47
4.1	Peak-aligned comparison of the dominant gravitational-wave strain between the Einstein–Maxwell reference waveform, $(\alpha_0, \alpha_1) = (0, 0)$, and the strongest-coupling EMDA case, $(\alpha_0, \alpha_1) = (100, 100)$. The figure shows the plus polarization of the strain.	62
4.2	Hamiltonian constraint violation for the different coupling choices. The curves show the evolution of the Hamiltonian constraint norm for the Einstein–Maxwell reference case and the corresponding EMD and EMDA evolutions.	63
4.3	Electric divergence constraint violation for the different coupling choices. The curves show the evolution of the electric constraint norm across the Einstein–Maxwell, EMD, and EMDA evolutions.	64
4.4	Magnetic divergence constraint violation for the different coupling choices. The curves show the evolution of the magnetic constraint norm across the Einstein–Maxwell, EMD, and EMDA evolutions.	65

- C.1 Example of the compactified TwoPunctures grid in a slice through the two punctures. The physical coordinates are shown in units of the half-separation b , with the black-hole punctures located at $x = \pm b$. The coordinate lines illustrate how the pseudo-spectral collocation points are distributed in the compactified domain before the resulting initial data are interpolated onto the evolution grid. The outer compactified boundary corresponds to spatial infinity and is outside the finite plotting window. 118
- E.1 Schematic illustration of the nested dyadic grids used in wavelet adaptive mesh refinement. Points present on coarser grids are retained on finer grids, while the red points indicate the additional degrees of freedom introduced at each refinement level. These additional points correspond to the detail spaces W_j 129
- E.2 Representative comparison of the adaptive mesh generated by the baseline wavelet AMR criterion and the Sobolev-informed AMR criterion. The Sobolev-informed criterion includes derivative information in the refinement indicator, which changes where grid points are allocated. The oscillatory ring corresponds to the outward-propagating dipole pulse. 132
- E.3 Constraints for the baseline wavelet AMR and Sobolev-informed AMR refinement criteria. The left panel shows the L_2 norm of the electric-field divergence constraint, while the right panel shows the corresponding divergence for the vector potential. These quantities are used to assess whether the modified refinement indicator improves or degrades the constraint behavior of the evolution. 133

- E.4 Mesh size and grid-efficiency for the baseline wavelet AMR and Sobolev-informed AMR refinement criteria. The left panel shows the number of active mesh points N_{mesh} , which serves as a proxy for computational cost. The right panel shows $\eta = 1/(C_{\text{tot}}N_{\text{mesh}}^2)$. Larger values of η indicate smaller constraint violation for a given mesh size, and therefore a more efficient use of grid points. 134
- E.5 The Hamiltonian constraint $\mathcal{H}$ and the summed momentum constraint violation measured over the full simulation. The constraints remain controlled throughout the evolution, including the merger phase. 135
- E.6 The volume-weighted L_2 norms of the electromagnetic constraints measured over the full simulation. These include the magnetic no-monopole constraint, $\nabla \cdot B$, and the electric Gauss constraint, $\nabla \cdot E$ 136
- E.7 The sum of the volume-weighted Hamiltonian and momentum constraint violations over the first $20M$ of a head-on evolution of two equal-mass charged black holes with charge-to-mass ratio $q/m = 0.1$ and dimensionless spin $\chi = 0.4$. The simulations use wavelet tolerances $\varepsilon \in \{3 \times 10^{-5}, 1 \times 10^{-5}, 3 \times 10^{-6}\}$ and a maximum refinement level of $j_{\text{max}} = 15$. Lowering the wavelet tolerance improves the resolution of dynamically important regions and reduces the overall constraint violation. 138
- E.8 The sum of the volume-weighted Hamiltonian and momentum constraint violations over the first $20M$ of a head-on evolution of two equal-mass charged black holes with charge-to-mass ratio $q/m = 0.1$ and dimensionless spin $\chi = 0.4$. The simulations compare maximum refinement levels $j_{\text{max}} \in \{14, 15, 16, 17\}$, corresponding to finest grid spacings $h \in \{0.0325, 0.0162, 0.00813, 0.00407\}$, respectively. The wavelet tolerance is fixed at $\varepsilon = 1 \times 10^{-5}$. Increasing the maximum refinement level reduces the overall constraint violation by allowing additional grid refinement in under-resolved regions. 139

E.9	The Hamiltonian constraint $\mathcal{H}$ and the summed momentum constraint violation measured over the full inspiral simulation. The L_2 norm is normalized by the volume of the grid. This representative case has $q = 1$, $\chi = 0.4$, and $\lambda_+^+ = 0.3$	140
E.10	The volume-weighted L_2 norms of the electromagnetic constraints measured over the full inspiral simulation. These include the magnetic no-monopole constraint, $\nabla \cdot B$, and the electric Gauss constraint, $\nabla \cdot E$. This representative case has $q = 1$, $\chi = 0.4$, and $\lambda_+^+ = 0.3$	141
H.1	The ParaView interface with the Open File icon highlighted.	170
H.2	The ParaView interface with the Slice and Calculator tools highlighted.	170
H.3	ParaView controls for stepping through time, selecting the displayed field or color option, and switching between representations such as Surface and Wireframe . .	172
H.4	The Warp By Scalar settings in the Properties panel. The Scalars menu selects the field used for the warping, and the Scale Factor controls the strength of the warp. .	173
H.5	The Color Map Editor panel in ParaView. This panel controls the color scale, opacity, and transfer function used to visualize the selected field.	174
I.1	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.	178
I.2	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.	179
I.3	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0.8$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.	180

I.4	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 2$ and $\chi = 0$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.	181
I.5	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 2$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.	182
I.6	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 4$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations with $ \lambda_i = 0.05$ and $ \lambda_i = 0.10$	183
I.7	Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 8$ and $\chi = 0.2$	183
I.8	Extracted Φ_1 waveforms for $q = 1$ and $\chi = 0$	184
I.9	Extracted Φ_1 waveforms for $q = 1$ and $\chi = 0.4$	185
I.10	Extracted Φ_1 waveforms for $q = 1$ and $\chi = 0.8$	186
I.11	Extracted Φ_1 waveforms for $q = 2$ and $\chi = 0$	187
I.12	Extracted Φ_1 waveforms for $q = 2$ and $\chi = 0.4$	187
I.13	Extracted Φ_1 waveforms for $q = 4$ and $\chi = 0.4$	188
I.14	Extracted Φ_1 waveforms for $q = 8$ and $\chi = 0.2$	188
I.15	Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0$	189
I.16	Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0.4$	190
I.17	Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0.8$	191
I.18	Extracted Φ_2 waveforms for $q = 2$ and $\chi = 0$	192
I.19	Extracted Φ_2 waveforms for $q = 2$ and $\chi = 0.4$	192
I.20	Extracted Φ_2 waveforms for $q = 4$ and $\chi = 0.4$	193
I.21	Extracted Φ_2 waveforms for $q = 8$ and $\chi = 0.2$	193

List of Tables

- 3.1 EMDA binary black-hole simulation catalog. Here $q = m_1/m_2$ denotes the mass ratio, χ denotes the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$ denotes the charge-to-mass ratio of black hole i . The neutral case corresponds to $Q_1 = Q_2 = 0$. For charged configurations, the magnitude of the nonzero charge-to-mass ratios is denoted by λ , with the sign structure indicated by the charge configuration. The configurations ++, +−, and +0 denote like-signed charges, oppositely signed charges, and cases in which only one black hole is charged, respectively. Because the neutral configuration is independent of λ , it is listed only once for each mass ratio and spin. 51

- 3.2 Mismatch of charged EMDA binary black-hole waveforms relative to the corresponding neutral reference waveform Q0. Here $q = m_1/m_2$, χ is the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$. The column $|\lambda_i|$ gives the magnitude of the nonzero charge-to-mass ratio used in each charged configuration. The columns ++, +−, and +0 denote like-signed charges, oppositely signed charges, and configurations in which only one black hole is charged, respectively. The mismatch is computed from the $(\ell, m) = (2, 2)$ gravitational waveform using the numerical-relativity analysis package `kuibit`, after optimizing over relative time and polarization shifts. The values in parentheses give the approximate fixed-parameter critical signal-to-noise ratio, $\rho_{\text{crit}} \simeq 1/\sqrt{2\mathcal{M}}$. These values should be interpreted as rough waveform-separation estimates rather than full observational detection thresholds. 52
- 3.3 Measured orbital eccentricities for the EMDA waveform catalog. The entries are reported as $(\varepsilon_\Omega \pm \delta\varepsilon_\Omega) \times 10^3$. Here $q = m_1/m_2$ denotes the mass ratio, χ denotes the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$ denotes the charge-to-mass ratio of black hole i . The neutral case corresponds to $Q_1 = Q_2 = 0$. For charged configurations, the magnitude of the nonzero charge-to-mass ratios is denoted by λ , with the sign structure indicated by the charge configuration. The configurations ++, +−, and +0 denote like-signed charges, oppositely signed charges, and cases in which only one black hole is charged, respectively. 53

Chapter 1

Introduction

1.1 Introduction

Since Einstein introduced the theory of general relativity in 1915 [1], it has remained consistent with all experimental tests to date [2, 3]. General relativity has fundamentally transformed our understanding of space and time, explaining phenomena ranging from the anomalous precession of Mercury to the operation of atomic clocks in GPS satellites [4], and predicting the existence of compact objects such as neutron stars and black holes, as well as gravitational waves [5–7].

Gravitational waves were first directly detected by LIGO in 2015 [8]. These signals originate from some of the most energetic events in the universe, such as mergers of neutron stars and black holes. However, the signals detected by LIGO are extremely weak; the detectors are sensitive enough to measure relative length changes on the order of 10^{-21} , corresponding to displacements smaller than a proton diameter [9]. Despite this extraordinary sensitivity, gravitational wave signals remain weak relative to background noise. The highest signal-to-noise ratio (SNR) gravitational-wave event reported to date is GW250114, with a network matched-filter SNR of approximately 80 [10].

Interpreting gravitational-wave detections requires accurate theoretical models of the expected signals. In practice, this is achieved by comparing observational data against waveform templates constructed from analytical approximations, phenomenological models, and numerical simulations. This template-based comparison allows the physical properties of the source, including the component masses and spins, to be inferred statistically [11, 12].

Constructing theoretical models of binary black-hole mergers is challenging because the relevant dynamics are strongly nonlinear and only partially accessible through analytic methods. In general relativity, exact analytic solutions are typically restricted to highly symmetric systems and cannot describe the full inspiral, merger, and ringdown of a generic binary black-hole system. Numerical relativity therefore plays an essential role in modeling the late inspiral, merger, and ringdown, where fully nonlinear effects are important. A number of numerical relativity codes have been developed




Figure 1.1 Schematic illustration of the LIGO interferometer used to detect gravitational waves through differential arm length changes.

for this purpose, including the Einstein Toolkit, SpEC, GRChombo, BAM, GR-Athena++, and Dendro-GR [13–18].

Simulating binary black-hole mergers is computationally demanding. These simulations require solving a coupled system of nonlinear partial differential equations across multiple length scales, including the scale of the individual black holes, the orbital scale of the binary, and the radiation zone where gravitational waves are extracted [19, 20]. In this work, we make use of the Dendro numerical relativity framework, which combines adaptive mesh refinement with scalable parallel algorithms for binary black-hole simulations [18, 21, 22].

For standard general relativity (GR), substantial progress has been made, and extensive waveform catalogs now cover much of the binary-black-hole parameter space, particularly for near-equal-mass systems. Here we define the mass ratio as $q = M_{\text{larger}}/M_{\text{smaller}} \geq 1$. The LIGO–Virgo–KAGRA (LVK) collaboration has reported hundreds of compact-binary candidates using waveform models informed by analytical approximations and numerical relativity [14, 23–25]. However, two key challenges remain: improving the fidelity of GR simulations and developing waveform catalogs for alternative theories of gravity.

The next generation of gravitational-wave detectors is expected to significantly increase the signal-to-noise ratios of observed compact-binary signals. Meeting this challenge will require higher-fidelity numerical simulations and more accurate waveform models [26, 27].

In general, higher-fidelity simulations, particularly those with larger mass ratios, require finer grid resolution and therefore substantially greater computational cost. This trend is illustrated in Fig. 1.2. Current simulations with mass ratios $q < 8$ can require weeks of runtime on modern supercomputers, and further increases in resolution would significantly amplify this cost [14, 27].

To obtain more accurate waveforms, there are two main approaches: running existing codes at higher resolution or improving the underlying numerical methods. A drawback of the brute-force approach—simply increasing resolution—is that the timescale of the problem is constrained by the CFL factor. This, in turn, requires more timesteps, significantly increasing the total runtime of the simulation.

One potential avenue for improving efficiency in numerical relativity codes lies in the computation of spatial derivatives. Most current implementations rely on explicit finite difference methods, while spectral methods—although highly accurate—are often difficult to implement. In this work, we explore compact finite difference (CFD) operators as an alternative. Compact schemes are implicit methods that can achieve higher spectral resolution than comparable explicit schemes [28]. In tests performed with our Maxwell solver, the compact finite-difference schemes generally perform




Figure 1.2 Walltime of numerical relativity simulations as a function of mass ratio.

better than the corresponding explicit methods. In some cases, the compact schemes achieve comparable accuracy to explicit schemes at approximately half the resolution. Despite their success in computational fluid dynamics and related fields, their use in numerical relativity remains limited [29].

In addition to the need for improved waveform modeling in GR, there is also a need for waveform models in alternative theories of gravity. Gravitational-wave observations provide a powerful probe of gravity in the strongly nonlinear, dynamical regime, where deviations from general relativity (GR) may become apparent [3, 30, 31]. To search for such deviations, waveform templates from alternative theories of gravity are required [32–34]. While studies have been carried out in theories such as Chern–Simons gravity [35, 36], scalar–tensor theories [37–40], Einstein–Gauss–Bonnet gravity [41–46], and Einstein–Maxwell–dilaton theory [47, 48], waveform catalogs in these frameworks remain limited [49–51].

In this dissertation, we focus on Einstein–Maxwell–dilaton–axion (EMDA) theory. EMDA is of particular interest because it admits a well-posed initial value formulation and arises as the

low-energy limit of string theory. In contrast to the no-hair theorem in standard GR, which states the black holes can be completely defined by their mass spin and charge, EMDA black holes possess additional degrees of freedom associated with the dilaton and axion fields.

We present a numerical relativity framework for simulating binary black hole mergers in EMDA theory. We analyze the resulting waveforms and investigate whether deviations from GR may be observable in gravitational wave signals.

1.2 Background

We focus on the study of black hole dynamics. As such, we begin with Einstein's field equations:

$$G_{ab} = 8\pi T_{ab} + \Lambda g_{ab}. \quad (1.1)$$

The left-hand side describes the geometry of spacetime, while the right-hand side encodes the matter and energy content. [52, 53] We briefly examine each term in turn.

The tensor G_{ab} is the Einstein tensor, which is constructed from quantities that encode the curvature of spacetime. The stress-energy tensor T_{ab} describes the distribution of matter and energy. The metric tensor g_{ab} defines distances and causal structure in spacetime. The constant Λ is the cosmological constant, which is generally assumed to be very small. On the scales considered in this work, it is negligible, and we therefore set $\Lambda = 0$.

In standard general relativity simulations of binary black hole mergers, we consider vacuum spacetimes, for which

$$G_{ab} = 0. \quad (1.2)$$

This provides the system of partial differential equations that must be solved in vacuum GR. Before proceeding further, we review some essential geometric concepts.

We begin with the metric, which encodes the notion of separation in spacetime. As an example, the Minkowski metric describing flat spacetime can be written as

$$ds^2 = -dt^2 + dx^2 + dy^2 + dz^2, \quad (1.3)$$

or equivalently in matrix form as

$$\eta_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (1.4)$$

Here we use η_{ab} to denote the flat spacetime metric. The negative sign in the time component reflects our choice of signature $(-, +, +, +)$.

The Minkowski metric can also be expressed in different coordinate systems. For example, in spherical coordinates it takes the form

$$\eta_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}. \quad (1.5)$$

Although these representations appear different, they describe the same flat spacetime. This illustrates that the form of the metric depends on the choice of coordinates.

A natural question is how to determine whether a given metric describes flat spacetime. A necessary and sufficient condition is that the Riemann curvature tensor vanishes everywhere. Given a metric, the Riemann tensor is defined as

$$R^a{}_{bcd} = \partial_c \Gamma^a{}_{bd} - \partial_d \Gamma^a{}_{bc} + \Gamma^a{}_{ec} \Gamma^e{}_{bd} - \Gamma^a{}_{ed} \Gamma^e{}_{bc}, \quad (1.6)$$

where $\Gamma^a{}_{bc}$ are the Christoffel symbols, given by

$$\Gamma^a{}_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{dc} + \partial_c g_{bd} - \partial_d g_{bc}). \quad (1.7)$$

These quantities encode how the coordinate system varies from point to point.

Contracting the Riemann tensor yields the Ricci tensor,

$$R_{ab} = g^{cd} R_{acbd}, \quad (1.8)$$

and further contraction gives the Ricci scalar,

$$R = g^{ab} R_{ab}. \quad (1.9)$$

The Einstein tensor is then constructed as

$$G_{ab} = R_{ab} - \frac{1}{2} g_{ab} R, \quad (1.10)$$

which satisfies the contracted Bianchi identities and ensures local conservation of energy and momentum.

With the Einstein tensor in hand, we can rewrite the field equations as a set of coupled second-order partial differential equations. To evolve these equations numerically, we must decompose them into a form suitable for time evolution. This requires splitting spacetime into three spatial dimensions and one time dimension.

The most straightforward approach leads to the ADM (Arnowitt–Deser–Misner) formulation. However, the ADM system is only weakly hyperbolic and can become unstable over long evolutions. Several modified formulations have been developed to address this issue. In this work, we use the BSSN (Baumgarte–Shapiro–Shibata–Nakamura) formulation, which is strongly hyperbolic and can be evolved stably for long periods of time. Given this formulation, we require suitable initial data, after which the system can be evolved forward in time.

To provide an overview of this procedure, we briefly consider simpler examples: a scalar field and Maxwell’s equations.

We begin with the scalar wave equation with a source term:

$$\square\psi = \nabla_a \nabla^a \psi = 4\pi\rho. \quad (1.11)$$

This equation is written in fully covariant form. Expanding it in flat spacetime using the Minkowski metric, we obtain

$$\square\psi = \eta^{ab}\nabla_a\nabla_b\psi = \left(-\partial_t^2 + D^2\right)\psi, \quad (1.12)$$

where D denotes purely spatial derivatives.

This provides a simple example of a 3+1 decomposition. The scalar wave equation is a second-order hyperbolic equation with well-posed initial value properties: solutions exist and are unique. Importantly, there are no constraint equations in this system, we are free to choose initial data for ψ and $\partial_t\psi$. As we will see, this is not the case for Maxwell's equations or general relativity, where constraint equations must be satisfied.

We now turn to the more complex case of Maxwell's equations:

$$\nabla \cdot \mathbf{E} = 4\pi\rho, \quad (1.13)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (1.14)$$

$$\frac{\partial \mathbf{E}}{\partial t} = \nabla \times \mathbf{B} - 4\pi\mathbf{J}, \quad (1.15)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}. \quad (1.16)$$

We present the full 3+1 decomposition of Maxwell's equations in the appendix. However, an important feature is already evident: there are eight equations, but only six dynamical variables ($E_x, E_y, E_z, B_x, B_y, B_z$). The remaining two equations act as constraints on the system.

Maxwell's equations can be separated into evolution equations:

$$\frac{\partial \mathbf{E}}{\partial t} = \nabla \times \mathbf{B} - 4\pi\mathbf{J}, \quad (1.17)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}, \quad (1.18)$$

and constraint equations:

$$\nabla \cdot \mathbf{E} = 4\pi\rho, \quad (1.19)$$

$$\nabla \cdot \mathbf{B} = 0. \quad (1.20)$$

Specifically, Gauss’s law and the absence of magnetic monopoles must be satisfied at all times. If the initial data satisfy these constraints, then in the continuum limit the evolution equations will preserve them. However, in numerical simulations we work with finite precision, and violations of the constraint equations provide a useful measure of numerical error.

Solving the Cauchy problem for Einstein’s equations is nontrivial, as we must satisfy both the Hamiltonian and momentum constraints. There are established techniques for solving these constraints. In particular, the momentum constraints are often addressed using the Bowen–York formalism. The Hamiltonian constraint reduces to an elliptic equation that must be solved numerically.

In EMDA, solving the Cauchy problem is further complicated by the presence of additional source terms. The stress-energy tensor T_{ab} encodes all contributions from matter and energy apart from gravity. In standard GR this is 0, in EMDA, this includes electromagnetic fields as well as scalar fields. The solution for single black holes is known analytically for the certain coupling strengths ($\alpha_0 = \alpha_1 = 1$) but numerically one could also imagine varying the coupling strength to see how the evolutions change. This dissertation will address the challenges associated with constructing consistent initial data and evolving these systems.

Once suitable initial data have been constructed and the system has been evolved, we extract the gravitational waveform. In this work, we compute the waveform using the Newman–Penrose formalism, in which outgoing gravitational radiation is represented by the Weyl scalar Ψ_4 [54]. Details of how Ψ_4 is calculated are provided in Appendix ??.

The scalar Ψ_4 is related to the gravitational-wave strain $h = h_+ - ih_\times$ by

$$\Psi_4 = \frac{d^2 h}{dt^2}, \quad (1.21)$$

up to the sign conventions used for the null tetrad. From Ψ_4 , we reconstruct the strain and compare simulated waveforms with one another and, in principle, with observed gravitational-wave signals. To quantify the difference between two waveforms, we compute the mismatch,

$$\text{mismatch}(h_1, h_2) = 1 - \max \mathcal{O}(h_1, h_2), \quad (1.22)$$

where O is the normalized waveform overlap and the maximum is taken over time shifts and polarization angles.

The mismatch can be used to estimate the signal-to-noise ratio required to distinguish two waveforms. In the small-mismatch limit, this distinguishability threshold is approximately

$$\rho \sim \frac{1}{\sqrt{2, \text{mismatch}(h_1, h_2)}}. \quad (1.23)$$

This estimate provides a useful way to interpret waveform differences, although it should not be regarded as an absolute detection criterion. In practice, interpreting differences between waveforms remains challenging because the parameter space is often degenerate. For example, it may not be immediately clear whether observed differences arise from spin, charge, or other physical effects.

Improving the accuracy of these waveform comparisons requires improving the numerical simulations that produce them. We therefore also present our work incorporating compact finite differences into our numerical relativity code. To begin, we first review the finite-difference methods currently used in the code.

On a discrete grid, derivatives can be approximated using finite differences. For example, a second-order accurate centered approximation to the first derivative is

$$\left. \frac{df}{dx} \right|_i = \frac{f_{i+1} - f_{i-1}}{2\Delta x} + O(\Delta x^2). \quad (1.24)$$

Higher-order accurate formulas can be constructed by matching Taylor series expansions. For instance, a fourth-order accurate centered stencil is

$$\left. \frac{df}{dx} \right|_i = \frac{-f_{i+2} + 8f_{i+1} - 8f_{i-1} + f_{i-2}}{12\Delta x} + O(\Delta x^4). \quad (1.25)$$

These finite-difference approximations are standard tools in numerical solutions of partial differential equations [19].

These are examples of explicit finite-difference schemes. Alternatively, one can approximate the first derivative using a compact, or implicit, finite-difference scheme of the form

$$\beta f'_{i-2} + \alpha f'_{i-1} + f'_i + \alpha f'_{i+1} + \beta f'_{i+2} = c \frac{f_{i+3} - f_{i-3}}{6\Delta x} + b \frac{f_{i+2} - f_{i-2}}{4\Delta x} + a \frac{f_{i+1} - f_{i-1}}{2\Delta x}. \quad (1.26)$$

Here f'_i denotes the numerical approximation to df/dx at grid point i . Compact finite-difference schemes of this type were developed to obtain higher spectral resolution than comparable explicit schemes [28].

The coefficients are determined by matching Taylor series expansions to the desired order of accuracy. For the above stencil, the consistency conditions include

$$a + b + c = 1 + 2\alpha + 2\beta \quad (2\text{nd order}), \quad (1.27)$$

$$a + 2^2b + 3^2c = 2\frac{3!}{2!}(\alpha + 2^2\beta) \quad (4\text{th order}), \quad (1.28)$$

$$a + 2^4b + 3^4c = 2\frac{5!}{4!}(\alpha + 2^4\beta) \quad (6\text{th order}). \quad (1.29)$$

This approach is more complex than an explicit finite-difference stencil because the derivatives are obtained by solving an implicit system. The trade-off is that compact schemes can provide improved accuracy and enhanced spectral resolution compared to explicit methods [28]. This improvement in spectral resolution is illustrated in Fig. 1.3. There are many possible compact finite-difference operators that can be constructed. In this work, we describe our methodology for identifying accurate and stable compact derivative operators and evaluate their performance relative to their explicit counterparts.

1.3 Summary of Dissertation

This dissertation primarily presents results from binary black-hole simulations in Einstein–Maxwell–dilaton–axion (EMDA) theory, along with a study of compact finite-difference methods for numerical relativity. The dissertation is organized as follows. Chapter 2 discusses the formulation of the initial data and presents results from head-on collision simulations. Chapter 3 presents inspiral and merger results in EMDA theory. Chapter 4 explores interpolated EMDA models in which the coupling strengths are varied. Chapter 5 investigates the implementation of compact finite-difference operators in numerical relativity codes.



Figure 1.3 Spectral resolution of explicit and compact fourth-order first-derivative operators. The blue line represents perfect resolution, and the dashed line marks the cutoff frequency used in the optimization. The compact finite-difference operator maintains better agreement with the ideal resolution curve over a wider range of wavenumbers.

Several of the results presented in this dissertation are currently in preparation for publication.

Each of these projects involved multiple collaborators. The EMDA work builds on earlier contributions by Hyun Lim at Los Alamos National Laboratory and was carried out collaboratively with Eric Hirschmann at Brigham Young University, David Van Komen at the University of Utah, Sebastian Vander Ploeg Fallon at Cornell University, and David Neilsen at Brigham Young University. The compact finite-difference (CFD) project was conducted in collaboration with Eric Hirschmann, David Van Komen, David Neilsen, Nathaniel Garey, and Luke Papenfuss.

1.4 Conventions

In this work, we use geometrized units with $c = G = 1$. Unless otherwise stated, lengths and masses are expressed in units of a characteristic mass scale M . In physical units, a length scale $d = M$ corresponds to $d = GM/c^2$, while a time scale $t = M$ corresponds to $t = GM/c^3$.

We use the metric signature $(-, +, +, +)$ and assume the Einstein summation convention unless otherwise stated. Spacetime indices $a, b, c, \dots$ run over $0, 1, 2, 3$, while spatial indices $i, j, k, \dots$ run over $1, 2, 3$. The symmetric and antisymmetric parts of a tensor are defined as

$$T_{(ab)} \equiv \frac{1}{2} (T_{ab} + T_{ba}), \quad T_{[ab]} \equiv \frac{1}{2} (T_{ab} - T_{ba}). \quad (1.30)$$

Partial derivatives are denoted by $\partial_i g$ or equivalently $g_{,i}$. Covariant derivatives are written as $\nabla_a f_b$ or equivalently $f_{b;a}$. The Lie derivative of a tensor X^a along a vector field β^a is denoted by $\mathcal{L}_\beta X^a$.

Chapter 2

EMDA Chapter 1

2.1 Introduction

While general relativity (GR) has passed all current experimental tests, there exist a large number of potentially viable extensions to it. Ongoing improvements in detector sensitivity, combined with numerical relativity will continue to enable increasingly precise tests of GR and proposed alternatives [3, 10, 31, 34, 55–61]. Indeed, advances in gravitational wave (GW) astronomy have led to a dramatic increase in the total number of detections of signals from compact binary mergers, thereby providing access to strongly gravitating, nonlinear domains. Since the first detection of gravitational waves, the Advanced LIGO and Virgo detectors have completed multiple observing runs, producing a growing catalog of compact binary mergers [23–25, 62]. All observed mergers to date are consistent with the predictions of GR within current uncertainties [8, 23, 30]. Nevertheless, compact-binary coalescences continue to provide one of the most promising arenas for testing gravity beyond GR.

The increasing precision of gravitational wave observations has motivated effort to model compact object dynamics in theories beyond general relativity [49, 50, 63–65]. Numerical relativity is essential for this program because the late inspiral, merger, and ringdown probe regimes in which the dynamics are intrinsically nonlinear and analytic approximations are not uniformly reliable. Extending numerical relativity to alternative theories of gravity, however, requires careful control of the associated initial-value problem. Reliable nonlinear simulations require evolution systems with suitable hyperbolic structure, controlled constraint propagation, and stable gauge dynamics. These considerations have become central in beyond-GR numerical relativity, where the construction of robust evolution formulations is often a prerequisite for extracting physical predictions.

The challenge is particularly relevant for theories motivated by effective field theory or higher curvature corrections to GR. Such theories introduce higher derivatives, additional characteristic modes, or principal parts whose hyperbolicity depends on the background solution and coupling strength. In some cases, higher derivatives may also be associated with Ostrogradsky-type

instabilities, unless the theory is interpreted within a controlled effective description or reformulated appropriately. A significant number of previous studies have therefore been devoted to approaches that permit a more complete consideration of nonlinear aspects while maintaining control over the fundamental evolutionary problem. Examples of these include order reduction schemes in dynamical Chern-Simons gravity [35, 36] and Einstein-dilaton-Gauss-Bonnet theory [44, 66], modified gauge choices in Einstein-scalar-Gauss-Bonnet [42, 67], the so-called “fixing the equations” approach [68], and auxiliary field formulations for solving higher curvature systems including quadratic gravity [51, 69–72] (see also [73] for a more in-depth discussion and comparison of these and other approaches). These developments have expanded the range of beyond-GR theories accessible to numerical relativity while emphasizing that dynamical consistency and observational phenomenology must be addressed together.

From this perspective, theories whose field equations can be written as second order evolution systems provide particularly useful laboratories for beyond-GR numerical relativity. They allow one to study additional propagating degrees of freedom in fully nonlinear black hole (BH) spacetimes without relying entirely on perturbative or order reduced treatments of the gravitational field equations. Einstein–Maxwell–Dilaton–Axion (EMDA) [74, 75], arising as a low-energy limit of heterotic string theory, is one such theory well suited to this purpose. Unlike many higher curvature modifications to GR, EMDA does not introduce higher derivatives into the metric field equations. Instead, it extends the Einstein–Maxwell system by coupling the electromagnetic field to two additional propagating matter degrees of freedom: a dilaton and an axion. The resulting equations retain a structure closely related to the Einstein–Maxwell system coupled to nonlinear matter fields. Consequently, EMDA can be formulated as a second order hyperbolic evolution problem using standard numerical relativity methods, together with appropriate evolution equations for the electromagnetic, dilaton, and axion fields. In the limit of vanishing axion coupling, EMDA reduces to Einstein–Maxwell–Dilaton (EMD) theory.

Having a well posed initial value problem for EMDA is a crucial ingredient for sensible simulations, but it is also advantageous as it has analytically known black hole solutions at specific points in the theory space. Certain of these are the stationary Kerr–Sen black holes. They generalize the Kerr–Newman family of general relativity in that they carry nontrivial dilaton and axion fields in addition to mass, angular momentum, and electromagnetic charge. As is well known, in standard GR coupled to Maxwell fields, stationary black holes are characterized by their mass, angular momentum, and electric charge [76, 77]. In contrast, extensions of GR such as EMDA can admit black holes with additional matter fields. Kerr–Sen solutions therefore provide a natural setting in which to explore whether these additional degrees of freedom behave as stable black hole hair in the nonlinear strong field regime. Related questions have been studied in scalar-Gauss–Bonnet gravity, where scalarization and dynamical descalarization can occur during compact binary evolutions [45, 78]. The nonlinear behavior of scalar and axionic hair in EMDA is connected to several lines of previous work. Studies of EMD [47, 48] provide an important foundation for extending nonlinear simulations to the full EMDA system. Analytical investigations have examined aspects of the perturbative stability of EMDA black holes in certain parameter regimes [79]. In addition, astrophysical studies have explored possible observational signatures of Kerr–Sen BHs. For example, analyses of Event Horizon Telescope (EHT) shadow data suggest that Sgr A* may favor the Kerr–Sen solution, although similar data from M87* favor Kerr while remaining consistent with Kerr–Sen within current observational uncertainties [80, 81].

Fully nonlinear simulations of binary BHs in EMDA remain limited. In particular, it is not yet well understood whether dilaton and axion hair persist through merger, disperse via radiation, or settle into a stable remnant configuration. Addressing these questions is important for assessing the nonlinear stability of Kerr–Sen BHs and for determining whether EMDA can provide distinct strong field signatures for future GW observations. Although significant progress has been made in waveform modeling for selected beyond-GR theories, including scalar-Gauss-Bonnet and Einstein-

dilaton-Gauss-Bonnet gravity, waveform coverage in alternative theories remains far less developed than in GR. Consequently, many current tests still rely on phenomenological parameterizations, post-Newtonian approximations, or targeted numerical relativity simulations [45, 46, 82, 83].

In this study, we extend these investigations by simulating head-on black hole mergers governed by the full EMDA equations. We analyze the resulting waveforms and examine the stability of the remnants, thereby gaining insight into the persistence of scalar hair within the theory. Specifically, we evolve several head-on collisions of BHs in EMDA to determine the conditions under which the dilaton and axion fields survive merger.

The remainder of this paper is organized as follows. In Sec. 3.2, we review the EMDA action, equations of motion, and the formulation used for numerical evolution. In Sec. 3.3, we describe our numerical methods, including the modified Dendro-GR infrastructure, the construction of approximate binary BH initial data, and the extraction of gravitational, electromagnetic, dilaton, and axion radiation. In Sec. 3.4, we present simulations of head-on Kerr-Sen BH collisions, analyze the emitted radiation, and examine the persistence of scalar hair in the remnant. We also study initially unscalarized configurations to determine whether scalar hair can be generated dynamically. We conclude in Sec. 3.5 with a summary of our results and directions for future work. Several important details such as the equations of motion, the analytic form of the Kerr-Sen solution, initial data and wave extraction techniques and issues associated with code performance are relegated to a number of appendices in order not to disrupt the continuity of the discussion. Throughout this paper we use a mostly positive signature, Latin letters as spacetime indices, and work in geometrized units, $G = c = 1$.

2.2 Einstein–Maxwell–Dilaton–Axion

The general theory we consider in this work is an extension of Einstein–Maxwell on inclusion of two coupled scalar fields, the dilaton and the axion. The action for the EMDA system is

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{8\pi} - 2(\nabla\phi)^2 - e^{-2\alpha_0\phi} F^2 - \frac{1}{2} e^{4\alpha_1\phi} (\nabla\kappa)^2 - \kappa F_{ab} (*F)^{ab} \right], \quad (2.1)$$

where the first term is the standard Hilbert action and ϕ and κ are the dilaton and axion, respectively. F_{ab} is the U(1) field strength, and $(*F)_{ab}$ is its dual. The parameters α_0 and α_1 are real, positive constants that set the coupling strengths and allow us to consider a family of EMDA theories. Particular choices correspond to particular theories. For instance, $\alpha_0 = \alpha_1 = 0$ gives Einstein–Maxwell theory minimally coupled to scalar and axion fields, while $(\alpha_0, \alpha_1) = (\sqrt{3}, 0)$ corresponds to Kaluza–Klein. In this work we take $(\alpha_0, \alpha_1) = (1, 1)$, corresponding to a low-energy limit of heterotic string theory. For these coupling values, the theory admits the Kerr–Sen family of rotating, charged black hole solutions. These solutions reduce to Kerr when the electromagnetic charge vanishes. In contrast to the Kerr–Newman solution of Einstein–Maxwell theory, the Kerr–Sen geometry is accompanied by nontrivial dilaton and axion fields whose structure is determined by the black hole charge and spin. The explicit Kerr–Sen metric and associated fields are summarized in Appendix B.6.

2.3 Numerical methods

With the aim of studying the nonlinear dynamics of EMDA in black hole collisions, we express EMDA as a Cauchy problem and evolve the metric using the BSSN formulation [84–86], supplemented by evolution equations for the electromagnetic, dilaton, and axion fields. We also employ puncture gauge conditions [87, 88]. The full equations of motion and the 3 + 1 decomposition are presented in Appendix A.4. Initial data are prepared for head-on black hole collisions and are described below.

2.3.1 Code Description

We perform our black hole merger simulations using Dendro-GR, a well developed numerical relativity framework. Dendro-GR combines an octree-based adaptive mesh infrastructure with automatic code generation for the evolution equations [18, 21, 22]. The computational domain is refined dynamically using wavelet adaptive multiresolution (WAMR) in which the evolved fields are represented on an adaptive hierarchy of octree blocks. Wavelet coefficients computed from the evolved variables provide a local estimate of the truncation error or unresolved field content [18, 89, 90]. Regions where these wavelet coefficients exceed a prescribed tolerance are refined, while regions in which wavelet coefficients remain below a coarsening threshold are de-refined. This tends to concentrate resolution near the black holes, in the wave extraction region, and in regions where the electromagnetic, dilaton, or axion fields develop large spatial gradients, while smoother regions are evolved on coarser grids.

Spatial derivatives are computed using sixth-order finite-difference stencils, and time integration is performed using a fourth-order Runge–Kutta scheme. We include Kreiss–Oliger dissipation to damp poorly resolved high-frequency modes and improve numerical stability [91]. Additional details of the Dendro-GR infrastructure, including its WAMR implementation, scalability, and previous applications to binary black hole evolutions, can be found in [18, 21, 22, 27].

During the evolutions, we monitor the Hamiltonian and momentum constraints together with the electromagnetic divergence constraints throughout the computational domain. These quantities provide a global check on the consistency of the evolution and are used to assess the efficacy of the WAMR refinement parameters. Representative plots of the constraints and their violation are shown in Appendix ??.

We also compute quasilocal quantities on apparent horizons using the BHaHAHA apparent horizon finder [92], which we have extended to evaluate the horizon mass, spin, and electric charge. These

quantities provide independent diagnostics of remnant relaxation and of the conservation of the black hole charges during the evolution.

As a check on our overall numerical strategy, we perform a limited type of convergence study by varying both the wavelet tolerance and maximum refinement level for a short head-on binary evolution. We confirm that with increased numbers of refinement levels and tighter refinement criteria constraint violations are reduced. The details of this study are presented in Appendix ??.

2.3.2 Initial Data

Valid initial data must satisfy the relevant constraint equations. In particular, the Hamiltonian, momentum, and Maxwell constraints are

$$R + K^2 - K_{ij}K^{ij} = 16\pi\rho, \quad (2.2)$$

$$D_j (K^{ij} - \gamma^{ij}K) = 8\pi S^i, \quad (2.3)$$

$$D_i E^i = \rho_E, \quad (2.4)$$

$$D_i B^i = 0, \quad (2.5)$$

where R is the Ricci scalar associated with the spatial metric γ_{ij} , K_{ij} is the extrinsic curvature, K is its trace, and D_i is the covariant derivative compatible with γ_{ij} . The electric and magnetic fields are E^i and B^i with electric charge density, ρ_E , while quantities ρ and S^i are the matter energy and current densities, respectively. The latter, in EMDA, take the form

$$\rho_E \equiv 2\alpha_0 E^i D_i \phi + e^{2\alpha_0 \phi} B^i D_i \kappa \quad (2.6)$$

$$\begin{aligned} \rho \equiv & \Pi^2 + (D\phi)^2 + e^{-2\alpha_0 \phi} (E_i E^i + B_i B^i), \\ & + \frac{1}{4} e^{4\alpha_1 \phi} (\Xi^2 + (D\kappa)^2), \end{aligned} \quad (2.7)$$

$$S^i \equiv 2\Pi D^i \phi + 2e^{-2\alpha_0 \phi} \epsilon^{ijk} E_j B_k + \frac{1}{2} e^{4\alpha_1 \phi} \Xi D^i \kappa. \quad (2.8)$$

where Π and Ξ denote the momenta conjugate to the dilaton ϕ and axion κ , respectively.

Binary black hole initial data in EMDA have not been extensively studied, although several methods have been developed for constructing initial data for charged black holes in related settings [93–97]. In this work, we adapt some of these techniques to the EMDA system by using an elliptic solver based on a modified version of the two-charged-punctures solver introduced in [97]. Our implementation includes contributions from the dilaton and axion fields. A more detailed discussion of the initial data construction is provided in Appendix C.1.


To analyze the radiation emitted, we compute the Newman–Penrose scalar Ψ_4 , along with the electromagnetic radiation. We extract their multipolar components at $r \approx 100M$. Additional details of the waveform extraction procedure are given in Appendix ???. A summary of the initial data parameters used in our simulations is provided in Appendix 2.6.

2.4 Results


We now describe the simulations of two broad classes of binary black holes in EMDA. The first consist of two Kerr–Sen black holes with at least one having nontrivial dilaton and axion fields attached. This is a nice, even stringent test of the persistence of these scalar fields through merger and the settling down of the final object to a new and different Kerr–Sen black hole. The second is the simulation of two Kerr–Newman black holes with perturbative dilaton and axion fields in the vicinity. This latter case is to test whether the merging black holes can acquire these scalar fields and whether the merger then results in a final Kerr–Sen remnant.

2.4.1 Initially Scalarized Black Holes


We perform simulations across a range of binary Kerr–Sen black hole configurations in the initial data space. For the current work, we focus on equal-mass binaries with low to moderate charge and low spin. In particular, the total mass of the system is normalized to $M = 1$. The charge to



(a) Real part of the (2,0) mode of Ψ_4 extracted at $R_{\text{ex}} = 100M$.



(b) Real part of the (2,0) mode of Φ_1 extracted at $R_{\text{ex}} = 100M$.



(c) Real part of the (2,0) mode of Φ_2 extracted at $R_{\text{ex}} = 100M$.

Figure 2.1 The real portions of the (2,0) modes of the gravitational radiation Ψ_4 , electromagnetic radiation Φ_1 , and Φ_2 , extracted at $R_{\text{ex}} = 100M$ for a binary with charge-to-mass ratio $q_i/m_i = 0.1$, dimensionless spin $\chi = 0.4$, and an initial coordinate separation of $16M$.

mass ratio on each black hole ranges between $0.0625 \leq q_i/m_i \leq 0.5$ where m_i and q_i are the initial masses and charges of the individual black holes. For spinning cases, the spins of both black holes are usually taken to be equal and aligned. The spin magnitude of each is taken to lie in the range $0.05 \leq \chi \leq 0.4$. A more complete summary of the parameter values used in our tests is provided in two tables as Appendix C.1.

Figure 2.1 shows the waveforms from one such representative head-on collision of two identical Kerr–Sen black holes. For this system, both BHs have $\chi = 0.4$ and $q/m = 0.1$, are initially separated by $16M$ and fall together from rest. The waveforms are of the gravitational radiation, Ψ_4 and electromagnetic radiation, Φ_1 and Φ_2 . The radiated energy as a function of time is shown in Figure 2.2. In this case, the GW is clearly the dominant form of energy released followed by the electromagnetic radiation. The scalar radiation is much smaller than either the EM or GW radiation. We conclude that there is very little energy radiated in the scalar field channels through the evolution. In all similar experiments, we find that the scalar fields on the black holes persist and do not radiate away through or following the merger. Nontrivial dilaton and axion fields are clearly present and remain attached to the black hole after the merger, as illustrated in Figure 2.3.

We monitor quasilocal quantities on the apparent horizon to verify that the remnant black hole is stable. In particular, we find that the horizon mass, dimensionless spin, and electric charge settle down to approximately constant values. Figure 2.4 shows the percent variation in these quantities after the final black hole has settled into a stationary configuration. In general, these quantities exhibit only small late-time variations, indicating that the remnant has settled to a stable final configuration. Indeed, with these calculated black hole values, we are able to confirm that on comparison with the value of the dilaton on the horizon, the resulting black hole is another, larger mass Kerr–Sen black hole with dilaton and axion fields.






Figure 2.2 Plot of the radiated energy for the case $\chi = 0.4$ and $q_i/m_i = 0.1$. The curves have been aligned such that $t = 0$ corresponds to the global maximum. On this plot the axion and dilaton radiation energy remain at least 3 orders of magnitude smaller than the GW or EM contributions

2.4.2 Initially Unscalarized Black Holes

To further assess the stability of scalar hair, we conducted a series of tests initialized with a Gaussian distribution of axion and dilaton fields centered at the origin, with amplitudes ranging from 10^{-7} up to almost 10^{-2} . Despite not strictly satisfying the constraints, these superposed perturbative scalar fields propagate outward and enable us to examine whether initially unscalarized black holes can dynamically acquire and retain scalar fields. Our results confirm that they do: a Kerr–Newman black hole in EMDA rapidly scalarizes, developing both dilaton and axion hair. In addition, the change to the constraint violations both from the initial perturbations as well as through the subsequent evolution are small enough to leave us confident in our results.



(a) Axion



(b) Dilaton

Figure 2.3 Scalar fields on the $z=0$ plane for the case $\chi=0.4$ and $q_i/m_i=0.1$ Kerr–Sen black holes. We show the amplitudes of the axion, κ (top four panels), and dilaton, ϕ (bottom four panels), fields at the beginning of the evolution (top left), shortly before merger (top right), just after merger (bottom left), and at a later time after merger (bottom right).





Figure 2.4 Percent deviation of the remnant black hole's horizon mass M , electric charge Q , and dimensionless spin χ from their calculated average values: $\langle M \rangle = 0.9682$, $\langle Q \rangle = 0.2001$, and $\langle \chi \rangle = 0.1961$. These quantities are measured on the apparent horizon of the remnant black hole for an initially spinning and charged binary black hole configuration with individual charge-to-mass ratios $q_i/m_i = 0.1$, initial dimensionless spins $\chi = 0.4$, and total mass $M = 1$.

Even in the absence of any initial scalar fields in the background, the system evolves to generate nontrivial scalar structure. This demonstrates that scalarization occurs dynamically and does not require pre-existing scalar hair. The scalarization occurs quickly with initially Kerr–Newman BHs acquiring Kerr–Sen aspects, i.e. scalar hair prior to merger. Mass and electromagnetic charge get radiated with a very small amount of scalar hair, but the final object is a Kerr–Sen BH. Figure 2.5 illustrates the evolution of the scalar fields for the case without initial scalar hair on the BHs.


2.4.3 Constraint and refinement tests

As a consistency check on the evolutions, we monitor the constraint violations throughout the simulations. Representative full evolution constraint plots are shown in Appendix ???. These diagnostics show that the Hamiltonian, momentum, and electromagnetic constraint violations remain controlled over the time scales considered here, including through the merger and ringdown.

We also perform a limited refinement study for a short head-on binary evolution. In this study, we vary both the wavelet tolerance and the maximum number of refinement levels. Both are WAMR related parameters that help dictate the amount and extent of refinement. The wavelet tolerance is varied between 3×10^{-5} and 3×10^{-6} , and we consider a maximum number of refinement levels between 14 and 17, corresponding to grid spacings between $h_{14} = 0.0325$ and $h_{17} = 0.00407$. Additional details are given in Appendix ???. Tightening the refinement criteria reduces the volume weighted Hamiltonian and momentum constraint violations, consistent with the expected behavior of the WAMR refinement scheme. Taken together, the full evolution constraint monitoring and refinement tests indicate that the simulations remain well behaved over the time scales considered here.



(a) Axion



(b) Dilaton

Figure 2.5 Scalar fields on the $z = 0$ plane for Kerr–Newman black holes with initial dimensionless spin $\chi = 0.4$ and charge-to-mass ratio $q_i/m_i = 0.1$. Initially, neither black hole carries dilaton or axion hair. Shown are the amplitudes of the scalar fields κ and ϕ at various stages of the evolution: the initial stage (top left), where both fields begin at zero; a short time later as the fields start to develop (top right); just before merger (bottom left); and some time after merger (bottom right).

2.5 Conclusion

We have simulated a range of different parameters for the head-on collision of Kerr–Sen black holes. We have primarily been concerned with the persistence of scalar hair through this highly nonlinear, dynamical process. In summary, our simulations show that the scalar fields do not radiate away so long as there is a nonzero charge (and spin for the axion). We have found that Kerr–Sen black holes are stable. In particular, we find that if the remnant black hole has a nonzero charge, the dilaton field persists. If it has both nonzero charge and spin, the axion field also persists after merger. These results suggest that Kerr–Sen black holes are stable under the conditions and time frames we considered.

A natural follow-up study would be the analysis of waveforms produced by inspiraling black holes, an area where the available results remain relatively limited. Further, while we have assumed a particular member of the family of EMDA theories in this work, namely a low-energy limit of heterotic string theory, there is a much larger theory space parametrized by the dilaton and axion couplings, α_0 and α_1 , that remains largely unexplored. A more detailed investigation of this theory space could be intriguing; possibly giving rise to broader classes of stable black holes as well as detectable departures from the predicted general relativistic gravitational wave signal.

2.6 Summary of Tests Done

In addition to some of the tests mentioned in the main body of this paper, we have also performed a broader set of simulations to test how the late-time dilaton and axion fields depend on the initial charges and spins of the binary components. In particular, we considered Kerr–Sen binaries with same-sign and opposite-sign charges, spin-aligned and spin-anti-aligned configurations, as well as mergers involving neutral GR black holes. For binaries with opposite charges, the net charge vanishes and the remnant settles to a Kerr black hole, with neither a persistent dilaton nor axion field.

For binaries with same-sign charges but oppositely aligned spins, the axion field decays, while the dilaton field remains nontrivial, corresponding to an EMD-like remnant. We further find that when charged, spinless black holes are merged with Kerr black holes, both axion and dilaton fields persist. Finally, we find that mergers of initially nonscalarized Kerr–Newman black holes produce remnants with both dilaton and axion hair. The configurations considered are summarized in Table 2.6, where a green check mark indicates that the corresponding field remains persistent and nontrivial at late times.

Binary black hole configurations

(a) Kerr–Sen Same-Sign Charge and Spin-Aligned Black Hole Tests

(χ_1, q_1)	(χ_2, q_2)	Dilaton ϕ	Axion κ
$\chi_1 = 0.10, q_1 \in \{0.0624, 0.125, 0.25, 0.50\}$	$\chi_2 = 0.10, q_2 \in \{0.0624, 0.125, 0.25, 0.50\}$	✓	✓
$\chi_1 = 0.20, q_1 \in \{0.0624, 0.125, 0.25, 0.50\}$	$\chi_2 \in \{0.10, 0.20\}, q_2 \text{ as above}$	✓	✓
$\chi_1 = 0.40, q_1 \in \{0.0624, 0.125, 0.25, 0.50\}$	$\chi_2 \in \{0.10, 0.20, 0.40\}, q_2 \in \{0.0624, 0.125, 0.25, 0.50\}$	✓	✓

(b) Collisions with Other Black Holes

Black Hole 1 (q, χ)	Black Hole 2 (q, χ)	ϕ	κ
Kerr–Sen (0.5, 0.4)	Kerr–Sen (0.5, 0.4)	✓	✓
Kerr–Sen (0.5, 0.4)	Kerr (0.0, 0.4)	✓	✓
Kerr–Sen (0.5, 0.4)	Kerr–Sen (−0.5, 0.4)	✗	✗
Kerr–Sen (0.5, −0.4)	Kerr–Sen (0.5, 0.4)	✓	✗
EMD (0.5, 0.0)	Kerr (0.0, 0.4)	✓	✓
KN (0.5, 0.4) [†]	KN (0.5, 0.4) [†]	✓	✓

[†] These configurations were initialized as nonscalarized Kerr–Newman black holes.

Abbreviations: EMD denotes Einstein–Maxwell–dilaton, corresponding here to a charged black hole with dilaton hair;

KN denotes Kerr–Newman, corresponding to a black hole with charge and spin; and Kerr–Sen denotes a rotating charged black hole with dilaton and axion hair.

Chapter 3

EMDA Chapter 2

3.1 Introduction

Gravitational-wave observations now provide direct access to the strong-field, dynamical regime of compact-binary coalescence. As detector sensitivities improve and high-signal-to-noise-ratio detections become more likely, the late inspiral, merger, and ringdown will offer increasingly stringent tests of the general-relativistic description of black holes [3, 10, 31, 34, 59]. Recent detections have reached signal-to-noise ratios as high as 80, enabling tests of Hawking’s area law and improved measurements of the remnant black hole [98]. These developments motivate numerical studies of how specific beyond-GR effects could appear in the late-inspiral, merger, and ringdown waveform.

A central difficulty in such tests is that the merger regime is both highly nonlinear and theory dependent. For many alternative theories of gravity, analytic approximations are insufficient near merger, while numerical simulations require a well-posed initial-value formulation, stable gauge choices, and controlled constraint behavior. Considerable effort has been devoted to developing formulations and approximation schemes that permit nonlinear simulations beyond GR [49, 50, 57, 63, 65]. Examples include studies of dynamical Chern–Simons gravity [35, 36], Einstein-dilaton-Gauss-Bonnet theory [44, 66], modified gauge choices in Einstein-scalar-Gauss-Bonnet theory [42, 67], the “fixing the equations” approach [68], and auxiliary-field formulations for higher-curvature systems, including quadratic gravity [51, 69–72]. See also Ref. [73] for a more detailed comparison of these approaches.

In this chapter, we focus on Einstein–Maxwell–dilaton–axion (EMDA) theory. EMDA extends Einstein–Maxwell theory by coupling the electromagnetic field to a dilaton and an axion, and it arises as a low-energy limit of heterotic string theory [74, 75]. The theory admits rotating charged black-hole solutions, such as the Kerr–Sen spacetime, which differs from Kerr–Newman by the presence of nontrivial dilaton and axion fields. Previous work has investigated the stability of black holes and black-hole hair in EMDA theory [79, 99], and black-hole shadow studies using

Event Horizon Telescope observations have not ruled out Kerr–Sen black holes [80, 81]. These features make EMDA a useful setting in which to study how additional matter fields can influence compact-binary dynamics and gravitational radiation.

Charged binary black holes have also been studied in Einstein–Maxwell gravity [95–97, 100–102]. These simulations show that electromagnetic charge can modify the orbital dynamics, emitted radiation, and recoil of the remnant. In one analysis, the GW150914 signal was found to be compatible with charge-to-mass ratios as large as $Q/M \sim 0.3$ [100]. These studies demonstrate that charge alone can alter the inspiral, merger, radiation content, and remnant properties. In EMDA theory, charge also sources additional scalar and pseudoscalar fields, so charged binaries provide a natural setting in which to study how electromagnetic and beyond-GR matter jointly affect the gravitational waveform.

The goal of this chapter is to take a step toward an exploratory waveform catalog for charged binary black holes in EMDA theory. We simulate inspiraling charged binary black holes and compare the resulting gravitational waveforms with neutral general-relativistic reference waveforms having the same mass ratio and spin. The cases considered here include charge-to-mass ratios larger than those expected for astrophysical black holes [103, 104]. Our aim is therefore not to construct immediately realistic astrophysical templates, but to identify regimes in which charge and the additional EMDA fields produce appreciable changes in the binary dynamics and gravitational-wave signal.

The catalog considered here samples several choices of mass ratio, spin, charge magnitude, and charge configuration. By comparing each charged EMDA waveform with its corresponding neutral GR reference waveform, we ask how waveform differences emerge as one moves through the EMDA parameter space and how these differences depend on the binary parameters.

These comparisons should be interpreted as fixed-parameter waveform comparisons rather than as a complete parameter-estimation study. Compact-binary waveforms are highly degenerate,

and differences between an EMDA waveform and a GR waveform may be partially mimicked by varying intrinsic binary parameters such as the mass ratio, spins, total mass scale, or eccentricity. Moreover, because the comparisons in this chapter are made against neutral GR rather than charged Einstein–Maxwell binaries, the resulting differences should be understood as the combined effect of charge, modified orbital dynamics, and the additional EMDA fields, rather than as isolated dilaton or axion signatures.

The catalog presented here is therefore intended as an initial step toward a more complete study of EMDA binary black-hole phenomenology. In particular, it identifies regions of parameter space where charged EMDA binaries produce appreciable waveform differences relative to neutral GR and motivates future comparisons among neutral GR, Einstein–Maxwell, and EMDA binaries.

3.2 Einstein–Maxwell–dilaton–axion Theory

We consider Einstein–Maxwell–dilaton–axion (EMDA) theory, in which the Einstein–Maxwell system is supplemented by a scalar dilaton field ϕ and a pseudoscalar axion field κ . With the conventions used here, the action is

$$S = \int d^4x, \sqrt{-g} \left[\frac{R}{8\pi} - 2\nabla_a \phi \nabla^a \phi - e^{-2\alpha_0 \phi} F_{ab} F^{ab} - \frac{1}{2} e^{4\alpha_1 \phi} \nabla_a \kappa \nabla^a \kappa - \kappa F_{ab} ({}^*F)^{ab} \right], \quad (3.1)$$

where R is the Ricci scalar, F_{ab} is the $U(1)$ field strength, and $({}^*F)_{ab}$ is its dual. The constants α_0 and α_1 determine the coupling strengths between the electromagnetic field and the dilaton and axion fields.

Different choices of α_0 and α_1 correspond to different members of this class of theories. The parameter α_0 controls the exponential coupling between the dilaton and the electromagnetic field, while α_1 controls the exponential coupling of the dilaton to the axion kinetic term. For example, the

choice $(\alpha_0, \alpha_1) = (\sqrt{3}, 0)$ gives the Kaluza–Klein value of the dilaton coupling [105, 106]. In the simulations presented here, we fix

$$(\alpha_0, \alpha_1) = (1, 1), \quad (3.2)$$

which corresponds to the coupling values associated with the low-energy limit of heterotic string theory [74, 75].

For this choice of couplings, the theory admits rotating charged black-hole solutions of the Kerr–Sen type [74]. These solutions reduce to Kerr in the limit of vanishing electromagnetic charge, but for nonzero charge the black hole is accompanied by a nontrivial dilaton field and, in the rotating case, a nontrivial axion field. The structure of these fields is known analytically for a single isolated black hole; the solution is summarized in Appendix B.6.

To simulate inspiraling binary black holes, we construct approximate charged binary black-hole initial data and evolve the full EMDA system numerically. This allows us to compare the resulting gravitational waveforms against neutral general-relativistic reference waveforms and to identify how the combined electromagnetic, dilaton, and axion fields influence the inspiral and merger.

3.3 Numerical Methods

To study the strong-field dynamics and gravitational radiation from inspiraling charged black-hole binaries in Einstein–Maxwell–dilaton–axion (EMDA) theory, we evolve the full EMDA system as a Cauchy initial-value problem. The spacetime metric is evolved using the BSSN formulation [84–86], together with puncture gauge conditions [87, 88]. The electromagnetic, dilaton, and axion fields are evolved using the matter evolution equations obtained from the 3 + 1 decomposition of the EMDA equations of motion. The full evolution system is summarized in Appendix A.4.

The simulations are initialized with approximate charged binary black-hole initial data and evolved through the late inspiral, merger, and ringdown. During the evolution, we extract gravitational

radiation and monitor the matter fields, constraints, and quasilocal horizon quantities in order to assess the dynamics and numerical consistency of the solutions.

3.3.1 Numerical implementation

For the simulations presented here, we evolve the EMDA system using *Dendro-GR*, the numerical relativity code also used in Ref. [99]. *Dendro-GR* is an adaptive mesh-refinement code based on an octree representation of the computational domain. The mesh hierarchy is adjusted dynamically using wavelet adaptive multiresolution, in which wavelet coefficients provide a local measure of unresolved field content and are used to determine where refinement or coarsening is required [18, 89, 90]. This allows resolution to be concentrated in regions where the evolved fields contain sharp gradients or otherwise require finer spatial resolution. In the simulations considered here, this places additional resolution near the black holes and in regions with nontrivial electromagnetic, dilaton, or axion structure, while avoiding uniform refinement of the full computational domain.

Spatial derivatives are evaluated using sixth-order finite-difference operators, and the evolution is advanced in time with a fourth-order Runge–Kutta method. We also add Kreiss–Oliger dissipation of the corresponding order to suppress poorly resolved high-frequency numerical modes. Further details on the *Dendro-GR* infrastructure, including its adaptive wavelet refinement strategy, code-generation framework, scalability, and previous applications to binary black-hole simulations, can be found in Refs. [18, 21, 22, 27]. In particular, *Dendro-GR* has been shown to perform well for intermediate-mass-ratio inspiral simulations, where adaptive refinement is especially useful for resolving disparate length scales [22, 27].

Throughout each evolution, we monitor the Hamiltonian and momentum constraints, as well as the electromagnetic divergence constraints. These quantities are used to track the consistency of the numerical solution and to assess whether the chosen refinement parameters adequately resolve the relevant dynamics. Representative constraint behavior is shown in Appendix E.3.1.

We also extract quasilocal horizon quantities using a modified version of the BHaHAHA apparent-horizon finder [92]. In addition to locating the apparent horizons, this modified version is used to compute the horizon mass, spin, and electric charge during the evolution.

3.3.2 Controlling the Eccentricity

In this work, we focus on approximately quasicircular inspirals. For uncharged binary black holes, post-Newtonian methods can be used to choose initial momenta that reduce the orbital eccentricity. In the EMDA system, however, the presence of electromagnetic, dilaton, and axion fields complicates the construction of low-eccentricity initial data. Post-Newtonian treatments have been developed for Einstein–Maxwell theory and for several scalar-modified theories of gravity [107–110]. Here we adopt a simpler prescription based on the charged-binary argument used in Ref. [111].

At Newtonian order, both the gravitational and electromagnetic interactions are central forces. For two charged compact objects, the Coulomb interaction modifies the effective strength of the attractive force. Denoting the charge-to-mass ratio of each black hole by

$$\lambda_i = \frac{Q_i}{M_i}, \quad (3.3)$$

the leading-order effect of the electromagnetic interaction can be incorporated through the replacement

$$G \rightarrow \tilde{G} = G (1 - \lambda_1 \lambda_2). \quad (3.4)$$

In units where $G = 1$, this becomes

$$\tilde{G} = 1 - \lambda_1 \lambda_2. \quad (3.5)$$

Thus, like-signed charges reduce the effective attraction, while oppositely signed charges increase it. We first compute the momenta required for the corresponding uncharged binary to be approximately quasicircular using post-Newtonian estimates, and then rescale the orbital momenta according to the

charge-dependent effective coupling. At leading Newtonian order, this corresponds to scaling the tangential momentum by

$$p_t \rightarrow p_t \sqrt{1 - \lambda_1 \lambda_2}. \quad (3.6)$$

This prescription assumes that the dominant correction to the initial orbital parameters comes from the electromagnetic interaction; possible direct corrections from the dilaton and axion fields are not included in the initial momentum estimate.

The initial momenta for the corresponding uncharged binaries are computed using the NRPyPN package [112]. This package provides post-Newtonian estimates for low-eccentricity binary black-hole initial data compatible with Bowen–York/TwoPunctures data [113, 114]. We then apply the charge-dependent rescaling described above to obtain the momenta used in the charged evolutions.

Although approximate, this prescription yields eccentricities that are adequate for the exploratory comparisons performed here, as illustrated in Figs. 3.1 and 3.2. The charge configuration also has a significant effect on the inspiral time, as expected from the change in the effective attractive force. Like-signed charges tend to weaken the attraction and delay the merger, while oppositely signed charges tend to strengthen the attraction and shorten the inspiral.

We estimate the orbital eccentricity using an estimator based on the orbital frequency [114],

$$\varepsilon_\Omega(t) = \frac{1}{2} \left[\frac{\Omega(t) - \Omega_{e=0}(t)}{\Omega_{e=0}(t)} \right], \quad (3.7)$$

where $\Omega(t)$ is the measured orbital frequency and $\Omega_{e=0}(t)$ denotes the corresponding non-eccentric reference trend.

We use an orbital-frequency-based estimator because the orbital frequency is less sensitive to gauge effects than the coordinate separation, although it is still not fully gauge invariant. We track the coordinate positions of each black hole during the inspiral, compute the coordinate separation, and then compute the orbital frequency. The orbital-frequency data are then fit using `NonlinearModelFit` in `Mathematica` with a post-Newtonian-inspired fitting model [114].





(a) $q = 1$: black-hole separation.(b) $q = 1$: orbital trajectories.(c) $q = 2$: black-hole separation.(d) $q = 2$: orbital trajectories.

Figure 3.1 Orbital dynamics for the $q = 1$ and $q = 2$ configurations. For each mass ratio, the left panel shows the coordinate separation as a function of time, while the right panel shows representative coordinate trajectories in the orbital plane. The charge configuration affects both the inspiral time and the orbital phasing through the modified effective attraction.





(a) $q = 4$: black-hole separation.(b) $q = 4$: orbital trajectories.(c) $q = 8$: black-hole separation.(d) $q = 8$: orbital trajectories.

Figure 3.2 Orbital dynamics for the $q = 4$ and $q = 8$ configurations. For each mass ratio, the left panel shows the coordinate separation as a function of time, while the right panel shows representative coordinate trajectories in the orbital plane.

When fitting the orbital-frequency data, it is useful to include as many orbits as possible in order to reduce the uncertainty in the eccentricity estimate. However, the early part of each simulation contains gauge-settling effects that are not associated with the physical orbital eccentricity, while the late part of the simulation is affected by the onset of plunge. We therefore exclude the first $\sim 200M$ of the simulation and choose the end of the fitting window before the rapid late-inspiral growth of the orbital frequency becomes dominant. In practice, this typically corresponds to ending the fit when the coordinate separation is of order $6M$.

The measured orbital eccentricities for the catalog runs are shown in Table 3.3. The quoted eccentricities are extrapolated from the fit to $t = 0M$. The initial coordinate separations are $r = 10M$, $r = 9M$, and $r = 8M$ for the $q = 1$, $q = 2$, and $q = 4$ or $q = 8$ configurations, respectively. Most of the lower-spin and lower-charge configurations have eccentricities of order 10^{-3} , while the largest eccentricities occur in the high-spin and high-charge equal-mass cases.

Figure 3.3 shows an example of the orbital-frequency eccentricity fit for one catalog run. This case has $q = 1$, $\chi = 0.4$, $\lambda = 0.3$, and charge configuration $++$. The fit uses approximately four orbits, with $t_{\min} = 200M$ and $t_{\max} = 1300M$. The measured eccentricity extrapolated to $t = 0M$ is $\varepsilon_{\Omega} = (14.44 \pm 0.33) \times 10^{-3}$. This comparatively large eccentricity is consistent with the fact that this is a high-charge, like-signed configuration, for which the reduced effective attraction makes the initial momentum prescription less accurate.

In principle, lower eccentricities could be obtained through an iterative eccentricity-reduction procedure, in which the initial momenta are corrected after an initial eccentricity measurement and the simulation is repeated. We do not perform iterative eccentricity reduction for this exploratory catalog. Standard post-Newtonian eccentricity-reduction methods may also need modification in the EMDA case, since the electromagnetic, dilaton, and axion fields can affect the orbital dynamics. Developing eccentricity-reduction procedures that incorporate these effects is left for future work.




Figure 3.3 Measured orbital-frequency eccentricity and fitted eccentricity model for a binary black-hole inspiral with $q = 1$, $\chi = 0.4$, $\lambda = 0.3$, and charge configuration $++$. The fit is performed from $t_{\min} = 200M$ to $t_{\max} = 1300M$, excluding early gauge-settling effects and ending before the onset of plunge. The measured eccentricity extrapolated to $t = 0M$ is $\varepsilon_{\Omega} = (14.44 \pm 0.33) \times 10^{-3}$.

3.3.3 Waveform Mismatch and Distinguishability

Following Refs. [100, 111, 115–117], we use waveform overlaps and mismatches to quantify differences between the simulated gravitational-wave signals.

A direct comparison between charged EMDA and neutral GR waveforms does not, by itself, isolate the contribution of the dilaton and axion fields. Part of the measured difference may arise from the electromagnetic charge and the associated change in the orbital dynamics, rather than from the additional EMDA fields alone. A more complete study would compare EMDA waveforms not only with neutral GR waveforms, but also with charged Einstein–Maxwell evolutions having the same charge configurations. Such a comparison is beyond the scope of the present work. Our goal here is instead to quantify the fixed-parameter waveform differences between charged EMDA binaries and their neutral GR counterparts.

To quantify the difference between two gravitational-wave signals, we compute the mismatch

$$\mathcal{M}(h_1, h_2) = 1 - \max_{\Delta t, \Delta \varphi} \mathcal{O}(h_1, h_2), \quad (3.8)$$

where the maximum is taken over relative time shifts Δt and polarization-angle shifts $\Delta\varphi$. The overlap O is defined by

$$O(h_1, h_2) = \frac{(h_1, h_2)}{\sqrt{(h_1, h_1)(h_2, h_2)}}, \quad (3.9)$$

where (h_1, h_2) denotes the detector-noise-weighted inner product between the two waveforms. In practice, this inner product is evaluated over the finite frequency range used in the analysis, with the same preprocessing choices applied to each waveform pair.

The comparison performed here does not maximize over intrinsic binary parameters such as the masses, spins, eccentricity, or charge. These degeneracies could partially absorb the waveform differences found in a fixed-parameter comparison and would likely make charge-induced deviations more difficult to identify in a full parameter-estimation study. The mismatch values reported below should therefore be interpreted as a measure of waveform separation at fixed intrinsic parameters, rather than as evidence for a unique observational signature of EMDA theory.

As a rough estimate of distinguishability, we associate a mismatch $\mathcal{M}$ with the critical signal-to-noise ratio

$$\rho_{\text{crit}} \simeq \frac{1}{\sqrt{2\mathcal{M}}}. \quad (3.10)$$

This estimate gives the signal-to-noise ratio at which two waveforms with mismatch $\mathcal{M}$ would be expected to become distinguishable in an idealized fixed-parameter comparison. It should be interpreted cautiously, since it does not account for correlations with the full set of binary parameters, detector systematics, numerical error, or the possibility that a nearby GR waveform with different intrinsic parameters could provide a comparable fit.

3.4 Results

We compare charged EMDA waveforms with neutral GR reference waveforms for a representative set of binary black-hole configurations. The purpose of this comparison is to determine how the

waveform differences depend on charge magnitude, charge configuration, mass ratio, and spin. In each case, the charged EMDA waveform is compared against a neutral reference waveform with the same mass ratio and spin.

The simulation catalog considered in this work is summarized in Table 3.1. For each mass ratio $q = m_1/m_2$, spin χ , and charge magnitude λ , we consider the neutral case, $Q_1 = Q_2 = 0$, together with several charged configurations. The charged cases are denoted by ++, +−, and +0, corresponding respectively to like-signed charges, oppositely signed charges, and configurations in which only one black hole is charged.

Representative peak-aligned waveforms are shown in Fig. 3.4. The figure compares the strain component rh_+ for selected charged EMDA configurations against the corresponding neutral GR reference waveform. The waveforms are aligned at the peak amplitude in order to emphasize differences in the late inspiral, merger, and ringdown morphology. Even after this alignment, the charged and neutral waveforms show visible phase and amplitude differences, indicating that charge and the additional EMDA fields can modify the binary dynamics prior to merger as well as the waveform near peak emission.

The full set of waveform mismatches is computed using the numerical-relativity analysis toolkit `kuibit` [118], and the results are given in Table 3.2. The mismatch values generally increase with charge magnitude, although the trend is not monotonic across all spin and charge configurations. For example, in the $q = 1$, $\chi = 0.4$, ++ sequence, the mismatch increases from 2.60×10^{-4} for $\lambda = 0.1$ to 5.28×10^{-3} for $\lambda = 0.3$. Similar growth with charge is seen in several other equal-mass configurations.

The charge configuration also plays an important role. The ++, +−, and +0 cases do not produce equivalent waveform differences, even at the same charge magnitude. In many cases, the singly charged +0 configurations remain closer to the neutral reference waveform than the corresponding two-charged configurations. This is consistent with the expectation that the strength and symmetry




Figure 3.4 Peak-aligned comparisons of the gravitational-wave strain component rh_+ for representative charged EMDA configurations against the corresponding neutral GR reference waveforms. In each panel, the time coordinate is shifted so that the waveform peak occurs at $t - t_{\text{peak}} = 0$. The remaining differences after peak alignment reflect changes in the accumulated phase, merger time, and waveform morphology.

of the charge distribution influence both the orbital dynamics and the matter-field content of the spacetime. Oppositely signed charges can also produce comparatively large mismatches, since the electromagnetic interaction increases the effective attraction and changes the inspiral rate.

The $\chi = 0.8$ equal-mass configurations are somewhat anomalous compared with the lower-spin cases. While the mismatch generally grows with charge for the lower-spin sequences, the $\chi = 0.8$ cases show larger and less monotonic variations across the different charge configurations. These runs also have noticeably larger eccentricities, especially at the highest charge magnitudes, suggesting that the approximate charged initial-data prescription is being pushed beyond the regime where it

gives uniformly low-eccentricity inspirals. We therefore include these cases as useful indicators of where large waveform differences can occur, but we do not interpret their detailed mismatch hierarchy as a clean physical trend.

The high-spin configurations nevertheless show the largest waveform separations in the catalog. For the $q = 1$, $\chi = 0.8$ cases, some mismatches reach the 10^{-2} – 10^{-1} level. These large values indicate that the charged and neutral waveforms are substantially separated at fixed intrinsic parameters. However, because these cases are also more sensitive to eccentricity and initial-data systematics, they should not be interpreted as isolated signatures of the dilaton or axion fields.

The critical signal-to-noise ratio estimate associated with the mismatch values spans approximately $\rho_{\text{crit}} \sim 2.3$ to $\rho_{\text{crit}} \sim 537$ across the catalog. These values should be understood as fixed-parameter distinguishability estimates: they indicate the signal-to-noise ratio at which the charged EMDA waveform and the corresponding neutral GR waveform would become distinguishable if the intrinsic binary parameters were held fixed. In a full parameter-estimation study, changes in the masses, spins, eccentricity, or other parameters could partially absorb these differences. The mismatch values therefore quantify waveform separation within this catalog, rather than providing direct observational bounds on EMDA theory.

3.5 Conclusion

In this chapter, we presented an exploratory waveform catalog for charged binary black-hole simulations in Einstein–Maxwell–dilaton–axion (EMDA) theory. We compared the resulting charged EMDA waveforms with neutral general-relativistic reference waveforms having the same mass ratio and spin. Across much of the catalog, the waveform differences increase with the charge magnitude, indicating that charge and the additional EMDA fields can produce appreciable changes in the gravitational-wave signal. The trend is clearest in the lower-spin configurations, while the

highest-spin cases show larger and less monotonic differences that are likely more sensitive to eccentricity and the approximate charged initial-data prescription.

Using the mismatch as a rough fixed-parameter distinguishability estimate, the largest waveform separations in the catalog correspond to critical signal-to-noise ratios, with the smallest values approximately $\rho_{\text{crit}} \sim 2.3$. These values indicate that the largest charged EMDA deviations from the corresponding neutral GR waveforms are substantial at fixed intrinsic parameters. However, they should not be interpreted as direct observational thresholds or constraints on EMDA theory. A full parameter-estimation study would be required to determine whether these differences could be absorbed by changes in the masses, spins, eccentricity, total mass scale, or other binary parameters.

The waveform differences found here should also be interpreted with care because the comparison is made against neutral GR rather than against charged Einstein–Maxwell binaries. As a result, the reported mismatches include the combined effect of electric charge, modified orbital dynamics, and the additional dilaton and axion fields. They cannot be attributed uniquely to the dilaton or axion fields alone. A more complete comparison would include corresponding Einstein–Maxwell simulations with the same charge configurations, allowing the purely electromagnetic contribution to be separated from the additional EMDA-field effects.

These results demonstrate that EMDA theory provides a useful numerical-relativity testbed for studying charged beyond-GR effects in binary black-hole mergers. Even within this exploratory catalog, charged EMDA binaries can produce waveform differences that are large enough to motivate further study. Natural next steps include larger parameter surveys, improved charged initial data, iterative eccentricity reduction, longer inspirals, convergence studies of the waveform mismatches, and direct comparisons among neutral GR, charged Einstein–Maxwell, and charged EMDA binaries.

3.6 Simulation Catalog

The simulations used in this chapter are summarized in Table 3.1. The catalog samples several mass ratios, spin values, charge magnitudes, and charge configurations. For each mass ratio and spin, we include a neutral reference simulation and compare the charged EMDA evolutions against that corresponding neutral case.

3.7 Measured Eccentricities

Table 3.3 summarizes the measured orbital eccentricities for the EMDA waveform catalog. The values are obtained from the orbital-frequency eccentricity estimator described in Sec. 3.3.2 and are reported as $(\varepsilon_\Omega \pm \delta\varepsilon_\Omega) \times 10^3$.

The eccentricity estimates for the $q = 8$ configurations are poorly constrained compared with the rest of the catalog, as indicated by their large uncertainties. We therefore use these values only as rough indicators of the orbital behavior and do not place significant weight on them when interpreting the mismatch trends.

Table 3.1 EMDA binary black-hole simulation catalog. Here $q = m_1/m_2$ denotes the mass ratio, χ denotes the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$ denotes the charge-to-mass ratio of black hole i . The neutral case corresponds to $Q_1 = Q_2 = 0$. For charged configurations, the magnitude of the nonzero charge-to-mass ratios is denoted by λ , with the sign structure indicated by the charge configuration. The configurations ++, +−, and +0 denote like-signed charges, oppositely signed charges, and cases in which only one black hole is charged, respectively. Because the neutral configuration is independent of λ , it is listed only once for each mass ratio and spin.

Mass Ratio q	Spin χ	Charge Magnitude λ	Neutral	++	+−	+0
1	0.0	0.1	✓	✓	✓	✓
1	0.0	0.2	−	✓	✓	✓
1	0.0	0.3	−	✓	✓	✓
1	0.4	0.1	✓	✓	✓	✓
1	0.4	0.2	−	✓	✓	✓
1	0.4	0.3	−	✓	✓	✓
1	0.8	0.1	✓	✓	✓	✓
1	0.8	0.2	−	✓	✓	✓
1	0.8	0.3	−	✓	✓	✓
2	0.0	0.05	✓	✓	✓	✓
2	0.0	0.15	−	✓	✓	✓
2	0.4	0.05	✓	✓	✓	✓
2	0.4	0.15	−	✓	✓	✓
4	0.4	0.05	✓	✓	✓	✓
4	0.4	0.10	−	✓	✓	✓
8	-0.2	0.10	✓	✓	−	−

Table 3.2 Mismatch of charged EMDA binary black-hole waveforms relative to the corresponding neutral reference waveform Q0. Here $q = m_1/m_2$, χ is the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$. The column $|\lambda_i|$ gives the magnitude of the nonzero charge-to-mass ratio used in each charged configuration. The columns ++, +−, and +0 denote like-signed charges, oppositely signed charges, and configurations in which only one black hole is charged, respectively. The mismatch is computed from the $(\ell, m) = (2, 2)$ gravitational waveform using the numerical-relativity analysis package *kuibit*, after optimizing over relative time and polarization shifts. The values in parentheses give the approximate fixed-parameter critical signal-to-noise ratio, $\rho_{\text{crit}} \simeq 1/\sqrt{2\mathcal{M}}$. These values should be interpreted as rough waveform-separation estimates rather than full observational detection thresholds.

Mass Ratio q	Spin χ	Charge Magnitude $ \lambda_i $	++	+−	+0
1	0.0	0.10	3.53×10^{-4} ; (37.6)	2.74×10^{-4} ; (42.8)	1.55×10^{-4} ; (56.8)
1	0.0	0.20	6.71×10^{-4} ; (27.3)	1.45×10^{-3} ; (18.5)	6.91×10^{-4} ; (26.9)
1	0.0	0.30	1.94×10^{-3} ; (16.0)	5.32×10^{-3} ; (9.7)	7.54×10^{-4} ; (25.8)
1	0.4	0.10	2.60×10^{-4} ; (43.9)	2.87×10^{-4} ; (41.8)	4.08×10^{-4} ; (35.0)
1	0.4	0.20	7.10×10^{-4} ; (26.5)	7.40×10^{-3} ; (8.2)	7.42×10^{-4} ; (26.0)
1	0.4	0.30	5.28×10^{-3} ; (9.7)	2.90×10^{-2} ; (4.1)	1.04×10^{-3} ; (21.9)
1	0.8	0.10	2.77×10^{-4} ; (42.5)	9.44×10^{-4} ; (23.0)	1.03×10^{-3} ; (22.0)
1	0.8	0.20	4.66×10^{-2} ; (3.3)	9.30×10^{-2} ; (2.3)	1.67×10^{-2} ; (5.5)
1	0.8	0.30	6.19×10^{-2} ; (2.8)	2.53×10^{-2} ; (4.4)	7.99×10^{-3} ; (7.9)
2	0.0	0.05	1.58×10^{-3} ; (17.8)	1.34×10^{-3} ; (19.3)	2.56×10^{-5} ; (139.7)
2	0.0	0.15	1.83×10^{-3} ; (16.5)	3.21×10^{-3} ; (12.5)	7.47×10^{-4} ; (25.9)
2	0.4	0.05	7.27×10^{-5} ; (82.9)	1.33×10^{-4} ; (61.3)	1.73×10^{-6} ; (537.4)
2	0.4	0.15	6.17×10^{-4} ; (28.5)	2.34×10^{-4} ; (46.2)	7.58×10^{-5} ; (81.2)
4	0.4	0.05	5.10×10^{-5} ; (99.0)	7.34×10^{-5} ; (82.5)	7.17×10^{-5} ; (83.5)
4	0.4	0.10	3.17×10^{-4} ; (39.7)	1.82×10^{-4} ; (52.5)	8.80×10^{-5} ; (75.4)
8	-0.2	0.10	1.14×10^{-3} ; (21.0)	—	—

Table 3.3 Measured orbital eccentricities for the EMDA waveform catalog. The entries are reported as $(\varepsilon_\Omega \pm \delta\varepsilon_\Omega) \times 10^3$. Here $q = m_1/m_2$ denotes the mass ratio, χ denotes the dimensionless spin parameter, and $\lambda_i = Q_i/M_i$ denotes the charge-to-mass ratio of black hole i . The neutral case corresponds to $Q_1 = Q_2 = 0$. For charged configurations, the magnitude of the nonzero charge-to-mass ratios is denoted by λ , with the sign structure indicated by the charge configuration. The configurations ++, +−, and +0 denote like-signed charges, oppositely signed charges, and cases in which only one black hole is charged, respectively.

Mass Ratio q	Spin χ	Charge Magnitude λ	Neutral	++	+−	+0
1	0.0	0.10	1.59 ± 0.21	2.57 ± 0.17	1.97 ± 0.27	2.54 ± 0.23
1	0.0	0.20	–	6.26 ± 0.26	3.52 ± 0.35	4.41 ± 0.24
1	0.0	0.30	–	14.99 ± 0.40	2.08 ± 0.36	5.28 ± 0.45
1	0.4	0.10	1.39 ± 0.10	1.80 ± 0.11	1.30 ± 0.13	1.26 ± 0.11
1	0.4	0.20	–	5.17 ± 0.18	1.04 ± 0.22	2.68 ± 0.18
1	0.4	0.30	–	14.44 ± 0.33	4.89 ± 0.53	4.19 ± 0.19
1	0.8	0.10	3.47 ± 0.23	3.23 ± 0.18	2.31 ± 0.23	2.31 ± 0.21
1	0.8	0.20	–	4.95 ± 0.64	5.44 ± 0.71	4.57 ± 0.27
1	0.8	0.30	–	44.75 ± 0.70	22.70 ± 6.36	13.03 ± 2.02
2	0.0	0.05	2.16 ± 0.60	1.98 ± 0.22	1.22 ± 0.62	1.62 ± 0.31
2	0.0	0.15	–	4.65 ± 0.36	2.18 ± 0.33	3.05 ± 0.37
2	0.4	0.05	1.68 ± 0.11	1.46 ± 0.24	1.78 ± 0.14	1.48 ± 0.17
2	0.4	0.15	–	5.19 ± 0.15	1.35 ± 0.24	2.09 ± 0.18
4	0.4	0.05	1.66 ± 0.43	1.44 ± 0.32	1.93 ± 0.34	1.49 ± 0.50
4	0.4	0.10	–	1.01 ± 0.09	0.81 ± 0.21	0.79 ± 0.21
8	-0.2	0.10	4.88 ± 61.91	1.45 ± 5.10	–	–

Chapter 4

EMDA Chapter 3

4.1 Introduction

The growing number of detected binary black-hole mergers provides an increasingly broad sample of gravitational-wave signals with which to test general relativity in the strong-field, dynamical regime. Continued improvements to the LIGO, Virgo, and KAGRA detectors are expected to further increase the rate of observed compact-binary mergers, expanding the available catalog of gravitational-wave events [24, 25]. As this catalog grows, gravitational-wave observations provide an increasingly powerful way to search for small deviations from general relativity and to constrain alternative theories of gravity [3, 34, 59].

Numerical relativity plays an important role in this effort because the merger regime is highly nonlinear and theory dependent. However, numerical-relativity models for alternative theories of gravity are less mature than their general-relativistic counterparts. Simulating such theories presents several challenges. Some modified theories suffer from pathologies such as ill-posed evolution systems or ghost-like degrees of freedom associated with higher-derivative terms [119–121]. Numerical studies of these theories therefore require formulations in which the initial-value problem can be posed consistently and the subsequent evolution remains well behaved. In particular, one must construct suitable initial data and evolve the system in a way that controls the constraints.

Many tests of gravity can be understood as constraints on additional structure that is absent in vacuum general relativity. Previous work has used gravitational-wave observations to bound higher-curvature corrections, placing limits on the length scales over which such terms can become important [3, 33, 122]. Other studies have considered constraints on black-hole charge [111, 123], as well as tests based on black-hole shadows [80, 81]. In this chapter, we investigate another possible extension: Einstein–Maxwell–dilaton–axion (EMDA) theory. EMDA theory contains, in addition to the metric and electromagnetic field, a scalar dilaton field and a pseudoscalar axion field. These additional degrees of freedom can modify the structure of charged black holes and may leave observable imprints on binary mergers and their emitted radiation [47, 74, 75, 105].

Analytic black-hole solutions in EMDA-related theories are known for special choices of the coupling parameters. These include the Kerr–Sen black hole for $\alpha_0 = \alpha_1 = 1$, Einstein–Maxwell–dilaton black holes for $\alpha_1 = 0$, and Kaluza–Klein black holes for $\alpha_0 = \sqrt{3}$ [74, 75, 105, 106]. The stability of Kerr–Sen black holes has recently been studied in both perturbative and fully nonlinear settings [79, 99], and previous numerical work has explored portions of the EMD parameter space [47]. However, less is known about how varying the EMDA coupling parameters affects the nonlinear dynamics of binary black-hole mergers.

Here, we focus primarily on the $\alpha_0 = \alpha_1$ sector of the EMDA parameter space. By varying this common coupling, we study how the strength of the dilaton and axion interactions influences the merger dynamics and the resulting gravitational-wave signals. This provides a first step toward understanding how binary black-hole observations may constrain broader regions of EMDA theory beyond the analytically known special cases.

The goal of this chapter is not to produce catalog-level simulations. Instead, we use approximate charged binary black-hole initial data to explore how the dynamics depend on the EMDA coupling parameters. The initial configurations are designed to be close to the corresponding charged black-hole solutions, after which the black holes and surrounding fields are allowed to relax dynamically during the evolution. We then use the subsequent inspiral and merger to search for differences in the dynamics and emitted radiation that may distinguish between different choices of the coupling parameters.

4.2 Theory space

As our starting point, we consider Einstein–Maxwell–dilaton–axion (EMDA) theory, in which the Einstein–Maxwell system is supplemented by a scalar dilaton field ϕ and a pseudoscalar axion field κ . The action is

$$S = \int d^4x, \sqrt{-g} \left[\frac{R}{8\pi} - 2\nabla_a \phi \nabla^a \phi - e^{-2\alpha_0 \phi} F_{ab} F^{ab} - \frac{1}{2} e^{4\alpha_1 \phi} \nabla_a \kappa \nabla^a \kappa - \kappa F_{ab} (*F)^{ab} \right], \quad (4.1)$$

where R is the Ricci scalar, F_{ab} is the $U(1)$ field strength, and $(*F)_{ab}$ is its dual. The constants α_0 and α_1 determine the coupling strengths between the dilaton, axion, and electromagnetic field. In this chapter, we focus primarily on the one-parameter diagonal sector

$$\alpha_0 = \alpha_1 \equiv \alpha. \quad (4.2)$$

Different choices of α_0 and α_1 correspond to different members of this class of theories. The choice $(\alpha_0, \alpha_1) = (1, 1)$ corresponds to the coupling values associated with the low-energy limit of heterotic string theory and admits the Kerr–Sen black-hole solution [74, 75]. Setting $\alpha_0 = 1, \alpha_1 = 0$ gives an Einstein–Maxwell–dilaton-type system, with the Kaluza–Klein value corresponding to $\alpha_0 = \sqrt{3}$ [105, 106]. The case $(\alpha_0, \alpha_1) = (0, 0)$ removes the exponential dilaton dependence from the Maxwell and axion kinetic terms and provides a useful reference point for comparison.

For the comparisons presented here, we consider the diagonal EMDA values

$$(\alpha_0, \alpha_1) = (0, 0), \quad (1, 1), \quad (10, 10), \quad (100, 100). \quad (4.3)$$

We also include selected EMD benchmark cases,

$$(\alpha_0, \alpha_1) = (1, 0), \quad (\sqrt{3}, 0), \quad (4.4)$$

to compare the diagonal EMDA evolutions with representative dilatonic cases. These choices allow us to compare a weakly coupled reference case, the standard EMDA/Kerr–Sen coupling, two larger-coupling EMDA cases, and known EMD limits.

Analytic stationary black-hole solutions are not known for all of the coupling choices considered here. Moreover, even when stationary single-black-hole solutions are known, our binary configurations are constructed using approximate initial data rather than exact binary black-hole solutions. We therefore construct approximate charged binary black-hole initial data and evolve the full EMDA system numerically. During the early part of the evolution, the black holes and surrounding fields are allowed to relax dynamically before inspiral and merger.

This approach allows us to compare the resulting gravitational waveforms against a common reference waveform and to identify how the electromagnetic, dilaton, and axion fields influence the inspiral and merger dynamics. The known analytic black-hole metrics and associated fields relevant to this theory space are summarized in Appendix B.6.

4.3 Numerical methods

To study the strong-field dynamics and gravitational radiation from inspiraling charged black-hole binaries in Einstein–Maxwell–dilaton–axion theory, we evolve the full EMDA system as a Cauchy initial-value problem. The spacetime metric is evolved using the BSSN formulation [84–86], together with puncture gauge conditions [87, 88]. The electromagnetic, dilaton, and axion fields are evolved using the matter evolution equations obtained from the $3+1$ decomposition of the EMDA equations of motion. The full evolution system is summarized in Appendix A.4.

We use the numerical code developed in Ref. [99]. In contrast to the head-on collisions considered there, the simulations presented here describe binary black holes with nonzero orbital angular momentum. The black holes are assigned initial linear momenta chosen to approximate quasicircular inspiral. This allows us to follow the late inspiral, merger, and ringdown and to extract gravitational, electromagnetic, dilaton, and axion radiation from the resulting spacetime.

Because the initial data are approximate, the electromagnetic constraints do not vanish identically at the initial time. To control these constraint violations, we evolve the electromagnetic fields with hyperbolic divergence cleaning. In this approach, auxiliary damping fields propagate violations of the electric and magnetic constraints away from the strong-field region and damp them during the evolution [124]. This prevents electromagnetic constraint violations from accumulating during the relaxation, inspiral, and merger.

4.3.1 Initial data

Constructing initial data for binary black holes in EMDA theory is challenging because exact binary black-hole solutions are not known. In addition, for general choices of the coupling parameters, the corresponding stationary single-black-hole solutions are not available in closed analytic form. We therefore use an approximate initial-data prescription guided by the known Kerr–Newman and Kerr–Sen solutions [74, 75].

Our starting point is a charged Kerr–Newman-like binary black-hole configuration constructed using approximate charged puncture data [93, 97, 125]. We supplement these data with approximate dilaton and axion fields motivated by the Kerr–Sen solution. Since we are interested in varying the coupling parameters, we rescale the scalar and pseudoscalar fields according to the chosen coupling. In particular, for the diagonal choice $\alpha_0 = \alpha_1 \equiv \alpha$, we take the initial dilaton and axion profiles to scale with α , so that larger coupling values correspond to stronger initial scalar and pseudoscalar field profiles.

This prescription does not produce an exact solution of the EMDA constraint equations. Instead, it should be viewed as an approximate-data construction designed to place the system near the expected charged black-hole configuration for each coupling choice. After the evolution begins, the black holes and surrounding fields are allowed to relax dynamically before the late inspiral and merger.

For each coupling choice, we consider binaries with charge-to-mass ratio magnitude $\lambda = Q/M = 0.01$ and a small spin $\chi = 0.2$. We examine three representative charge configurations: $++$, $+-$, and $+0$. In the $++$ case, both black holes carry charge with the same sign; in the $+-$ case, the black holes carry equal and opposite charges; and in the $+0$ case, only one black hole is charged. These configurations allow us to compare systems with different net charge and electromagnetic field structure while keeping the individual charge scale fixed.

We then use the subsequent evolution to study how changing the coupling strength affects the electromagnetic, dilaton, and axion fields, the binary dynamics, and the emitted gravitational radiation.

4.4 Results

4.4.1 Waveform comparison

We first compare the gravitational waveforms produced by the different coupling choices. For this comparison, we use the Einstein–Maxwell case, $(\alpha_0, \alpha_1) = (0, 0)$, as the reference waveform and compare the dominant $(\ell, m) = (2, 2)$ mode against the corresponding modes from the EMD and EMDA evolutions.

For the charge considered here, the gravitational waveforms remain close to the Einstein–Maxwell reference waveform across all of the coupling choices. In most cases, the differences in the dominant strain are too small to interpret as clearly resolved coupling-dependent effects. Because the simulations use approximate initial data, the early relaxation of the black holes and surrounding fields can introduce waveform differences that are unrelated to the intended change in the coupling parameters. These effects, together with finite-resolution error, waveform-extraction error, and the short duration of the evolutions, likely dominate the very small differences observed among most of the runs.

The main robust conclusion is therefore qualitative. For the smaller-coupling EMD and EMDA cases, the dominant gravitational waveform is effectively indistinguishable from the Einstein–Maxwell reference waveform at the accuracy of the present simulations. The strongest-coupling case, $(\alpha_0, \alpha_1) = (100, 100)$, produces the largest departure from the reference waveform and is the only case in this comparison for which the coupling-dependent difference is visibly apparent. Even in this case, however, the dominant strain remains close to the Einstein–Maxwell waveform.

These results suggest that, for the relatively small charge considered here, coupling-dependent effects in the dominant gravitational-wave mode are subtle. Larger charges, longer inspirals, improved initial data, and explicit convergence tests may be required to separate physical coupling-dependent effects from relaxation and numerical errors. The present comparisons should therefore be interpreted as an exploratory test of the EMDA coupling dependence rather than as precision estimates of waveform distinguishability.

Figure 4.1 shows a representative comparison between the Einstein–Maxwell reference waveform and the strongest-coupling EMDA case. The two waveforms are peak-aligned to make the small differences near merger visible.

4.4.2 Constraints

Because the initial data are approximate, the constraint violations provide an important check on the quality of the evolutions. We monitor the Hamiltonian constraint, together with the electric and magnetic divergence constraints, throughout the simulations. These quantities are shown in Figs. 4.2–4.4. Across the set of coupling choices considered here, the constraints remain controlled during the initial relaxation, inspiral, and merger. The largest features occur during the early relaxation phase and near merger, where the fields vary most rapidly.

The electromagnetic constraints remain bounded throughout the evolutions, indicating that the divergence-cleaning scheme prevents the electric and magnetic constraint violations from growing




Figure 4.1 Peak-aligned comparison of the dominant gravitational-wave strain between the Einstein–Maxwell reference waveform, $(\alpha_0, \alpha_1) = (0, 0)$, and the strongest-coupling EMDA case, $(\alpha_0, \alpha_1) = (100, 100)$. The figure shows the plus polarization of the strain.

without bound. The Hamiltonian constraint shows the expected transient behavior during the initial relaxation and near merger, but it does not exhibit runaway growth. These results indicate that the simulations remain sufficiently well controlled for the qualitative waveform comparisons presented above.

4.5 Conclusion

In this chapter, we studied binary black-hole mergers in a broader region of Einstein–Maxwell–dilaton–axion theory by varying the coupling parameters that control the interactions between the electromagnetic, dilaton, and axion fields. We focused primarily on the diagonal sector, $\alpha_0 = \alpha_1$, and compared several representative coupling choices against the Einstein–Maxwell reference case. Because exact binary black-hole initial data are not available for these theories, we used approximate charged binary black-hole initial data and allowed the systems to relax dynamically before merger.




Figure 4.2 Hamiltonian constraint violation for the different coupling choices. The curves show the evolution of the Hamiltonian constraint norm for the Einstein–Maxwell reference case and the corresponding EMD and EMDA evolutions.

The evolutions remained stable for the coupling choices considered here, and the electromagnetic and gravitational constraints remained controlled throughout the simulations. This provides evidence that the numerical framework can be used to explore binary black-hole dynamics beyond the analytically known special cases of EMDA theory. In particular, these simulations provide a first step toward studying how nonlinear merger dynamics depend on the theory’s coupling parameters.

For the small charge considered in this study, we found that the dominant gravitational-wave signal is only weakly affected by the choice of coupling. The smaller-coupling EMD and EMDA cases remained effectively indistinguishable from the Einstein–Maxwell reference waveform at the accuracy of the present simulations. The strongest-coupling case, $(\alpha_0, \alpha_1) = (100, 100)$, produced the largest departure from the reference waveform, but even this difference remained small in the dominant strain.




Figure 4.3 Electric divergence constraint violation for the different coupling choices. The curves show the evolution of the electric constraint norm across the Einstein–Maxwell, EMD, and EMDA evolutions.

These results suggest that coupling-dependent effects in the gravitational radiation are subtle at small charge. Because the simulations use approximate initial data, very small waveform differences may be dominated by relaxation effects, finite-resolution error, and waveform-extraction uncertainty rather than by physical coupling dependence. Larger charges, longer inspirals, improved initial data, and explicit convergence tests may be required to clearly separate physical effects from numerical and initial-data uncertainties.




Figure 4.4 Magnetic divergence constraint violation for the different coupling choices. The curves show the evolution of the magnetic constraint norm across the Einstein–Maxwell, EMD, and EMDA evolutions.

Chapter 5

CFD Chapter 1

5.1 CFD

Chapter 6

Conclusion

6.1 Conclusion

Throughout this dissertation, we have pursued two complementary themes in numerical relativity. The first is the numerical modeling of black-hole spacetimes in Einstein–Maxwell–dilaton–axion theory, an extension of general relativity that includes electromagnetic, dilaton, and axion fields. Within this setting, we studied the nonlinear stability of charged black holes and investigated how these additional fields can affect gravitational waveforms from binary black-hole mergers. This work provides an example of how numerical relativity can be used to explore strong-field dynamics in theories beyond general relativity.

The second theme is the improvement of numerical relativity methods through the use of compact finite-difference operators. These methods are designed to improve the accuracy and efficiency of simulations of strong-field gravitational systems. In this final chapter, we summarize the main conclusions from each part of the dissertation and discuss directions for future work.

In the first EMDA portion of this dissertation, we developed a method for constructing approximate binary black-hole initial data in Einstein–Maxwell–dilaton–axion theory. This allowed us to evolve head-on collisions of charged black holes across a range of initial spins, charges, and field configurations. These simulations were used to study whether the additional fields present in EMDA theory remain associated with black holes in the fully nonlinear merger regime.

We found that scalar hair persists through merger for initially scalarized Kerr–Sen black holes. When the black holes carry nontrivial electric charge, the dilaton field remains present after merger. When both charge and spin are present, the remnant also retains an axion field. The horizon quantities of the final black hole settle to approximately constant values, supporting the interpretation that the remnant approaches another Kerr–Sen black hole. We also showed that initially unscalarized black holes can dynamically develop scalar hair. In particular, initially Kerr–Newman-like black holes in EMDA evolve toward configurations with nontrivial scalar structure. These results suggest that scalar and axionic hair can be robust features of charged black holes in EMDA theory.

For inspiraling black-hole binaries, we found that increasing the charge produces small changes in the gravitational waveform relative to the corresponding neutral GR case. These waveform differences provide a first indication of how EMDA fields may affect binary black-hole signals in the strong-field regime. We also constructed a modest waveform catalog from the simulations performed in this dissertation, including several charge configurations, spins, and mass ratios. In particular, this catalog includes EMDA binary black-hole mergers with mass ratios as large as $q = 8$, demonstrating that these simulations can be extended beyond the equal-mass case.

In the broader EMDA theory-space study, we investigated how binary black-hole evolutions change as the coupling parameters are varied. We explored a range of EMDA coupling choices and compared these results with simulations in Einstein–Maxwell theory, Kaluza–Klein-inspired theory, and related Einstein–Maxwell–dilaton models. This allowed us to search for significant differences in the dynamics and to assess how theory-dependent effects appear in numerical evolutions. For the small charges considered in that study, the dominant gravitational-wave signal was only weakly affected by the coupling choice, with the largest differences appearing in the strongest-coupling cases.

In the compact finite-difference work, we developed novel compact derivative operators and evaluated their performance against comparable explicit finite-difference operators. In the tests considered here, the fourth-order compact operators achieved accuracy comparable to fourth-order explicit operators while using roughly half the resolution. We also demonstrated that sixth-order compact derivatives can outperform their explicit counterparts. These improvements were shown for both first- and second-derivative operators and were tested across three systems: Maxwell theory, a nonlinear sigma model, and the BSSN equations. The improved accuracy of these operators suggests that compact finite differences may provide a useful tool for improving the fidelity of numerical relativity simulations.

Several directions remain for future work. First, compact finite-difference operators should be implemented and tested in full binary black-hole evolutions. Preliminary work in this direction has already shown some advantages, but additional simulations are needed to determine whether these operators provide a robust improvement over standard explicit finite differences in realistic merger calculations. If these advantages persist, compact finite differences could provide a path toward faster and higher-fidelity gravitational-wave simulations, which will be increasingly important as next-generation detectors require more accurate waveform models.

Second, future EMDA simulations could be used to make closer contact with gravitational-wave observations. One promising approach would be to simulate systems similar to high-significance observed binary black-hole mergers whose masses and spins are relatively well constrained. By comparing EMDA waveforms against observed signals, one could place constraints on the possible dilaton and axion charges of the merging black holes. Such a study would help connect the theoretical and numerical results of this dissertation to observational tests of deviations from general relativity.

Taken together, the results of this dissertation demonstrate how numerical relativity can both expand the study of black-hole dynamics beyond general relativity and improve the computational tools needed to model gravitational-wave sources with increasing accuracy.

Appendix A

Mathematical Overview

A.1 Electricity and Magnetism

We give here a simple example of how a system of field equations can be separated into evolution equations and constraint equations. We use Maxwell's equations as the example, following an approach similar to that of *Numerical Relativity: Starting from Scratch* [20]. This example is presented in flat spacetime in order to emphasize the basic structure of the decomposition before introducing the additional geometric terms that appear in curved spacetime.

We begin with Maxwell's equations written in terms of the Faraday tensor,

$$\nabla_b F^{ab} = 4\pi j^a, \quad (\text{A.1})$$

together with the homogeneous equation

$$\nabla_{[a} F_{bc]} = 0. \quad (\text{A.2})$$

Here j^a is the four-current, which we write schematically as

$$j^a = (\rho, J^i), \quad (\text{A.3})$$

where ρ is the charge density and J^i is the spatial current. Charge conservation is expressed by the continuity equation

$$\nabla_a j^a = 0. \quad (\text{A.4})$$

In general relativity, an analogous conservation law for the stress-energy tensor follows from the Bianchi identity.

The Faraday tensor can be written in terms of a vector potential A_a as

$$F_{ab} = \nabla_a A_b - \nabla_b A_a. \quad (\text{A.5})$$

With this definition, the homogeneous Maxwell equation $\nabla_{[a} F_{bc]} = 0$ is satisfied automatically. In flat spacetime, we decompose the vector potential into an electromagnetic scalar potential Φ and a spatial vector potential A_i . With metric signature $(-, +, +, +)$, we may write schematically

$$A_a = (-\Phi, A_i). \quad (\text{A.6})$$

In flat spacetime, the electric field can be written in terms of the potentials as

$$E_i = -\partial_t A_i - D_i \Phi, \quad (\text{A.7})$$

where D_i denotes the spatial derivative. Taking the divergence gives

$$D_i E^i = -\partial_t D_i A^i - D_i D^i \Phi. \quad (\text{A.8})$$

The time component of Eq. (A.1) gives

$$D_i E^i = 4\pi\rho, \quad (\text{A.9})$$

which is Gauss's Law and is a constraint equation. Equivalently, Eq. (A.7) may be rearranged to give

$$\partial_t A_i = -E_i - D_i \Phi. \quad (\text{A.10})$$

This equation gives the time evolution of the spatial vector potential once the scalar potential, or equivalently a gauge condition, has been chosen.

We now consider the spatial components. The magnetic field is defined in terms of the vector potential by

$$B^i = \epsilon^{ijk} D_j A_k, \quad (\text{A.11})$$

so that the magnetic constraint

$$D_i B^i = 0 \quad (\text{A.12})$$

is automatically satisfied in flat space. The spatial components of Maxwell's equations give the evolution equation for the electric field,

$$\partial_t E_i = -D_j D^j A_i + D_i D^j A_j - 4\pi J_i. \quad (\text{A.13})$$

This equation is equivalent to the usual curl form of Ampere's law,

$$\partial_t \mathbf{E} = \nabla \times \mathbf{B} - 4\pi \mathbf{J}. \quad (\text{A.14})$$

Thus, in this simple flat-space example, Maxwell's equations can be separated into evolution equations,

$$\partial_t A_i = -E_i - D_i \Phi, \quad (\text{A.15})$$

$$\partial_t E_i = -D_j D^j A_i + D_i D^j A_j - 4\pi J_i, \quad (\text{A.16})$$

and constraint equations,

$$D_i E^i = 4\pi \rho, \quad (\text{A.17})$$

$$D_i B^i = 0. \quad (\text{A.18})$$

This illustrates the same basic structure that appears in the 3 + 1 decomposition of general relativity: part of the system provides evolution equations, while the remaining equations constrain the allowed initial data. In curved spacetime, the same decomposition must be performed with the spacetime metric, lapse, shift, spatial metric, and spatial covariant derivative included explicitly.

A.2 Spacetime and Kinematics

We now allow for more general slicings of spacetime. This freedom is needed to formulate the Cauchy problem in general relativity and in alternative theories of gravity. We foliate the spacetime into a family of spacelike hypersurfaces Σ_t , which may be interpreted as slices of constant coordinate time t .

The gradient of the time function is

$$\nabla_a t = \partial_a t. \quad (\text{A.19})$$

This covector is normal to the constant-time hypersurfaces. We define the future-directed unit normal n_a to the slices by

$$n_a = -\alpha \nabla_a t, \quad (\text{A.20})$$

where α is the lapse function. The normalization condition is

$$n_a n^a = g^{ab} n_a n_b = -1. \quad (\text{A.21})$$

Because the hypersurfaces are spacelike, their normal is timelike, and we therefore normalize it to negative unity. The lapse α measures the rate at which proper time advances relative to coordinate time for observers moving normal to the slices.

In coordinates adapted to the foliation,

$$\nabla_a t = \partial_a t = \frac{\partial t}{\partial x^a}, \quad (\text{A.22})$$

so the only nonzero component is the time component. Therefore,

$$n_a = (-\alpha, 0, 0, 0). \quad (\text{A.23})$$

For an observer moving normal to the slice, the proper time interval satisfies

$$d\tau = \alpha dt. \quad (\text{A.24})$$

The coordinate time-evolution vector t^a can be decomposed into a part normal to the slice and a part tangent to the slice:

$$t^a = \alpha n^a + \beta^a. \quad (\text{A.25})$$

Here β^a is the shift vector. It is purely spatial, so

$$n_a \beta^a = 0. \quad (\text{A.26})$$

Equivalently,

$$\alpha n^a = t^a - \beta^a. \quad (\text{A.27})$$

In coordinates adapted to the foliation, $\beta^a = (0, \beta^i)$ and $t^a = (1, 0, 0, 0)$. Therefore,

$$n^a = \frac{1}{\alpha} (1, -\beta^i). \quad (\text{A.28})$$

The lapse controls the separation between neighboring slices in proper time, while the shift describes how the spatial coordinates move from one slice to the next.

With these definitions, the spacetime line element can be written in ADM form as

$$ds^2 = -\alpha^2 dt^2 + \gamma_{ij} (dx^i + \beta^i dt) (dx^j + \beta^j dt). \quad (\text{A.29})$$

This form makes explicit the split between the time direction, described by α and β^i , and the spatial geometry, described by the spatial metric γ_{ij} .

The spacetime metric can be decomposed into spatial and temporal parts using the normal vector. The spatial metric is

$$\gamma_{ab} = g_{ab} + n_a n_b, \quad (\text{A.30})$$

and the corresponding projection operator is

$$\gamma^a_b = \delta^a_b + n^a n_b. \quad (\text{A.31})$$

This operator projects spacetime tensors onto the spatial hypersurfaces.

The way the spatial hypersurfaces are embedded in spacetime is described by the extrinsic curvature. With the sign convention used in this work, we define

$$K_{ab} = -\gamma_a^c \gamma_b^d \nabla_c n_d. \quad (\text{A.32})$$

Equivalently, K_{ab} measures the spatially projected change of the normal vector from point to point on the hypersurface.

The Lie derivative measures how a tensor field changes as it is dragged along the flow generated by a vector field. For a covector A_d , the Lie derivative along n^a is

$$\mathcal{L}_n A_d = n^c \nabla_c A_d + A_c \nabla_d n^c. \quad (\text{A.33})$$

In the 3 + 1 decomposition, the relation

$$t^a = \alpha n^a + \beta^a \quad (\text{A.34})$$

allows coordinate time evolution to be separated into evolution normal to the slice and spatial transport along the shift.

Before moving on, we also recall the definition of the covariant derivative and the Christoffel symbols. For scalars, the ordinary partial derivative is sufficient. For tensors, however, the derivative must also account for changes in the basis vectors. This is done by introducing the connection coefficients, or Christoffel symbols.

For a vector A^b , the covariant derivative is defined by

$$\nabla_a A^b = \partial_a A^b + \Gamma^b_{ac} A^c. \quad (\text{A.35})$$

For a covector A_b , the corresponding expression is

$$\nabla_a A_b = \partial_a A_b - \Gamma^c_{ab} A_c. \quad (\text{A.36})$$

For the Levi-Civita connection, which is metric compatible and torsion-free, the Christoffel symbols are

$$\Gamma^a_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{dc} + \partial_c g_{db} - \partial_d g_{bc}). \quad (\text{A.37})$$

A.3 General Relativity

To decompose Einstein's equations in a similar way, we begin with the spacetime metric g_{ab} and Einstein's equation,

$$G_{ab} + \Lambda g_{ab} = 8\pi T_{ab}. \quad (\text{A.38})$$

For the remainder of this discussion we set $\Lambda = 0$, as is appropriate for the asymptotically flat spacetimes considered in this work.

At first, Einstein's equation appears to provide ten equations for the ten independent components of the spacetime metric. However, the Bianchi identity,

$$\nabla_a G^{ab} = 0, \quad (\text{A.39})$$

must always be satisfied. This identity plays a role analogous to constraint preservation in electromagnetism. It implies that four of Einstein's equations are not independent evolution equations. We therefore expect four constraint equations in general relativity, along with a fourfold gauge freedom associated with the choice of spacetime coordinates.

We now decompose the spacetime metric into spatial and temporal parts. The spatial metric is defined by

$$\gamma_{ab} = g_{ab} + n_a n_b, \quad (\text{A.40})$$

where n^a is the future-directed unit normal to the spatial hypersurfaces. Equivalently,

$$g_{ab} = \gamma_{ab} - n_a n_b. \quad (\text{A.41})$$

The tensor γ_{ab} acts as the metric intrinsic to each spatial slice. Since it is purely spatial, its contraction with the normal vanishes:

$$\gamma_{ab} n^b = 0. \quad (\text{A.42})$$

The inverse spacetime metric can be written as

$$g^{ab} = \gamma^{ab} - n^a n^b. \quad (\text{A.43})$$

In coordinates adapted to the foliation, this becomes

$$g^{ab} = \begin{pmatrix} -\alpha^{-2} & \alpha^{-2} \beta^i \\ \alpha^{-2} \beta^j & \gamma^{ij} - \alpha^{-2} \beta^i \beta^j \end{pmatrix}. \quad (\text{A.44})$$

The line element is then

$$ds^2 = -\alpha^2 dt^2 + \gamma_{ij} (dx^i + \beta^i dt) (dx^j + \beta^j dt). \quad (\text{A.45})$$

Here α is the lapse function and β^i is the shift vector.

The extrinsic curvature can be related to the time evolution of the spatial metric. With the sign convention adopted in this work,

$$K_{ab} = -\frac{1}{2} \mathcal{L}_n \gamma_{ab}. \quad (\text{A.46})$$

Using the decomposition $t^a = \alpha n^a + \beta^a$, this becomes

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij}. \quad (\text{A.47})$$

Thus, the extrinsic curvature measures how the spatial metric changes in time, up to the lapse and shift. The spatial metric γ_{ij} and the extrinsic curvature K_{ij} form the basic dynamical variables in the ADM decomposition.

We next introduce the spatial covariant derivative D_i , which is the derivative operator compatible with the spatial metric γ_{ij} . For a spatial vector V^j ,

$$D_i V^j = \partial_i V^j + \Gamma_{ik}^j V^k, \quad (\text{A.48})$$

where Γ_{jk}^i is the Christoffel symbol associated with γ_{ij} :

$$\Gamma_{jk}^i = \frac{1}{2} \gamma^{il} (\partial_j \gamma_{lk} + \partial_k \gamma_{lj} - \partial_l \gamma_{jk}). \quad (\text{A.49})$$

The spatial projections of the spacetime curvature are related to the intrinsic curvature of the spatial slice and the extrinsic curvature through the Gauss–Codazzi relations. These relations provide the geometric bridge between the four-dimensional spacetime curvature and the three-dimensional geometry of each spatial slice.

The Gauss equation is

$$\gamma^a_i \gamma^b_j \gamma^c_k \gamma^d_l {}^{(4)}R_{abcd} = {}^{(3)}R_{ijkl} + K_{ik} K_{jl} - K_{il} K_{jk}. \quad (\text{A.50})$$

Here ${}^{(4)}R_{abcd}$ is the four-dimensional spacetime Riemann tensor, while ${}^{(3)}R_{ijkl}$ is the three-dimensional Riemann tensor associated with the spatial metric γ_{ij} .

The Codazzi equation is

$$\gamma^a_i \gamma^b_j \gamma^c_k n^d {}^{(4)}R_{abcd} = D_j K_{ik} - D_i K_{jk}. \quad (\text{A.51})$$

This relates the mixed spatial-normal projection of the spacetime curvature to spatial derivatives of the extrinsic curvature.

The Ricci equation, which involves two projections along the normal direction, is

$$\gamma^a{}_i \gamma^b{}_j n^c n^{d(4)} R_{acbd} = \mathcal{L}_n K_{ij} + \frac{1}{\alpha} D_i D_j \alpha + K_{ik} K^k{}_j. \quad (\text{A.52})$$

To obtain the ADM equations, we insert these decompositions into Einstein's equation and equate the projections of the Einstein tensor with the corresponding projections of the stress-energy tensor in Eq. (A.130).

Projecting Einstein's equation twice along the normal direction gives the Hamiltonian constraint:

$$R + K^2 - K_{ij} K^{ij} = 16\pi\rho. \quad (\text{A.53})$$

Here

$$\rho = n^a n^b T_{ab} \quad (\text{A.54})$$

is the energy density measured by an observer moving normal to the spatial slice. The quantity K is the trace of the extrinsic curvature,

$$K = \gamma^{ij} K_{ij}. \quad (\text{A.55})$$

A mixed projection, with one index projected along the normal direction and one projected onto the spatial slice, gives the momentum constraint:

$$D_j (K^{ij} - \gamma^{ij} K) = 8\pi j^i. \quad (\text{A.56})$$

Here

$$j^i = -\gamma^{ia} n^b T_{ab} \quad (\text{A.57})$$

is the momentum density measured by the normal observer.

A complete spatial projection gives the evolution equation for the extrinsic curvature:

$$\partial_t K_{ij} = \alpha \left(R_{ij} - 2K_{ik} K^k{}_j + K K_{ij} \right) - D_i D_j \alpha - 4\pi\alpha M_{ij} + \mathcal{L}_\beta K_{ij}. \quad (\text{A.58})$$

In this expression,

$$M_{ij} = 2S_{ij} - \gamma_{ij} (S - \rho), \quad (\text{A.59})$$

where

$$S_{ij} = \gamma^a{}_i \gamma^b{}_j T_{ab} \quad (\text{A.60})$$

is the spatial projection of the stress-energy tensor, and

$$S = \gamma^{ij} S_{ij} \quad (\text{A.61})$$

is its trace.

Together with Eq. (A.47), these equations form the ADM system. The variables γ_{ij} and K_{ij} are the dynamical variables, while α and β^i encode the gauge freedom associated with the choice of spacetime coordinates.

$R + K^2 - K_{ij}K^{ij} = 16\pi\rho,$	Hamiltonian constraint,
$D_j (K^{ij} - \gamma^{ij} K) = 8\pi j^i,$	Momentum constraint,
$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij},$	Metric evolution,
$\partial_t K_{ij} = \alpha \left(R_{ij} - 2K_{ik}K^k{}_j + K K_{ij} \right) - D_i D_j \alpha$ $- 4\pi\alpha \left[2S_{ij} - \gamma_{ij}(S - \rho) \right] + \mathcal{L}_\beta K_{ij},$	Extrinsic-curvature evolution.

(A.62)

A.3.1 Reformulating the Einstein Equations

The ADM equations provide a natural 3 + 1 decomposition of Einstein's equations, but the standard ADM system is only weakly hyperbolic and is not generally well suited for stable numerical evolution. It is therefore useful to reformulate the equations into a system with better hyperbolic properties. The Baumgarte–Shapiro–Shibata–Nakamura (BSSN) formulation achieves this by introducing conformal variables and promoting certain combinations of spatial derivatives of the metric to independent evolved quantities [84–86].

A Maxwell Analogy

We begin with an analogy from electromagnetism. In a flat-space 3 + 1 decomposition, Maxwell's equations can be written in terms of the electric field E_i and vector potential A_i as

$$\partial_t E_i = -D_j D^j A_i + D_i D^j A_j - 4\pi J_i, \quad (\text{A.63})$$

together with

$$\partial_t A_i = -E_i - D_i \Phi. \quad (\text{A.64})$$

Here Φ is the electromagnetic scalar potential, and J_i is the spatial current. These evolution equations are supplemented by Gauss's law,

$$D_i E^i = 4\pi \rho. \quad (\text{A.65})$$

We define the electric constraint violation by

$$C \equiv D_i E^i - 4\pi \rho. \quad (\text{A.66})$$

For constraint-satisfying data,

$$C = 0. \quad (\text{A.67})$$

In exact arithmetic, if the constraint is satisfied initially, it remains satisfied during the evolution. In a numerical evolution, however, truncation error and roundoff error generally produce small but nonzero values of C . It is therefore useful to reformulate the evolution system so that constraint violations propagate in a controlled way.

We now introduce the auxiliary variable

$$\Gamma \equiv D_i A^i. \quad (\text{A.68})$$

This definition introduces an additional auxiliary constraint,

$$\mathcal{G} \equiv \Gamma - D_i A^i. \quad (\text{A.69})$$

For the original Maxwell system, $\mathcal{G} = 0$. We can nevertheless promote Γ to an independent evolved variable. Taking a time derivative of its defining relation gives

$$\partial_t \Gamma = D_i \partial_t A^i. \quad (\text{A.70})$$

Using the evolution equation for A_i , we find

$$\partial_t \Gamma = -D_i E^i - D_i D^i \Phi. \quad (\text{A.71})$$

Since $C = 0$ for constraint-satisfying data, we may add an arbitrary multiple of C to the right-hand side without changing the continuum solutions. Thus, we write

$$\partial_t \Gamma = -D_i E^i - D_i D^i \Phi + \eta C, \quad (\text{A.72})$$

where η is a constant. Using

$$C = D_i E^i - 4\pi\rho, \quad (\text{A.73})$$

this can also be written as

$$\partial_t \Gamma = (\eta - 1) D_i E^i - D_i D^i \Phi - 4\pi\eta\rho. \quad (\text{A.74})$$

With this new variable, the Maxwell evolution system becomes

$$\partial_t A_i = -E_i - D_i \Phi, \quad (\text{A.75})$$

$$\partial_t E_i = -D^2 A_i + D_i \Gamma - 4\pi J_i, \quad (\text{A.76})$$

$$\partial_t \Gamma = -D_i E^i - D_i D^i \Phi + \eta C. \quad (\text{A.77})$$

Here $D^2 \equiv D_i D^i$. This system is equivalent to the original Maxwell system when the constraints are satisfied, but it changes how constraint violations behave.

Starting from

$$C = D_i E^i - 4\pi\rho, \quad (\text{A.78})$$

we take a time derivative:

$$\partial_t C = D_i \partial_t E^i - 4\pi \partial_t \rho. \quad (\text{A.79})$$

Using charge conservation,

$$\partial_t \rho = -D_i J^i, \quad (\text{A.80})$$

we obtain

$$\partial_t C = D_i \partial_t E^i + 4\pi D_i J^i. \quad (\text{A.81})$$

Substituting the evolution equation for E_i gives

$$\partial_t C = D^i \left(-D^2 A_i + D_i \Gamma - 4\pi J_i \right) + 4\pi D_i J^i \quad (\text{A.82})$$

$$= -D^i D^2 A_i + D^i D_i \Gamma. \quad (\text{A.83})$$

Assuming that the spatial derivatives commute, this becomes

$$\partial_t C = -D^2 D^i A_i + D^2 \Gamma. \quad (\text{A.84})$$

Therefore,

$$\partial_t C = D^2 (\Gamma - D_i A^i) = D^2 \mathcal{G}. \quad (\text{A.85})$$

We next compute the evolution of $\mathcal{G}$. Taking a time derivative gives

$$\partial_t \mathcal{G} = \partial_t \Gamma - D_i \partial_t A^i. \quad (\text{A.86})$$

Using the evolution equations for Γ and A_i , we find

$$\partial_t \mathcal{G} = (-D_i E^i - D_i D^i \Phi + \eta C) - (-D_i E^i - D_i D^i \Phi) \quad (\text{A.87})$$

$$= \eta C. \quad (\text{A.88})$$

Combining this equation with

$$\partial_t C = D^2 \mathcal{G} \quad (\text{A.89})$$

gives

$$\partial_t^2 C = D^2 \partial_t \mathcal{G} = \eta D^2 C. \quad (\text{A.90})$$

Equivalently,

$$\boxed{\left(-\partial_t^2 + \eta D_i D^i\right) C = 0}. \quad (\text{A.91})$$

Thus, for $\eta > 0$, electric constraint violations propagate with speed $\sqrt{\eta}$. Analytically, this reformulation does not change the physical solutions of Maxwell's equations, because the added terms vanish for constraint-satisfying data. Numerically, however, it changes the behavior of constraint violations by allowing them to propagate rather than remain fixed in place.

The BSSN formulation uses a related idea. It introduces auxiliary variables, most notably the conformal connection functions, and adds suitable multiples of the constraints to improve the hyperbolic structure and numerical behavior of the Einstein evolution system.

BSSN Equations

We begin with the ADM equations:

$$R + K^2 - K_{ij} K^{ij} = 16\pi\rho, \quad \text{Hamiltonian constraint,} \quad (\text{A.92})$$

$$D_j (K^{ij} - \gamma^{ij} K) = 8\pi j^i, \quad \text{Momentum constraint,} \quad (\text{A.93})$$

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + \mathcal{L}_\beta \gamma_{ij}, \quad \text{Metric evolution,} \quad (\text{A.94})$$

$$\begin{aligned} \partial_t K_{ij} = & \alpha \left(R_{ij} - 2K_{ik} K^k_j + K K_{ij} \right) - D_i D_j \alpha \\ & - 4\pi\alpha \left[2S_{ij} - \gamma_{ij}(S - \rho) \right] + \mathcal{L}_\beta K_{ij}, \quad \text{Extrinsic-curvature evolution.} \end{aligned} \quad (\text{A.95})$$

The Ricci tensor can be written in several different forms. In particular, R_{ij} contains second derivatives of the metric:

$$\begin{aligned} R_{ij} = & -\frac{1}{2} \gamma^{kl} (\partial_k \partial_l \gamma_{ij} + \partial_i \partial_j \gamma_{kl} - \partial_i \partial_l \gamma_{kj} - \partial_k \partial_j \gamma_{il}) \\ & + \gamma^{kl} (\Gamma^m_{il} \Gamma_{mkj} - \Gamma^m_{ij} \Gamma_{mkl}). \end{aligned} \quad (\text{A.96})$$

The first second-derivative term, $\gamma^{kl} \partial_k \partial_l \gamma_{ij}$, acts like a Laplace operator on the metric. However, because of the remaining second-derivative terms, the system is not immediately in a simple

wave-equation form. In analogy with the Maxwell example above, we introduce an auxiliary variable related to derivatives of the metric. This allows us to rewrite the Ricci tensor in terms of the trace of the Christoffel symbols. We define

$$\Gamma^i \equiv \gamma^{jk} \Gamma^i_{jk}. \quad (\text{A.97})$$

In terms of this contracted Christoffel symbol, the Ricci tensor can be written as

$$\begin{aligned} R_{ij} = & -\frac{1}{2} \gamma^{kl} \partial_k \partial_l \gamma_{ij} + \gamma_{k(i} \partial_{j)} \Gamma^k + \Gamma^k \Gamma_{(ij)k} \\ & + 2\gamma^{lm} \Gamma^k_{m(i} \Gamma_{j)kl} + \gamma^{kl} \Gamma^m_{il} \Gamma_{mkj}. \end{aligned} \quad (\text{A.98})$$

We can also use the identity relating the contracted Christoffel symbols to the divergence of the inverse metric:

$$g^{bc} \Gamma^a_{bc} = -\frac{1}{\sqrt{|g|}} \partial_b \left(\sqrt{|g|} g^{ab} \right). \quad (\text{A.99})$$

Applying this identity to the spatial metric gives

$$\Gamma^i = -\frac{1}{\sqrt{\gamma}} \partial_j \left(\sqrt{\gamma} \gamma^{ij} \right). \quad (\text{A.100})$$

In the BSSN formulation, the corresponding conformal connection functions are treated as independent evolved variables. To define them, we introduce a conformal decomposition of the spatial metric:

$$\gamma_{ij} = e^{4\varphi} \tilde{\gamma}_{ij}. \quad (\text{A.101})$$

The conformal factor is chosen so that

$$\tilde{\gamma} = 1, \quad (\text{A.102})$$

where $\tilde{\gamma}$ is the determinant of $\tilde{\gamma}_{ij}$. Taking the determinant of the conformal decomposition then gives

$$\varphi = \frac{1}{12} \ln \gamma, \quad (\text{A.103})$$

where γ is the determinant of the physical spatial metric γ_{ij} .

The physical Ricci tensor can be split into two pieces,

$$R_{ij} = \tilde{R}_{ij} + R_{ij}^{\varphi}, \quad (\text{A.104})$$

where $\tilde{R}_{ij}$ is computed from the conformally related metric $\tilde{\gamma}_{ij}$, and R_{ij}^{φ} contains terms involving the conformal factor φ .

The conformal Ricci tensor is

$$\begin{aligned} \tilde{R}_{ij} = & -\frac{1}{2} \tilde{\gamma}^{kl} \partial_k \partial_l \tilde{\gamma}_{ij} + \tilde{\gamma}_{k(i} \partial_{j)} \tilde{\Gamma}^k + \tilde{\Gamma}^k \tilde{\Gamma}_{(ij)k} \\ & + 2 \tilde{\gamma}^{lm} \tilde{\Gamma}_{m(i} \tilde{\Gamma}_{j)kl} + \tilde{\gamma}^{kl} \tilde{\Gamma}^m_{il} \tilde{\Gamma}_{mkj}. \end{aligned} \quad (\text{A.105})$$

The conformal-factor contribution is

$$\begin{aligned} R_{ij}^{\varphi} = & -2 \left(\tilde{D}_i \tilde{D}_j \varphi + \tilde{\gamma}_{ij} \tilde{\gamma}^{lm} \tilde{D}_l \tilde{D}_m \varphi \right) \\ & + 4 \left[(\tilde{D}_i \varphi) (\tilde{D}_j \varphi) - \tilde{\gamma}_{ij} \tilde{\gamma}^{lm} (\tilde{D}_l \varphi) (\tilde{D}_m \varphi) \right]. \end{aligned} \quad (\text{A.106})$$

The contracted conformal Christoffel symbols are defined by

$$\tilde{\Gamma}^i \equiv \tilde{\gamma}^{jk} \tilde{\Gamma}_{jk}^i = -\frac{1}{\tilde{\gamma}^{1/2}} \partial_j \left(\tilde{\gamma}^{1/2} \tilde{\gamma}^{ij} \right). \quad (\text{A.107})$$

With $\tilde{\gamma} = 1$, this reduces to

$$\tilde{\Gamma}^i = -\partial_j \tilde{\gamma}^{ij}. \quad (\text{A.108})$$

The extrinsic curvature is decomposed into its trace and trace-free parts:

$$K_{ij} = A_{ij} + \frac{1}{3} \gamma_{ij} K, \quad (\text{A.109})$$

where

$$K = \gamma^{ij} K_{ij}. \quad (\text{A.110})$$

The trace-free part is then conformally rescaled according to

$$A_{ij} = e^{4\varphi} \tilde{A}_{ij}, \quad (\text{A.111})$$

or equivalently,

$$\tilde{A}_{ij} = e^{-4\varphi} \left(K_{ij} - \frac{1}{3} \gamma_{ij} K \right). \quad (\text{A.112})$$

The evolved BSSN variables are therefore

$$\varphi, \quad \tilde{\gamma}_{ij}, \quad \tilde{A}_{ij}, \quad K, \quad \tilde{\Gamma}^i. \quad (\text{A.113})$$

These variables, together with appropriate gauge conditions for the lapse and shift, form the standard BSSN formulation.

The BSSN equations are

$\partial_t \varphi = -\frac{1}{6} \alpha K + \beta^i \partial_i \varphi + \frac{1}{6} \partial_i \beta^i,$	Conformal exponent evolution,
$\begin{aligned} \partial_t K = & -\gamma^{ij} D_i D_j \alpha + \alpha \left(\tilde{A}_{ij} \tilde{A}^{ij} + \frac{1}{3} K^2 \right) + 4\pi \alpha (\rho + S) \\ & + \beta^i \partial_i K, \end{aligned}$	Mean-curvature evolution,
$\begin{aligned} \partial_t \tilde{\gamma}_{ij} = & -2\alpha \tilde{A}_{ij} + \beta^k \partial_k \tilde{\gamma}_{ij} + \tilde{\gamma}_{ik} \partial_j \beta^k \\ & + \tilde{\gamma}_{kj} \partial_i \beta^k - \frac{2}{3} \tilde{\gamma}_{ij} \partial_k \beta^k, \end{aligned}$	Conformal-metric evolution,
$\begin{aligned} \partial_t \tilde{A}_{ij} = & e^{-4\varphi} \left[- (D_i D_j \alpha)^{\text{TF}} + \alpha \left(R_{ij}^{\text{TF}} - 8\pi S_{ij}^{\text{TF}} \right) \right] \\ & + \alpha \left(K \tilde{A}_{ij} - 2 \tilde{A}_{il} \tilde{A}^l{}_j \right) \\ & + \beta^k \partial_k \tilde{A}_{ij} + \tilde{A}_{ik} \partial_j \beta^k + \tilde{A}_{kj} \partial_i \beta^k - \frac{2}{3} \tilde{A}_{ij} \partial_k \beta^k, \end{aligned}$	Trace-free curvature evolution,
$\begin{aligned} \partial_t \tilde{\Gamma}^i = & -2 \tilde{A}^{ij} \partial_j \alpha + 2\alpha \left(\tilde{\Gamma}^i{}_{jk} \tilde{A}^{kj} - \frac{2}{3} \tilde{\gamma}^{ij} \partial_j K - 8\pi \tilde{\gamma}^{ij} j_j + 6 \tilde{A}^{ij} \partial_j \varphi \right) \\ & + \beta^j \partial_j \tilde{\Gamma}^i - \tilde{\Gamma}^j \partial_j \beta^i + \frac{2}{3} \tilde{\Gamma}^i \partial_j \beta^j \\ & + \frac{1}{3} \tilde{\gamma}^{li} \partial_l \partial_j \beta^j + \tilde{\gamma}^{lj} \partial_j \partial_l \beta^i, \end{aligned}$	Conformal-connection evolution.

(A.114)

Gauge and Slicing Conditions

The BSSN equations do not fully determine the evolution until the coordinate gauge is specified. In particular, one must choose evolution conditions for the lapse α and the shift β^i . These choices determine how the spatial slices are threaded through spacetime and how the spatial coordinates move from one slice to the next. We summarize several common choices below.

Geodesic slicing. A simple choice is

$$\alpha = 1, \quad \beta^i = 0. \quad (\text{A.115})$$

This is called geodesic slicing. Although this gauge is simple, it is usually a poor choice for numerical relativity simulations of black holes. The acceleration of the normal observers is

$$a_a = D_a \ln \alpha. \quad (\text{A.116})$$

For geodesic slicing, $\alpha = 1$, so $a_a = 0$. Thus, the normal observers follow geodesics. In a strong gravitational field, these freely falling observers can focus and form coordinate singularities, making geodesic slicing unsuitable for most black-hole evolutions.

Maximal slicing. Another possibility is to control the convergence of the normal observers by controlling the mean curvature. In maximal slicing, one imposes

$$K = 0. \quad (\text{A.117})$$

If this condition is maintained throughout the evolution, then the slicing condition implies

$$\partial_t K = 0, \quad (\text{A.118})$$

up to the corresponding shift terms. Maximal slicing has several desirable properties, including singularity avoidance. In particular, the lapse tends to collapse toward zero near strong-field regions,

causing those parts of the slice to advance more slowly in coordinate time. This behavior is known as collapse of the lapse. However, maximal slicing also has disadvantages, such as grid stretching, which can produce sharp coordinate gradients that are difficult to resolve numerically.

Harmonic coordinates. Harmonic coordinates are defined by requiring the spacetime coordinates to satisfy wave equations. Equivalently, one may impose

$$\Gamma^a = g^{bc}\Gamma^a_{bc} = 0. \quad (\text{A.119})$$

A related condition is harmonic slicing, in which only the time component of this condition is imposed. If we also set $\beta^i = 0$, then

$$\Gamma^0 = -\frac{1}{\alpha\sqrt{\gamma}}\partial_t\left(\frac{\sqrt{\gamma}}{\alpha}\right) = 0. \quad (\text{A.120})$$

This gives the lapse evolution equation

$$\partial_t\alpha = -\alpha^2 K. \quad (\text{A.121})$$

Harmonic slicing is closely related to wave coordinates and often leads to well-behaved hyperbolic evolution systems. However, modified harmonic slicing conditions are typically more useful for black-hole evolutions.

1 + log slicing and the Gamma-driver condition. A common generalization of harmonic slicing is the Bona–Masso family of slicing conditions,

$$\partial_t\alpha = -\alpha^2 f(\alpha)K. \quad (\text{A.122})$$

For $f(\alpha) = 1$, this reduces to harmonic slicing. For $f(\alpha) = 0$, the lapse is constant in time, giving geodesic slicing when $\alpha = 1$. In the formal limit $f(\alpha) \rightarrow \infty$, this condition approaches maximal slicing.

A particularly useful choice is

$$f(\alpha) = \frac{2}{\alpha}. \quad (\text{A.123})$$

This gives

$$\partial_t \alpha = -2\alpha K, \quad (\text{A.124})$$

which is the standard 1 + log slicing condition. This choice has singularity-avoidance properties similar to maximal slicing while retaining good behavior in nearly flat regions.

When the shift is nonzero, it is common to include an advective term and write

$$(\partial_t - \beta^i \partial_i) \alpha = -\alpha^2 f(\alpha) K. \quad (\text{A.125})$$

For $f(\alpha) = 2/\alpha$, this becomes

$$(\partial_t - \beta^i \partial_i) \alpha = -2\alpha K. \quad (\text{A.126})$$

We also need an evolution condition for the shift. A common choice is the Gamma-driver condition,

$$(\partial_t - \beta^j \partial_j) \beta^i = \mu B^i, \quad (\text{A.127})$$

together with

$$(\partial_t - \beta^j \partial_j) B^i = (\partial_t - \beta^j \partial_j) \tilde{\Gamma}^i - \eta B^i. \quad (\text{A.128})$$

Here B^i is an auxiliary variable, while μ and η are parameters that control the characteristic speed and damping of the shift condition. The combination of 1 + log slicing and the Gamma-driver shift condition is called the moving-puncture gauge. A major advantage of this gauge is that it allows black-hole spacetimes to be evolved without excising the singularity from the computational domain.

A.4 EMDA Equations of Motion

The action for EMDA in the Einstein frame is given as:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{8\pi} - 2(\nabla\phi)^2 - e^{-2\alpha_0\phi} F^2 - \frac{1}{2} e^{4\alpha_1\phi} (\nabla\kappa)^2 - \kappa F_{ab} (*F)^{ab} \right]. \quad (\text{A.129})$$

where ϕ and κ are the dilaton and axion, respectively. The tensor F_{ab} is the U(1) field strength, and $(*F)_{ab}$ is its dual. In this work we set the coupling constants to $\alpha_0 = \alpha_1 = 1$, corresponding to a low-energy limit of heterotic string theory.

The Einstein equations are:

$$G_{ab} = 8\pi G \left[2 \left(\nabla_a \phi \nabla_b \phi - \frac{1}{2} g_{ab} (\nabla \phi)^2 \right) + 2e^{-2\alpha_0 \phi} \left(F_{ac} F_b{}^c - \frac{1}{4} g_{ab} F^2 \right) + \frac{1}{2} e^{4\alpha_1 \phi} \left(\nabla_a \kappa \nabla_b \kappa - \frac{1}{2} g_{ab} (\nabla \kappa)^2 \right) \right]. \quad (\text{A.130})$$

while the matter equations can be written

$$0 = \nabla^a \nabla_a \phi + \frac{1}{2} \alpha_0 e^{-2\alpha_0 \phi} F^2 - \frac{1}{2} \alpha_1 e^{4\alpha_1 \phi} (\nabla \kappa)^2, \quad (\text{A.131})$$

$$0 = \nabla^a \left(e^{4\alpha_1 \phi} \nabla_a \kappa \right) - F_{ab} (*F)^{ab}, \quad (\text{A.132})$$

$$0 = \nabla_a \left(e^{-2\alpha_0 \phi} F^{ab} \right) + (*F)^{ab} \nabla_a \kappa. \quad (\text{A.133})$$

We include a form of hyperbolic constraint damping into the Maxwell equations so that violations propagate away at the speed of light. In particular, we introduce two scalar fields Φ and Ψ into Eq. (A.133) and the identity $\nabla_a (*F)^{ab} = 0$ as follows:

$$\nabla_a \left(F^{ab} + g^{ab} \Psi \right) = \eta_1 n^a \Psi + 2\alpha_0 \nabla_a \phi F^{ab} - e^{2\alpha_0 \phi} (*F)^{ab} \nabla_a \kappa, \quad (\text{A.134})$$

$$\nabla_a \left((*F)^{ab} + g^{ab} \Phi \right) = \eta_2 n^a \Phi. \quad (\text{A.135})$$

where η_1 and η_2 are taken as positive constants and serve as tunable parameters to effect the damping of the electromagnetic constraints. Note that on the right-hand sides, we have also included n^a , the usual unit timelike normal to the spatial hypersurfaces in anticipation of performing the usual 3 + 1 split of spacetime. It is worth noting that these damped versions of Maxwell are now no longer strictly four-covariant. We introduce the usual 3 + 1 decomposition of spacetime by defining the line element as

$$ds^2 = -\alpha^2 dt^2 + \gamma_{ij} (dx^i + \beta^i dt) (dx^j + \beta^j dt) \quad (\text{A.136})$$

with lapse, α , shift, β^i , 3-metric, γ_{ij} , unit timelike normal $n_a = (-\alpha, 0, 0, 0)$ and extrinsic curvature $K_{ab} = -\gamma_a^c \gamma_b^d \nabla_c n_d$.

Using the BSSN formulation, we solve the standard BSSN evolution equations for the usual $\chi, \tilde{\gamma}_{ij}, K, \tilde{A}_{ij}$ and $\tilde{\Gamma}^i$ variables with EMDA matter terms.¹ These equations take the form

$$\partial_0 \tilde{\gamma}_{ij} = -2\alpha \tilde{A}_{ij}, \quad (\text{A.137})$$

$$\partial_0 \chi = \frac{2}{3} \alpha \chi K, \quad (\text{A.138})$$

$$\partial_0 K = [\dots] + 4\pi\alpha(\rho + T), \quad (\text{A.139})$$

$$\partial_0 \tilde{A}_{ij} = [\dots] - 8\pi\chi\alpha \left(T_{ij} - \frac{1}{3} \gamma_{ij} T_k^k \right), \quad (\text{A.140})$$

$$\partial_0 \tilde{\Gamma}^i = [\dots] - 16\pi\alpha \chi^{-1} S^i, \quad (\text{A.141})$$

where $\partial_0 \equiv \partial_t - \mathcal{L}_\beta$ with $\mathcal{L}_\beta$ the Lie derivative along the shift and the $[\dots]$ are a stand-in for the relevant right hand sides which can be found in standard references, e.g. [86]

The quantities S^i , T_{ij} , and ρ are defined as

$$\begin{aligned} S^i &\equiv -\gamma^{ia} T_{ab} n^b \\ &= 2\Pi D^i \phi + 2e^{-2\alpha_0 \phi} \epsilon^{ijk} E_j B_k + \frac{1}{2} e^{4\alpha_1 \phi} \Xi D^i \kappa, \end{aligned} \quad (\text{A.142})$$

$$\begin{aligned} T_{ij} &\equiv T_{ab} \gamma^a_i \gamma^b_j \\ &= 2 \left[D_i \phi D_j \phi - \frac{1}{2} \tilde{\gamma}_{ij} ((D\phi)^2 - \Pi^2) \right] - 2e^{-2\alpha_0 \phi} \left(E_i E_j + B_i B_j - \frac{1}{2} \tilde{\gamma}_{ij} (E_k E^k + B_k B^k) \right) \\ &\quad + \frac{1}{2} e^{4\alpha_1 \phi} \left[D_i \kappa D_j \kappa - \frac{1}{2} \tilde{\gamma}_{ij} ((D\kappa)^2 - \Xi^2) \right]. \end{aligned} \quad (\text{A.143})$$

$$\rho \equiv \Pi^2 + (D\phi)^2 + e^{-2\alpha_0 \phi} (E_a E^a + B_a B^a) + \frac{1}{4} e^{4\alpha_1 \phi} (\Xi^2 + (D\kappa)^2). \quad (\text{A.144})$$

¹Note that in this Appendix only, χ represents the conformal factor and not the dimensionless spin parameter.

We use standard puncture gauge conditions for the lapse and the shift:

$$\partial_t \alpha = \beta^i \partial_i - 2\alpha K, \quad (\text{A.145})$$

$$\partial_t \beta^i = \beta^j \partial_j \beta^i + \frac{3}{4} B^i, \quad (\text{A.146})$$

$$\partial_t B^i = \beta^j \partial_j B^i + \partial_t \tilde{\Gamma}^i - \beta^j \partial_j \tilde{\Gamma}^i - \eta B^i. \quad (\text{A.147})$$

We also employ a slow-start lapse condition [126] to reduce transients at early times. This amounts to adding the term

$$-\chi^{1/2} (\varpi e^{-t^2/2\sigma^2}) (\alpha - \chi^{1/2}) \quad (\text{A.148})$$

to the equation for the lapse. Following [126], we take the constants $\varpi = 3/5$ and $\sigma = 20$.

For the matter equations we start with the dilaton and axion:

$$\partial_0 \phi = -\alpha \Pi, \quad (\text{A.149})$$

$$\partial_0 \kappa = -\Xi \quad (\text{A.150})$$

$$\begin{aligned} \partial_0 \Pi = & \alpha K \Pi - \alpha \chi \tilde{\gamma}^{ij} \partial_i \partial_j \phi - \chi \tilde{\gamma}^{ij} \partial_i \alpha \partial_j \phi + \alpha \chi \tilde{\Gamma}^i \partial_i \phi + \frac{\alpha}{2} \tilde{\gamma}^{ij} \partial_i \chi \partial_j \phi \\ & + \frac{\alpha \alpha_0}{\chi} e^{-2\alpha_0 \phi} \tilde{\gamma}_{ij} (E^i E^j - B^i B^j) + \frac{\alpha \alpha_1}{2} e^{4\alpha_1 \phi} \left(\chi \tilde{\gamma}^{ij} \partial_i \kappa \partial_j \kappa - \Xi^2 \right). \end{aligned} \quad (\text{A.151})$$

$$\begin{aligned} \partial_0 \Xi = & \alpha K \Xi - \alpha \chi \tilde{\gamma}^{ij} \partial_i \partial_j \kappa - \chi \tilde{\gamma}^{ij} \partial_i \alpha \partial_j \kappa + \alpha \chi \tilde{\Gamma}^i \partial_i \kappa + \frac{\alpha}{2} \tilde{\gamma}^{ij} \partial_i \kappa \partial_j \chi \\ & - 4\alpha \alpha_1 \chi \tilde{\gamma}^{ij} \partial_i \kappa \partial_j \phi + 4\alpha \alpha_1 \Pi \Xi + \frac{4\alpha}{\chi} e^{-4\alpha_1 \phi} \tilde{\gamma}_{ij} B^i E^j. \end{aligned} \quad (\text{A.152})$$

The equations for the electromagnetic fields, including hyperbolic divergence cleaning become

$$\partial_0 \Psi = -\alpha \partial_i E^i + \frac{3\alpha}{2\chi} E^i \partial_i \chi + 2\alpha_0 \alpha E^i \partial_i \phi + \alpha e^{2\alpha_0 \phi} B^i \partial_i \kappa - \alpha \eta_1 \Psi. \quad (\text{A.153})$$

$$\partial_0 \Phi = \alpha \partial_i B^i - \frac{3\alpha}{2\chi} B^i \partial_i \chi - \alpha \eta_2 \Phi. \quad (\text{A.154})$$

$$\begin{aligned} \partial_0 E^i = & \alpha K E^i + 2\alpha \alpha_0 \chi^{1/2} \epsilon^{jik} \tilde{\gamma}_{kl} B^l \partial_j \phi - 2\alpha \alpha_0 E^i \Pi \\ & + \alpha e^{2\alpha_0 \phi} \chi^{1/2} \epsilon^{ijk} \tilde{\gamma}_{kl} E^l \partial_j \kappa - \alpha e^{2\alpha_0 \phi} \Xi B^i - \alpha \chi \tilde{\gamma}^{ij} \partial_j \Psi - \chi^{1/2} \epsilon^{jik} \tilde{\gamma}_{kl} B^l \partial_j \alpha \\ & - \alpha \chi^{1/2} \epsilon^{jik} B^l \partial_j \tilde{\gamma}_{kl} - \alpha \chi^{1/2} \epsilon^{jik} \tilde{\gamma}_{kl} \partial_j B^l + \alpha \chi^{-1/2} \epsilon^{jik} \tilde{\gamma}_{kl} B^l \partial_j \chi. \end{aligned} \quad (\text{A.155})$$

$$\partial_0 B^i = \alpha K B^i + \alpha \chi \tilde{\gamma}^{ij} \partial_j \Phi$$

$$- \chi^{1/2} \epsilon^{ijk} \tilde{\gamma}_{kl} E^l \partial_j \alpha + \alpha \chi^{-1/2} \epsilon^{ijk} \tilde{\gamma}_{kl} E^l \partial_j \chi - \alpha \chi^{1/2} \epsilon^{ijk} E^l \partial_j \tilde{\gamma}_{kl} - \alpha \chi^{1/2} \epsilon^{ijk} \tilde{\gamma}_{kl} \partial_j E^l.$$

(A.156)

Appendix B

EMDA Black Holes

B.1 Schwarzschild Metric

We present the Schwarzschild solution in standard Schwarzschild coordinates. This solution describes a static, spherically symmetric, uncharged black hole in general relativity. The metric takes the form

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 - \frac{2M}{r}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2. \quad (\text{B.1})$$

Equivalently, this can be written as

$$ds^2 = -f(r) dt^2 + f(r)^{-1} dr^2 + r^2 d\Omega^2, \quad (\text{B.2})$$

where

$$f(r) = 1 - \frac{2M}{r}, \quad (\text{B.3})$$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2. \quad (\text{B.4})$$

The Schwarzschild solution is characterized by a single physical parameter, the ADM mass M . Since the spacetime is nonrotating and uncharged, the electric charge, angular momentum, and magnetic dipole moment vanish:

$$Q = 0, \quad (\text{B.5})$$

$$J = 0, \quad (\text{B.6})$$

$$\mu = 0. \quad (\text{B.7})$$

In the absence of additional matter fields, the associated gauge, dilaton, and axion fields are trivial:

$$A_a dx^a = 0, \quad (\text{B.8})$$

$$\phi = 0, \quad (\text{B.9})$$

$$\kappa = 0. \quad (\text{B.10})$$

With respect to the Schwarzschild radial coordinate r , the event horizon is located at

$$r_H = 2M. \quad (\text{B.11})$$

The curvature singularity occurs at

$$r = 0, \quad (\text{B.12})$$

while the apparent coordinate singularity at $r = 2M$ corresponds to the event horizon rather than a physical singularity.

The Schwarzschild solution may be obtained as the nonrotating, uncharged limit of the Kerr–Sen solution by taking

$$a \rightarrow 0, \quad \alpha \rightarrow 0, \quad m \rightarrow M. \quad (\text{B.13})$$

In this limit, the Kerr–Sen metric functions reduce to

$$\Delta = r^2 - 2Mr, \quad (\text{B.14})$$

$$\rho^2 = r^2, \quad (\text{B.15})$$

$$\Sigma = r^4, \quad (\text{B.16})$$

and the Kerr–Sen line element becomes the Schwarzschild line element.

B.2 Reissner–Nordström Metric

We present the Reissner–Nordström solution in standard Schwarzschild-like coordinates. This solution describes a static, spherically symmetric, electrically charged black hole in general relativity coupled to electromagnetism. The metric takes the form

$$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2. \quad (\text{B.17})$$

Equivalently, this can be written as

$$ds^2 = -f(r) dt^2 + f(r)^{-1} dr^2 + r^2 d\Omega^2, \quad (\text{B.18})$$

where

$$f(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2}, \quad (\text{B.19})$$

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2. \quad (\text{B.20})$$

The associated electromagnetic potential may be written as

$$A_a dx^a = -\frac{Q}{r} dt, \quad (\text{B.21})$$

up to a gauge transformation. Equivalently, one may choose the opposite sign convention,

$$A_a dx^a = \frac{Q}{r} dt, \quad (\text{B.22})$$

depending on the convention used for the Maxwell tensor. In either case, the corresponding electric field has magnitude proportional to Q/r^2 .

The Reissner–Nordström solution is characterized by two physical parameters: the ADM mass M and the electric charge Q . Since the spacetime is nonrotating, the angular momentum and magnetic dipole moment vanish:

$$J = 0, \quad (\text{B.23})$$

$$\mu = 0. \quad (\text{B.24})$$

In the absence of additional scalar fields, the dilaton and axion are trivial:

$$\phi = 0, \quad (\text{B.25})$$

$$\kappa = 0. \quad (\text{B.26})$$

With respect to the Schwarzschild-like radial coordinate r , the horizons are located at the roots of $f(r) = 0$:

$$r_{\pm} = M \pm \sqrt{M^2 - Q^2}. \quad (\text{B.27})$$

The outer event horizon is therefore

$$r_H = r_+ = M + \sqrt{M^2 - Q^2}. \quad (\text{B.28})$$

The condition for the existence of a nonextremal Reissner–Nordström black hole is

$$M^2 > Q^2. \quad (\text{B.29})$$

Equivalently, in terms of the dimensionless charge-to-mass ratio,

$$\lambda = \frac{Q}{M}, \quad (\text{B.30})$$

the nonextremality condition is

$$|\lambda| < 1. \quad (\text{B.31})$$

The curvature singularity occurs at

$$r = 0. \quad (\text{B.32})$$

The apparent coordinate singularities at $r = r_{\pm}$ correspond to the inner and outer horizons.

The Reissner–Nordström solution reduces to the Schwarzschild solution in the uncharged limit,

$$Q \rightarrow 0. \quad (\text{B.33})$$

In this limit,

$$f(r) \rightarrow 1 - \frac{2M}{r}, \quad (\text{B.34})$$

$$r_+ \rightarrow 2M, \quad (\text{B.35})$$

$$r_- \rightarrow 0, \quad (\text{B.36})$$

and the line element becomes the Schwarzschild line element.

B.3 Kerr Metric

We present the Kerr solution in Boyer–Lindquist coordinates. This solution describes a stationary, axisymmetric, uncharged rotating black hole in general relativity. The metric takes the form

$$ds^2 = -\left(1 - \frac{2Mr}{\rho^2}\right) dt^2 + \rho^2 \left(\frac{dr^2}{\Delta} + d\theta^2\right) + \frac{\Sigma}{\rho^2} \sin^2 \theta d\varphi^2 - \frac{4Mar}{\rho^2} \sin^2 \theta dt d\varphi, \quad (\text{B.37})$$

where the metric functions are defined by

$$\Delta = r^2 - 2Mr + a^2, \quad (\text{B.38})$$

$$\rho^2 = r^2 + a^2 \cos^2 \theta, \quad (\text{B.39})$$

$$\Sigma = (r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta. \quad (\text{B.40})$$

The Kerr solution is characterized by two physical parameters: the ADM mass M and the angular momentum J . The spin parameter a is related to these quantities by

$$a = \frac{J}{M}. \quad (\text{B.41})$$

Since the Kerr black hole is uncharged, the electric charge and magnetic dipole moment vanish:

$$Q = 0, \quad (\text{B.42})$$

$$\mu = 0. \quad (\text{B.43})$$

In the absence of additional matter fields, the associated gauge, dilaton, and axion fields are trivial:

$$A_a dx^a = 0, \quad (\text{B.44})$$

$$\phi = 0, \quad (\text{B.45})$$

$$\kappa = 0. \quad (\text{B.46})$$

With respect to the Boyer–Lindquist radial coordinate r , the event horizons are located at the roots of $\Delta = 0$:

$$r_{\pm} = M \pm \sqrt{M^2 - a^2}. \quad (\text{B.47})$$

The outer event horizon is therefore

$$r_H = r_+ = M + \sqrt{M^2 - a^2}. \quad (\text{B.48})$$

The condition for the existence of a nonextremal Kerr black hole is

$$M^2 > a^2 \quad \Leftrightarrow \quad M^4 > J^2. \quad (\text{B.49})$$

Equivalently, in terms of the dimensionless spin parameter

$$\chi = \frac{J}{M^2} = \frac{a}{M}, \quad (\text{B.50})$$

the nonextremality condition is

$$|\chi| < 1. \quad (\text{B.51})$$

The curvature singularity occurs where

$$\rho^2 = r^2 + a^2 \cos^2 \theta = 0, \quad (\text{B.52})$$

which corresponds to the Kerr ring singularity at

$$r = 0, \quad \theta = \frac{\pi}{2}. \quad (\text{B.53})$$

The Kerr solution may be obtained as the uncharged limit of the Kerr–Sen solution by taking

$$\alpha \rightarrow 0, \quad m \rightarrow M. \quad (\text{B.54})$$

In this limit, the Kerr–Sen fields become trivial,

$$A_a dx^a = 0, \quad (\text{B.55})$$

$$\phi = 0, \quad (\text{B.56})$$

$$\kappa = 0, \quad (\text{B.57})$$

and the Kerr–Sen metric functions reduce to the Kerr metric functions

$$\Delta = r^2 - 2Mr + a^2, \quad (\text{B.58})$$

$$\rho^2 = r^2 + a^2 \cos^2 \theta, \quad (\text{B.59})$$

$$\Sigma = (r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta. \quad (\text{B.60})$$

B.4 Kerr–Newman Metric

We present the Kerr–Newman solution in Boyer–Lindquist coordinates. This solution describes a stationary, axisymmetric, electrically charged rotating black hole in Einstein–Maxwell theory. The metric takes the form

$$ds^2 = - \left(1 - \frac{2Mr - Q^2}{\rho^2} \right) dt^2 + \rho^2 \left(\frac{dr^2}{\Delta} + d\theta^2 \right) + \frac{\Sigma}{\rho^2} \sin^2 \theta d\varphi^2 - \frac{2a(2Mr - Q^2)}{\rho^2} \sin^2 \theta dt d\varphi, \quad (\text{B.61})$$

where the metric functions are defined by

$$\Delta = r^2 - 2Mr + a^2 + Q^2, \quad (\text{B.62})$$

$$\rho^2 = r^2 + a^2 \cos^2 \theta, \quad (\text{B.63})$$

$$\Sigma = (r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta. \quad (\text{B.64})$$

The associated electromagnetic potential may be written as

$$A_a dx^a = - \frac{Qr}{\rho^2} \left(dt - a \sin^2 \theta d\varphi \right), \quad (\text{B.65})$$

up to a gauge transformation and sign convention.

The Kerr–Newman solution is characterized by three physical parameters: the ADM mass M , electric charge Q , and angular momentum J . The spin parameter a is related to the angular momentum by

$$a = \frac{J}{M}. \quad (\text{B.66})$$

The magnetic dipole moment is

$$\mu = aQ. \quad (\text{B.67})$$

In the absence of additional scalar fields, the dilaton and axion are trivial:

$$\phi = 0, \quad (\text{B.68})$$

$$\kappa = 0. \quad (\text{B.69})$$

With respect to the Boyer–Lindquist radial coordinate r , the horizons are located at the roots of $\Delta = 0$:

$$r_{\pm} = M \pm \sqrt{M^2 - a^2 - Q^2}. \quad (\text{B.70})$$

The outer event horizon is therefore

$$r_H = r_+ = M + \sqrt{M^2 - a^2 - Q^2}. \quad (\text{B.71})$$

The condition for the existence of a nonextremal Kerr–Newman black hole is

$$M^2 > a^2 + Q^2. \quad (\text{B.72})$$

Equivalently, using $a = J/M$, this condition may be written as

$$M^4 > J^2 + M^2 Q^2. \quad (\text{B.73})$$

In terms of the dimensionless spin and charge parameters

$$\chi = \frac{J}{M^2} = \frac{a}{M}, \quad (\text{B.74})$$

$$\lambda = \frac{Q}{M}, \quad (\text{B.75})$$

the nonextremality condition becomes

$$\chi^2 + \lambda^2 < 1. \quad (\text{B.76})$$

The curvature singularity occurs where

$$\rho^2 = r^2 + a^2 \cos^2 \theta = 0, \quad (\text{B.77})$$

which corresponds to the ring singularity

$$r = 0, \quad \theta = \frac{\pi}{2}. \quad (\text{B.78})$$

The Kerr–Newman solution reduces to the Kerr solution in the uncharged limit,

$$Q \rightarrow 0, \quad (\text{B.79})$$

for which

$$\Delta \rightarrow r^2 - 2Mr + a^2, \quad (\text{B.80})$$

$$r_+ \rightarrow M + \sqrt{M^2 - a^2}. \quad (\text{B.81})$$

It reduces to the Reissner–Nordström solution in the nonrotating limit,

$$a \rightarrow 0, \quad (\text{B.82})$$

for which

$$\Delta \rightarrow r^2 - 2Mr + Q^2, \quad (\text{B.83})$$

$$r_+ \rightarrow M + \sqrt{M^2 - Q^2}. \quad (\text{B.84})$$

Finally, in the uncharged and nonrotating limit,

$$Q \rightarrow 0, \quad a \rightarrow 0, \quad (\text{B.85})$$

the Kerr–Newman solution reduces to the Schwarzschild solution.

B.5 Einstein–Maxwell–Dilaton Black Holes

We present a static, spherically symmetric black-hole solution in Einstein–Maxwell–Dilaton theory.

In Schwarzschild-like coordinates, the magnetically charged solution takes the form

$$ds^2 = - \left(1 - \frac{r_+}{r}\right) \left(1 - \frac{r_-}{r}\right)^{1-\alpha_1} dt^2 + \left(1 - \frac{r_+}{r}\right)^{-1} \left(1 - \frac{r_-}{r}\right)^{\alpha_1-1} dr^2 + r^2 \left(1 - \frac{r_-}{r}\right)^{\alpha_1} d\Omega^2, \quad (\text{B.86})$$

where

$$d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2. \quad (\text{B.87})$$

The coupling-dependent parameter α_1 is defined by

$$\alpha_1 = \frac{2\alpha_0^2}{1 + \alpha_0^2}, \quad (\text{B.88})$$

where α_0 is the dilaton coupling constant.

For the magnetic solution, the Maxwell field and dilaton are

$$F_{\theta\varphi} = Q_m \sin \theta, \quad (\text{B.89})$$

$$e^{-2\alpha_0\phi} = e^{-2\alpha_0\phi_0} \left(1 - \frac{r_-}{r}\right)^{\alpha_1}, \quad (\text{B.90})$$

where Q_m is the magnetic charge and ϕ_0 is the asymptotic value of the dilaton at spatial infinity.

The constants r_+ and r_- are related to the ADM mass M , magnetic charge Q_m , and asymptotic dilaton value ϕ_0 by

$$2M = r_+ + (1 - \alpha_1)r_-, \quad (\text{B.91})$$

$$2Q_m^2 = e^{2\alpha_0\phi_0} r_+ r_- (2 - \alpha_1). \quad (\text{B.92})$$

The surface $r = r_+$ corresponds to the event horizon, while $r = r_-$ corresponds to a curvature singularity. For $r > r_-$, the dilaton is larger than its asymptotic value and monotonically decreases toward ϕ_0 at spatial infinity.

The theory also admits an electrically charged solution related to the magnetic solution by a discrete electromagnetic duality. Under this duality, the metric is unchanged, while the Maxwell field and dilaton transform according to

$$F_{ab} \rightarrow e^{-2\alpha_0\phi} (*F)_{ab}, \quad \phi \rightarrow -\phi. \quad (\text{B.93})$$

For the electric solution, the Maxwell field and dilaton are

$$F_{tr} = \frac{Q_e}{r^2}, \quad (\text{B.94})$$

$$e^{2\alpha_0\phi} = e^{2\alpha_0\phi_0} \left(1 - \frac{r_-}{r}\right)^{\alpha_1}, \quad (\text{B.95})$$

where Q_e is the electric charge. The constants r_+ and r_- then satisfy

$$2M = r_+ + (1 - \alpha_1)r_-, \quad (\text{B.96})$$

$$2Q_e^2 = e^{-2\alpha_0\phi_0} r_+ r_- (2 - \alpha_1). \quad (\text{B.97})$$

In this case, the solution has ADM mass M , electric charge Q_e , and asymptotic dilaton value ϕ_0 . The event horizon is located at $r = r_+$, and the curvature singularity is located at $r = r_-$. For $r > r_-$, the dilaton monotonically increases toward ϕ_0 at spatial infinity.

Since this solution is static and spherically symmetric, the angular momentum and magnetic dipole moment vanish:

$$J = 0, \quad (\text{B.98})$$

$$\mu = 0. \quad (\text{B.99})$$

Several familiar black-hole solutions are recovered as special limits. In the limit of vanishing dilaton coupling,

$$\alpha_0 \rightarrow 0, \quad \alpha_1 \rightarrow 0, \quad (\text{B.100})$$

the solution reduces to the Reissner–Nordström black hole. In the uncharged limit,

$$Q_m \rightarrow 0 \quad \text{or} \quad Q_e \rightarrow 0, \quad (\text{B.101})$$

one has $r_- \rightarrow 0$, and the solution reduces to the Schwarzschild black hole with

$$r_+ = 2M. \quad (\text{B.102})$$

B.6 Kerr–Sen Metric

We present the Kerr–Sen solution in the Einstein frame, written in Boyer–Lindquist type coordinates.

The metric takes the form

$$ds^2 = - \left(1 - \frac{2mr \cosh^2 \alpha}{\rho^2} \right) dt^2 + \rho^2 \left(\frac{dr^2}{\Delta} + d\theta^2 \right) + \frac{\Sigma}{\rho^2} \sin^2 \theta d\varphi^2 - \frac{4amr \cosh^2 \alpha}{\rho^2} \sin^2 \theta dt d\varphi \quad (\text{B.103})$$

where the metric functions are defined by

$$\Delta = r^2 - 2mr + a^2, \quad (\text{B.104})$$

$$\rho^2 = r^2 + a^2 \cos^2 \theta + 2mr \sinh^2 \alpha, \quad (\text{B.105})$$

$$\Sigma = \left(r^2 + a^2 + 2mr \sinh^2 \alpha \right)^2 - \Delta a^2 \sin^2 \theta. \quad (\text{B.106})$$

The associated gauge, dilaton, and axion fields are given by

$$A_a dx^a = \frac{mr \sinh 2\alpha}{\sqrt{2} \rho^2} \left(dt - a \sin^2 \theta d\varphi \right), \quad (\text{B.107})$$

$$e^{-2\phi} = \frac{\rho^2}{r^2 + a^2 \cos^2 \theta}, \quad (\text{B.108})$$

$$\kappa = -2ma \sinh^2 \alpha \frac{\cos \theta}{r^2 + a^2 \cos^2 \theta}. \quad (\text{B.109})$$

The parameters m , α , and a arise as integration constants. The physical mass M , electric charge Q , angular momentum J , and magnetic dipole moment μ are related to these parameters via

$$M = m \cosh^2 \alpha, \quad (\text{B.110})$$

$$Q = \frac{m}{\sqrt{2}} \sinh 2\alpha, \quad (\text{B.111})$$

$$J = aM, \quad (\text{B.112})$$

$$\mu = aQ. \quad (\text{B.113})$$

With respect to the Boyer–Lindquist radial coordinate, r , the event horizon is located at the familiar

$$r_H = m + \sqrt{m^2 - a^2}, \quad (\text{B.114})$$

and the condition for the existence of a nonextremal black hole is

$$m^2 > a^2 \quad \Leftrightarrow \quad M^2 > |J| + \frac{Q^2}{2}. \quad (\text{B.115})$$

The inverse relations are

$$m = M - \frac{Q^2}{2M}, \quad (\text{B.116})$$

$$\sinh 2\alpha = \frac{2\sqrt{2}QM}{2M^2 - Q^2}, \quad (\text{B.117})$$

$$a = \frac{J}{M}. \quad (\text{B.118})$$

Appendix C

EMDA Initital data

C.1 Initial Data

It is necessary to have good initial data for our EMDA BH binaries. Of course, exact solutions are not known. But we can generalize standard approaches for the construction of initial data in vacuum GR to these additional fields on making certain approximations. We follow closely the approach based on the conformal transverse-traceless decomposition used with punctures. Indeed, we extend this approach as it has been incorporated into the `TwoChargedPunctures` code of Bozzola and Paschalidis [97, 123]. We mostly follow their notation below.

As usual, we decompose the metric, but using notation slightly modified from the previous appendix. In particular we define

$$\gamma_{ij} = \psi^4 \bar{\gamma}_{ij}, \quad (\text{C.1})$$

with ψ the conformal factor and the conformal metric assumed to be the flat metric, $\bar{\gamma}_{ij} = \eta_{ij}$. The latter assumption simplifies the following computations and is a reasonable choice for the systems we study. It is a limiting assumption preventing the construction of highly spinning black holes [127, 128]. Notwithstanding, Bowen–York black holes can reach spin values of order $\chi \sim 0.9$ [129]. As our simulations remain below this limit, the approximation should be reasonable. While we anticipate the appearance of “junk radiation” partly in consequence of this assumption [111], we find that it does decay as the fields relax to quasi-equilibrium which results in decreased constraint violation over the course of the evolutions. We adopt the maximal slicing condition, $K = 0$, and define a conformally rescaled tracefree extrinsic curvature, $\bar{A}_{ij} = \psi^2 (K_{ij} - \frac{1}{3} \gamma_{ij} K)$. Ignoring the latter’s radiative degrees of freedom, we can express it in terms of a vector, V^i , as

$$\bar{A}_{ij} = \partial_i V_j + \partial_j V_i - \frac{2}{3} \eta_{ij} \partial_k V^k. \quad (\text{C.2})$$

The initial data problem then reduces to determining V^i and ψ for the gravitational pieces. In particular, the Hamiltonian and momentum constraint equations take the form

$$\partial_k \partial^k \psi + \frac{1}{8} \psi^{-7} \bar{A}_{ij} \bar{A}^{ij} = -2\pi \psi^5 \rho \quad (\text{C.3})$$

$$\partial_k \partial^k V^i + \frac{1}{3} \eta^{ij} \partial_j (\partial_k V^k) = 8\pi S^i. \quad (\text{C.4})$$

The momentum constraint is linear and independent of ψ . This permits dividing the solution between homogeneous (vacuum gravitational) and inhomogeneous (matter) contributions:

$$V^i = V_{\text{GR}}^i + V_{\text{M}}^i. \quad (\text{C.5})$$

For V_{GR}^i , we use the standard Bowen–York solutions [125] while for the matter contribution, we must solve Eq. C.4 with

$$S^i = 2\Pi D^i \phi + 2e^{-2\alpha_0 \phi} \epsilon^{ijk} E_j B_k + \frac{1}{2} e^{4\alpha_1 \phi} \Xi D^i \kappa. \quad (\text{C.6})$$

If we assume a moment of time symmetry and neglect contributions from the magnetic field this source term vanishes. On imposing asymptotically flat boundary conditions, the solution to the matter part therefore becomes $V_{\text{M}}^i = 0$.

We must also impose the electromagnetic constraints. At this point we depart somewhat from [97] because the EMDA version of these constraints are no longer linear, containing rather a nontrivial current that couples the dilaton and axion to the electromagnetic fields:

$$\nabla_i E^i = 2\alpha_0 E^i \partial_i \phi + e^{2\alpha_0 \phi} B^i \partial_i \kappa \quad (\text{C.7})$$

$$\nabla_i B^i = 0 \quad (\text{C.8})$$

Were the source in the Gauss constraint zero, we would follow [93, 96] and perform a conformal rescaling of the fields as

$$\bar{E}^i = \psi^6 E^i, \quad \bar{B}^i = \psi^6 B^i. \quad (\text{C.9})$$

This results in the sourceless Maxwell constraints taking the form

$$\partial_i \bar{E}^i = 0, \quad \partial_i \bar{B}^i = 0, \quad (\text{C.10})$$

independent of ψ and therefore solvable separately from the spacetime variables. Since these equations are linear, solutions may be superposed. The advantages of this are significant so we choose to make the approximation that this source term is, in fact, negligible and we solve the EMDA electromagnetic constraints in the way just described. Of course, this failure to self-consistently solve these constraints necessarily introduces violations. Partly for this reason, we have introduced hyperbolic divergence cleaning to drive these violations smaller over the course of the evolution. We have measured these violations on the initial slice. For the simulations considered in this work we confirm that they are consistently small and driven smaller via the constraint damping. The expected damping behavior is demonstrated in Figure ???. There is an initial violation of the EM constraints, which damps over time.

The Hamiltonian constraint above, not being linear, must be treated differently. We assume that we have two black holes charged under the U(1) gauge field as well as possessing scalar dilaton and axion fields. Following [97], we approximate this with a conformal factor of the form

$$\psi^2 = (1 + u + \bar{\eta})^2 - \bar{\xi}^2 \quad (\text{C.11})$$

for which we define

$$\bar{\eta} = \frac{m_1}{2R_1} + \frac{m_2}{2R_2}, \quad (\text{C.12})$$

$$\bar{\xi} = \frac{q_1}{2R_1} + \frac{q_2}{2R_2}, \quad (\text{C.13})$$

$$\bar{\kappa} = 1 + u + \bar{\eta}, \quad (\text{C.14})$$

and where m_i , q_i and R_i here are the masses, electric charges and the coordinate distances from the origin of the BHs. This is intended to correspond to two electrically charged BHs with the singular part of the conformal factor separated out with u giving finite corrections, including the necessary corrections for the additional scalar fields.

On rearranging the Hamiltonian constraint in terms of u , we find

$$\partial_k \partial^k u = -\frac{1}{\bar{\kappa}} \left(\partial_a \bar{\kappa} \partial^a \bar{\kappa} - \partial_a \bar{\xi} \partial^a \bar{\xi} - \partial_a \psi \partial^a \psi + \frac{1}{8} \psi^{-6} \bar{A}_{ij} \bar{A}^{ij} + 2\pi \psi^6 \rho \right), \quad (\text{C.15})$$

with

$$\begin{aligned} \rho \equiv & \left[\Pi^2 + (D\phi)^2 \right] + e^{-2\alpha_0\phi} \left[E_a E^a + B_a B^a \right] \\ & + \frac{1}{4} e^{4\alpha_1\phi} \left[\Xi^2 + (D\kappa)^2 \right]. \end{aligned} \quad (\text{C.16})$$

Using simplifying assumptions similar to above, namely a moment of time symmetry and neglecting the magnetic field, ρ reduces to

$$\rho = (D\phi)^2 + e^{-2\alpha_0\phi} E_a E^a + \frac{1}{4} e^{4\alpha_1\phi} (D\kappa)^2. \quad (\text{C.17})$$

Again, this follows [97] quite closely, except that we have now included dilaton and axion contributions in ρ . Specifically, we employ a Reissner–Nordström electric field for the electric component and add a dilaton field for a single, spinless black hole. In practice, the axion is initialized to zero and allowed to relax to a quasi-equilibrium state during an evolution.

The electric field is initially set to be:

$$E^i = \psi^{-6} \left[\frac{q_1}{R_1^2} + \frac{q_2}{R_2^2} \right]. \quad (\text{C.18})$$

while the dilaton fields are initially set as:

$$\phi = \phi_1(r_1, \theta_1) + \phi_2(r_2, \theta_2) \quad (\text{C.19})$$

where Eq. B.108 gives the form for a dilaton field for a single Kerr–Sen black hole (albeit in Boyer–Lindquist type coordinates) for our current theory parameters of $(\alpha_0, \alpha_1) = (1, 1)$.

C.1.1 TPID Solver

Our charged-puncture initial-data solver is based on existing TwoPunctures-style methods. The discussion here follows closely the presentation of Ref. [97], as well as the original TwoPunctures paper [130]. The solver is used to construct initial data for two charged black holes separated along the x axis, with punctures located at $x = \pm b$. The user specifies the bare parameters of the punctures, including their masses, charges, linear momenta, and spins. These parameters determine the source terms that enter the constraint equations.

The TPID solver uses a pseudo-spectral collocation method. In contrast to a finite-difference method, where variables are stored on a grid and derivatives are approximated using local stencils, a pseudo-spectral method represents the unknown correction functions using global basis functions. The equations are then enforced at a discrete set of collocation points. Once the solution is obtained at these points, it is interpolated onto the numerical grid used for the time evolution.

The coordinate system used by TwoPunctures is designed to cover all of $\mathbb{R}^3$ with a single compactified computational domain. The physical coordinates (x, y, z) are parameterized by computational coordinates (A, B, ϕ) , where $A, B \in [-1, 1]$ and $\phi \in [0, 2\pi]$. The transformation is given by

$$x = b \frac{(1+A)^2 + 4}{(1+A)^2 - 4} \frac{2B}{1+B^2}, \quad (\text{C.20})$$

$$y = b \frac{4(1+A)}{4 - (1+A)^2} \frac{1 - B^2}{1+B^2} \cos \phi, \quad (\text{C.21})$$

$$z = b \frac{4(1+A)}{4 - (1+A)^2} \frac{1 - B^2}{1+B^2} \sin \phi. \quad (\text{C.22})$$

These coordinates form a cylindrical-like coordinate system around the x axis, with the mapping depending on both A and B . The punctures are located on the x axis, while the outer boundary of the compactified domain corresponds to spatial infinity. In particular, the limit $A \rightarrow 1$ maps to spatial infinity. This compactification makes it straightforward to impose the boundary condition that the correction to the conformal factor vanishes at infinity.

The computational coordinates are discretized using n_A , n_B , and n_ϕ collocation points. An example of this grid is shown in Fig. C.1. In the A and B directions, these points are chosen from the roots of Chebyshev polynomials, which concentrate resolution near the punctures and in regions where the solution varies rapidly. The unknown correction functions are expanded spectrally, and the expansion coefficients are determined by requiring the constraint equations to be satisfied at the collocation points.

Because the constraint equations are nonlinear, the resulting algebraic system is solved iteratively using a Newton–Raphson method. At each iteration, the residuals of the constraint equations are evaluated at the collocation points. These residuals measure the extent to which the current approximate solution fails to satisfy the constraint equations. For an exact solution, the residuals would vanish at every collocation point. In practice, the Newton–Raphson iteration is continued until a chosen norm of the residuals falls below a prescribed tolerance, indicating that the constraints are satisfied to the desired numerical accuracy. The resulting spectral solution is then interpolated from the TwoPunctures collocation points onto the evolution grid used in the numerical-relativity simulation.




Figure C.1 Example of the compactified TwoPunctures grid in a slice through the two punctures. The physical coordinates are shown in units of the half-separation b , with the black-hole punctures located at $x = \pm b$. The coordinate lines illustrate how the pseudo-spectral collocation points are distributed in the compactified domain before the resulting initial data are interpolated onto the evolution grid. The outer compactified boundary corresponds to spatial infinity and is outside the finite plotting window.

Appendix D

Computation of Ψ_4

D.1 Wave Extraction

This appendix summarizes the wave-extraction procedure used for the gravitational, electromagnetic, dilaton, and axion radiation. The gravitational radiation is extracted from the Newman–Penrose scalar Ψ_4 , while the electromagnetic radiation is extracted from the Maxwell Newman–Penrose scalars Φ_1 and Φ_2 . We also define analogous outgoing scalar quantities for the dilaton and axion fields by projecting their gradients along an outgoing null direction.

D.1.1 Null Tetrad and Spatial Triad

The Newman–Penrose scalar Ψ_4 is defined by projecting the Weyl tensor onto a null tetrad:

$$\Psi_4 = C_{abcd} k^a \bar{m}^b k^c \bar{m}^d. \quad (\text{D.1})$$

Here C_{abcd} is the Weyl tensor, and k^a and $\bar{m}^a$ are elements of a null tetrad. The tetrad is constructed from the future-directed unit normal n^a to the spatial hypersurface and an orthonormal spatial triad r^a , θ^a , and φ^a , which asymptotically approach the usual radial, polar, and azimuthal directions. We define

$$\begin{aligned} l^a &= \frac{1}{\sqrt{2}} (n^a + r^a), \\ k^a &= \frac{1}{\sqrt{2}} (n^a - r^a), \\ m^a &= \frac{1}{\sqrt{2}} (\theta^a + i\varphi^a), \\ \bar{m}^a &= \frac{1}{\sqrt{2}} (\theta^a - i\varphi^a). \end{aligned} \quad (\text{D.2})$$

With metric signature $(-, +, +, +)$, these vectors satisfy

$$l^a k_a = -1, \quad m^a \bar{m}_a = 1, \quad (\text{D.3})$$

with all other tetrad inner products vanishing. In this appendix, l^a denotes the null tetrad vector, while ℓ is reserved for the spherical-harmonic mode index.

In practice, the spatial triad is constructed from coordinate vectors that approach the spherical-coordinate basis vectors in the wave zone. We begin with

$$\begin{aligned} u^a &= (0, x, y, z), \\ v^a &= (0, xz, yz, -x^2 - y^2), \\ w^a &= (0, -y, x, 0). \end{aligned} \tag{D.4}$$

Using the spatial metric γ_{ij} , we define the spatial inner product

$$\langle u, v \rangle = \gamma_{ij} u^i v^j \tag{D.5}$$

and the associated norm

$$|u| = \sqrt{\langle u, u \rangle}. \tag{D.6}$$

The orthonormal triad is then obtained by Gram–Schmidt orthonormalization:

$$\begin{aligned} r^a &= \frac{u^a}{|u|}, \\ \theta^a &= \frac{v^a - \langle r, v \rangle r^a}{|v - \langle r, v \rangle r|}, \\ \varphi^a &= \frac{w^a - \langle r, w \rangle r^a - \langle \theta, w \rangle \theta^a}{|w - \langle r, w \rangle r - \langle \theta, w \rangle \theta|}. \end{aligned} \tag{D.7}$$

D.1.2 Gravitational Radiation

The Weyl tensor is the trace-free part of the spacetime curvature. In four dimensions, the Riemann tensor has twenty independent components. Ten of these are determined by the Ricci tensor, while the remaining ten are contained in the Weyl tensor. The Weyl tensor therefore captures the part of the curvature associated with tidal and radiative degrees of freedom.

In n spacetime dimensions, with $n \geq 4$, the Weyl tensor is

$$\begin{aligned} C_{abcd} &= R_{abcd} + \frac{1}{n-2} (g_{ad} R_{cb} + g_{bc} R_{da} - g_{ac} R_{db} - g_{bd} R_{ca}) \\ &\quad + \frac{1}{(n-1)(n-2)} (g_{ac} g_{db} - g_{ad} g_{cb}) R. \end{aligned} \tag{D.8}$$

In four dimensions, this reduces to

$$C_{abcd} = R_{abcd} + \frac{1}{2} (g_{ad}R_{cb} + g_{bc}R_{da} - g_{ac}R_{db} - g_{bd}R_{ca}) + \frac{1}{6} (g_{ac}g_{db} - g_{ad}g_{cb}) R. \quad (\text{D.9})$$

The Weyl tensor has the same index symmetries as the Riemann tensor:

$$C_{abcd} = -C_{abdc} = -C_{bacd} = C_{cdab}, \quad (\text{D.10})$$

together with the cyclic identity

$$C_{abcd} + C_{adbc} + C_{acdb} = 0. \quad (\text{D.11})$$

At future null infinity, and for an appropriate outgoing tetrad, Ψ_4 encodes the outgoing gravitational radiation. With the convention used here, the complex strain

$$h = h_+ - ih_\times \quad (\text{D.12})$$

is related to Ψ_4 by

$$\Psi_4 = \ddot{h}_+ - i\ddot{h}_\times = \ddot{h}, \quad (\text{D.13})$$

up to the overall sign convention chosen for the null tetrad. Therefore, the strain may be obtained by integrating Ψ_4 twice in time:

$$h(t) = \int^t dt' \int^{t'} dt'' \Psi_4(t''). \quad (\text{D.14})$$

In practice, this integration must be performed carefully because direct time-domain integration can introduce secular drifts. Frequency-domain methods, including fixed-frequency integration, are often used to control low-frequency numerical artifacts.

D.1.3 Electromagnetic, Dilaton, and Axion Radiation

The outgoing electromagnetic radiation is extracted using Newman–Penrose scalars associated with the Maxwell tensor [54, 95]. We define

$$\Phi_1 = \frac{1}{2} F_{ab} \left(l^a k^b + \bar{m}^a m^b \right), \quad (\text{D.15})$$

$$\Phi_2 = F_{ab} \bar{m}^a k^b. \quad (\text{D.16})$$

At large radius, Φ_2 represents the outgoing radiative electromagnetic degree of freedom.

We define analogous outgoing scalar quantities for the dilaton and axion fields by projecting their gradients along the outgoing null direction:

$$\mathcal{D} = l^a \nabla_a \phi, \quad \mathcal{A} = l^a \nabla_a \kappa. \quad (\text{D.17})$$

Here $\mathcal{D}$ and $\mathcal{A}$ denote the extracted dilaton and axion radiation variables, respectively.

D.1.4 Multipolar Decomposition

At an extraction radius R_{ex} , each radiative quantity is decomposed into spin-weighted spherical harmonics with the appropriate spin weight:

$$\Psi_4(t, \theta, \varphi) = \sum_{\ell, m} \Psi_4^{\ell m}(t) {}_{-2}Y_{\ell m}(\theta, \varphi), \quad (\text{D.18})$$

$$\Phi_1(t, \theta, \varphi) = \sum_{\ell, m} \Phi_1^{\ell m}(t) {}_0Y_{\ell m}(\theta, \varphi), \quad (\text{D.19})$$

$$\Phi_2(t, \theta, \varphi) = \sum_{\ell, m} \Phi_2^{\ell m}(t) {}_{-1}Y_{\ell m}(\theta, \varphi), \quad (\text{D.20})$$

$$\mathcal{D}(t, \theta, \varphi) = \sum_{\ell, m} \mathcal{D}^{\ell m}(t) {}_0Y_{\ell m}(\theta, \varphi), \quad (\text{D.21})$$

$$\mathcal{A}(t, \theta, \varphi) = \sum_{\ell, m} \mathcal{A}^{\ell m}(t) {}_0Y_{\ell m}(\theta, \varphi). \quad (\text{D.22})$$

The strain is decomposed similarly as

$$h(t, \theta, \varphi) = \sum_{\ell, m} h_{\ell m}(t) {}_{-2}Y_{\ell m}(\theta, \varphi). \quad (\text{D.23})$$

For each gravitational-wave mode, Eq. (D.13) implies

$$\Psi_4^{\ell m}(t) = \ddot{h}_{\ell m}(t), \quad (\text{D.24})$$

or equivalently

$$h_{\ell m}(t) = \int^t dt' \int^{t'} dt'' \Psi_4^{\ell m}(t''). \quad (\text{D.25})$$

Waveform comparisons in this work primarily use the dominant $(\ell, m) = (2, 2)$ mode.

D.1.5 Radiated Energy Fluxes

The radiated energy fluxes in the gravitational, electromagnetic, dilaton, and axion channels are computed from the multipolar amplitudes. Following Ref. [96], we use

$$F_{\text{GW}} = \frac{dE_{\text{GW}}}{dt} = \lim_{r \rightarrow \infty} \frac{r^2}{16\pi} \sum_{\ell, m} \left| \int_{-\infty}^t \Psi_4^{\ell m}(t') dt' \right|^2, \quad (\text{D.26})$$

$$F_{\text{EM}} = \frac{dE_{\text{EM}}}{dt} = \lim_{r \rightarrow \infty} \frac{r^2}{4\pi} \sum_{\ell, m} |\Phi_2^{\ell m}(t)|^2, \quad (\text{D.27})$$

$$F_{\text{D}} = \frac{dE_{\text{D}}}{dt} = \lim_{r \rightarrow \infty} \frac{r^2}{4\pi} \sum_{\ell, m} |\mathcal{D}^{\ell m}(t)|^2, \quad (\text{D.28})$$

$$F_{\text{A}} = \frac{dE_{\text{A}}}{dt} = \lim_{r \rightarrow \infty} \frac{r^2}{4\pi} \sum_{\ell, m} |\mathcal{A}^{\ell m}(t)|^2. \quad (\text{D.29})$$

In numerical simulations, these quantities are evaluated at finite extraction radii. The expressions above should therefore be interpreted as asymptotic large-radius definitions, with the finite-radius values serving as numerical approximations.

D.1.6 Relation to the 3 + 1 Variables

In numerical relativity, Ψ_4 is computed from quantities defined on spatial hypersurfaces. The spacetime curvature entering the Weyl tensor can therefore be related to the spatial metric γ_{ij} , the extrinsic curvature K_{ij} , and their derivatives using the Gauss–Codazzi relations. Schematically,

$$\gamma^a_i \gamma^b_j \gamma^c_k \gamma^d_l {}^{(4)}R_{abcd} = {}^{(3)}R_{ijkl} + K_{ik}K_{jl} - K_{il}K_{jk}, \quad (\text{D.30})$$

$$\gamma^a_i \gamma^b_j \gamma^c_k n^d {}^{(4)}R_{abcd} = D_j K_{ik} - D_i K_{jk}, \quad (\text{D.31})$$

and

$$\gamma^a_i \gamma^b_j n^c n^d {}^{(4)}R_{acbd} = \mathcal{L}_n K_{ij} + K_{ik} K^k_j + \frac{1}{\alpha} D_i D_j \alpha. \quad (\text{D.32})$$

These identities allow the spacetime curvature appearing in Eq. (D.1) to be expressed in terms of 3 + 1 quantities on each slice.

After substituting the Gauss–Codazzi relations, Ψ_4 can be written in terms of spatial curvature and extrinsic-curvature quantities. A common schematic form is

$$\begin{aligned} \Psi_4 = & \left({}^{(3)}R_{bd} + K K_{bd} - K_{bc} K^c_d \right) \bar{m}^b \bar{m}^d \\ & + (D_c K_{bd} - D_d K_{bc}) r^c \bar{m}^b \bar{m}^d, \end{aligned} \quad (\text{D.33})$$

where the precise signs depend on the conventions used for the Riemann tensor, the extrinsic curvature, and the null tetrad. In non-vacuum evolutions, matter terms may also enter when the curvature is rewritten using the field equations. The important point is that Ψ_4 can be evaluated from the spatial geometry, the extrinsic curvature, and their derivatives on each time slice.

For BSSN-type formulations, the extrinsic curvature is decomposed as

$$K_{ij} = \chi^{-1} \left(\tilde{A}_{ij} + \frac{1}{3} \tilde{\gamma}_{ij} K \right), \quad (\text{D.34})$$

where $\tilde{\gamma}_{ij}$ is the conformal spatial metric, $\tilde{A}_{ij}$ is the trace-free conformal extrinsic curvature, K is the trace of the extrinsic curvature, and χ is the conformal factor used in the evolution system. Equivalently, if $\gamma_{ij} = e^{4\phi} \tilde{\gamma}_{ij}$, then $\chi = e^{-4\phi}$.

The resulting expressions for the real and imaginary parts of Ψ_4 in terms of BSSN variables are lengthy. They are therefore evaluated directly in the code using the evolved variables and their spatial derivatives. Symbolically, one may write

$$\Psi_4 = \text{Re}(\Psi_4) + i \text{Im}(\Psi_4), \quad (\text{D.35})$$

where the real and imaginary parts are obtained by projecting the curvature and extrinsic-curvature terms onto the complex dyad formed from m^a and $\bar{m}^a$.

D.1.7 Waveform Mismatch

To quantify the similarity between two gravitational waveforms, we compute a mismatch. A purely numerical comparison measures the waveform difference directly, while a detector-based comparison weights the difference by the noise curve of a particular detector.

For two complex waveforms $h_1(t)$ and $h_2(t)$, a simple normalized overlap is

$$O = \frac{|\langle h_1, h_2 \rangle|}{\sqrt{\langle h_1, h_1 \rangle \langle h_2, h_2 \rangle}}, \quad (\text{D.36})$$

where a simple time-domain inner product is

$$\langle h_1, h_2 \rangle = \int_{t_1}^{t_2} h_1(t) h_2^*(t) dt. \quad (\text{D.37})$$

The mismatch is then

$$\mathcal{M} = 1 - O. \quad (\text{D.38})$$

In practice, the overlap is maximized over relative time and phase shifts:

$$\mathcal{M} = 1 - \max_{\Delta t, \Delta \phi} \frac{\left| \langle h_1, h_2^{\Delta t, \Delta \phi} \rangle \right|}{\sqrt{\langle h_1, h_1 \rangle \langle h_2, h_2 \rangle}}. \quad (\text{D.39})$$

Here $h_2^{\Delta t, \Delta \phi}$ denotes the waveform h_2 shifted in time by Δt and rotated by an overall phase $\Delta \phi$. This maximization removes trivial offsets in arrival time and phase, leaving a measure of the physical disagreement between the waveform shapes. A smaller mismatch indicates better agreement.

For detector-weighted comparisons, one instead uses the frequency-domain inner product

$$(h_1, h_2) = 4 \operatorname{Re} \int_{f_{\min}}^{f_{\max}} \frac{\tilde{h}_1(f) \tilde{h}_2^*(f)}{S_n(f)} df, \quad (\text{D.40})$$

where $\tilde{h}(f)$ is the Fourier transform of $h(t)$ and $S_n(f)$ is the one-sided detector noise power spectral density. The detector-weighted overlap is

$$\mathcal{O} = \frac{(h_1, h_2)}{\sqrt{(h_1, h_1)(h_2, h_2)}}, \quad (\text{D.41})$$

again maximized over relative time and phase shifts. The corresponding mismatch is

$$\mathcal{M} = 1 - \max_{\Delta t, \Delta \phi} \mathcal{O}. \quad (\text{D.42})$$

This detector-weighted mismatch is the more relevant quantity when assessing whether two waveforms could be distinguished by a particular gravitational-wave detector.

Appendix E

Dendro Notes




Figure E.1 Schematic illustration of the nested dyadic grids used in wavelet adaptive mesh refinement. Points present on coarser grids are retained on finer grids, while the red points indicate the additional degrees of freedom introduced at each refinement level. These additional points correspond to the detail spaces W_j .

E.1 Wavelet Adaptive Mesh Refinement

Dendro employs wavelet adaptive mesh refinement (WAMR) to construct a sparse representation of the evolved fields while maintaining resolution in regions where the solution contains significant spatial structure. The basic idea is to represent the computational domain using a hierarchy of nested dyadic grids. In one spatial dimension, these grids may be written as

$$V_j = \{x^{j,k} = 2^{-j} k \Delta x\}, \quad (\text{E.1})$$

where j denotes the refinement level, k indexes the points on that level, and Δx is the spacing on the coarsest grid. The grids are nested, so points present on a coarse level also appear on all finer levels. The multidimensional AMR hierarchy used in the simulations is built from the same dyadic refinement idea.

Figure E.1 illustrates this nested structure in one dimension. The black points denote grid points inherited from coarser levels, while the red points denote the additional points introduced at each refinement level. These additional points form the detail spaces W_j , which contain the information added when passing from level $j - 1$ to level j .

Let $u^{j,k}$ denote the value of a field u at the grid point $x^{j,k}$. Since the grids are nested, values at points inherited from the previous level satisfy

$$u^{j+1,2k} = u^{j,k}. \quad (\text{E.2})$$

The values at newly introduced points may be estimated by interpolation from the previous level,

$$u^{j+1,2k+1} = \sum_m h_{j,m}^{j+1,2k+1} u^{j,m}, \quad (\text{E.3})$$

where $h_{j,m}^{j+1,2k+1}$ are the interpolation weights.

Equivalently, this interpolation can be described in terms of scaling functions. A basis function on level j may be represented using basis functions on the finer level $j+1$ as

$$\phi_{j,k}(x) = \sum_l c^l \phi_{j+1,l}(x), \quad (\text{E.4})$$

where the coefficients c^l encode the interpolation stencil. This nested basis construction allows the field to be represented by a coarse contribution together with corrections from the detail spaces. A sparse representation of the field can therefore be written schematically as

$$u(x) = \sum_{k \in S_0} u^{0,k} \phi_{0,k}(x) + \sum_{j=1}^J \sum_{k \in S_j} d^{j,k} \psi_{j,k}(x), \quad (\text{E.5})$$

where S_0 denotes the set of retained points on the coarsest level, S_j denotes the set of retained detail points on level j , $d^{j,k}$ are the wavelet coefficients, and $\psi_{j,k}$ denotes the corresponding detail basis function.

The wavelet coefficient at a detail point is computed by comparing the actual field value to the value predicted by interpolation from the previous refinement level,

$$d^{j,k} = u^{j,k} - P(x^{j,k}, j-1), \quad (\text{E.6})$$

where $P(x^{j,k}, j-1)$ denotes the interpolated value obtained from level $j-1$. Thus, the coefficient $d^{j,k}$ measures the local interpolation error associated with omitting the point $x^{j,k}$. Points for which the interpolation error is small can be discarded without significantly changing the represented field, while points with large wavelet coefficients are retained. In practice, refinement is controlled by a tolerance ε , and points are retained when, schematically,

$$|d^{j,k}| > \varepsilon. \quad (\text{E.7})$$

This criterion provides an efficient adaptive representation: smooth regions of the solution are represented using relatively few points, while regions containing sharp gradients, wave propagation, or strong-field structure are refined automatically.

E.2 Sobolev-Informed Refinement

In addition to the standard wavelet criterion, we also consider a Sobolev-inspired modification of the refinement indicator. The purpose of this modification is to make the refinement criterion sensitive not only to the local wavelet truncation estimate of a field, but also to local derivative content. This is useful in regions where the field itself may be relatively smooth in amplitude, while its gradients or curvature contain dynamically relevant structure.

For a given evolved variable u , the standard wavelet refinement indicator may be denoted by

$$I_W(u). \quad (\text{E.8})$$

This quantity represents the usual WAMR estimate based on the magnitude of the local wavelet coefficient. The Sobolev-informed indicator augments this quantity by including contributions from first and second spatial derivatives. We use an indicator of the form

$$I_{\text{Sob}}(u) = \sqrt{I_W(u)^2 + w_1 I_{\nabla u}^2 + w_2 I_{\nabla^2 u}^2}, \quad (\text{E.9})$$

where $I_{\nabla u}$ is a local first-derivative indicator, $I_{\nabla^2 u}$ is a local second-derivative indicator, and w_1 and w_2 are user-controlled weights. Any required normalization of the derivative indicators is absorbed into these weights. The first-derivative term increases refinement in regions with large gradients, while the second-derivative term increases refinement in regions with large curvature or rapidly changing gradients.

This refinement measure should be interpreted as a *Sobolev-inspired* or *Sobolev-weighted* WAMR indicator rather than as an exact discrete Sobolev norm. A true Sobolev norm is a global or




Figure E.2 Representative comparison of the adaptive mesh generated by the baseline wavelet AMR criterion and the Sobolev-informed AMR criterion. The Sobolev-informed criterion includes derivative information in the refinement indicator, which changes where grid points are allocated. The oscillatory ring corresponds to the outward-propagating dipole pulse.

integrated quantity involving a function and its derivatives. By contrast, Eq. (E.9) is a local AMR indicator designed to guide mesh refinement. Its role is to bias the adaptive mesh toward regions where derivative information suggests that additional resolution may be needed, while retaining the efficiency of the underlying wavelet-based refinement algorithm.

The following figures compare the baseline wavelet adaptive mesh refinement criterion with the Sobolev-informed refinement criterion. These diagnostics compare both the constraint behavior of the two refinement strategies and the computational cost associated with the resulting adaptive meshes.

Figure E.2 shows a representative side-by-side comparison of the adaptive mesh structure produced by the two refinement criteria. The Sobolev-informed criterion modifies the local refinement indicator by including derivative information, which changes the distribution of refined regions relative to the baseline wavelet criterion.




Figure E.3 Constraints for the baseline wavelet AMR and Sobolev-informed AMR refinement criteria. The left panel shows the L_2 norm of the electric-field divergence constraint, while the right panel shows the corresponding divergence for the vector potential. These quantities are used to assess whether the modified refinement indicator improves or degrades the constraint behavior of the evolution.

Figure E.3 shows the divergence constraints for the electric and vector-potential fields. These quantities provide a direct measure of the constraint violation during the evolution. The comparison shows how the Sobolev-informed refinement criterion affects the constraint-preserving behavior of the evolution relative to the baseline wavelet criterion.

Figure E.4 compares the number of active mesh points and a measure of the grid-efficiency. The number of active mesh points, N_{mesh} , measures the size of the adaptive grid and is therefore a proxy for the computational cost of the evolution. A refinement strategy that achieves comparable or smaller constraint violations using fewer mesh points is more efficient.

To quantify this tradeoff, we define the grid efficiency as:

$$\eta = \frac{1}{C_{\text{tot}} N_{\text{mesh}}^2}, \quad (\text{E.10})$$

where C_{tot} is the total constraint measure and N_{mesh} is the number of active mesh points. In this work, the total constraint measure is constructed from the divergence constraints as

$$C_{\text{tot}} = C_{\nabla E} + C_{\nabla A}, \quad (\text{E.11})$$




Figure E.4 Mesh size and grid-efficiency for the baseline wavelet AMR and Sobolev-informed AMR refinement criteria. The left panel shows the number of active mesh points N_{mesh} , which serves as a proxy for computational cost. The right panel shows $\eta = 1/(C_{\text{tot}} N_{\text{mesh}}^2)$. Larger values of η indicate smaller constraint violation for a given mesh size, and therefore a more efficient use of grid points.

where $C_{\nabla E}$ and $C_{\nabla A}$ denote the L_2 norms of the corresponding divergences. With this definition, larger values of η indicate that a simulation achieves smaller constraint violation for a given mesh size. The factor of N_{mesh}^2 penalizes simulations that reduce the constraint violation only by using substantially more grid points.

E.3 Performance of the Code

E.3.1 Constraint Quantities

We monitor the constraint violations throughout the evolution. In particular, we compute the L_2 norms of the Hamiltonian and momentum constraints, together with the electromagnetic constraints. When evaluating these norms, we excise small neighborhoods around the punctures to avoid allowing near-puncture behavior to dominate the constraint measure. Representative constraint violations for the full simulation are shown in Figs. E.5 and E.6.




Figure E.5 The Hamiltonian constraint $\mathcal{H}$ and the summed momentum constraint violation measured over the full simulation. The constraints remain controlled throughout the evolution, including the merger phase.




Figure E.6 The volume-weighted L_2 norms of the electromagnetic constraints measured over the full simulation. These include the magnetic no-monopole constraint, $\nabla \cdot B$, and the electric Gauss constraint, $\nabla \cdot E$.

A full convergence study for binary black hole evolutions is computationally expensive and is therefore not attempted here. Instead, we assess the behavior of the code by varying two parameters that directly affect the adaptive mesh refinement: the wavelet tolerance, ε , and the maximum refinement level, $j_{\max}$. The wavelet tolerance is the threshold value applied to the coefficients of the wavelet expansion in the WAMR scheme; it determines whether refinement or coarsening occurs. The maximum refinement level sets the finest grid spacing that can be reached in a simulation. We confirm that making these refinement criteria more stringent leads to a corresponding reduction in the constraint violations.

The tests performed here use the first $20M$ of a head-on evolution of two equal-mass, charged black holes. The black holes have an initial coordinate separation of $16M$, individual masses $m_1 = m_2 = 0.50$, and charge-to-mass ratios $q_1/m_1 = q_2/m_2 = 0.1$. Both black holes have dimensionless spin $\chi = 0.4$ and are initially at rest. In computing the constraints, we again exclude a small neighborhood of each puncture from the calculation of the norms. Throughout this section, the constraint measure shown is the sum of the volume-weighted Hamiltonian constraint and the three volume-weighted momentum constraints. This provides a single measure of the overall constraint violation during the evolution.

Comparisons of simulations with different wavelet tolerances are shown in Fig. E.7. Decreasing ε allows the grid to refine more aggressively in regions with larger truncation error. For the three wavelet tolerances $\varepsilon \in \{3 \times 10^{-5}, 1 \times 10^{-5}, 3 \times 10^{-6}\}$, the constraints decrease as ε is lowered.

Comparing simulations with different maximum refinement levels provides another way to test resolution dependence in an AMR setting. Figure E.8 compares simulations with maximum refinement levels $j_{\max} \in \{14, 15, 16, 17\}$, corresponding to finest grid spacings $h \in \{0.0325, 0.0162, 0.00813, 0.00407\}$, respectively. Increasing the allowed maximum depth provides additional resolution near the black holes and in regions where the fields develop sharp spatial




Figure E.7 The sum of the volume-weighted Hamiltonian and momentum constraint violations over the first $20M$ of a head-on evolution of two equal-mass charged black holes with charge-to-mass ratio $q/m = 0.1$ and dimensionless spin $\chi = 0.4$. The simulations use wavelet tolerances $\varepsilon \in \{3 \times 10^{-5}, 1 \times 10^{-5}, 3 \times 10^{-6}\}$ and a maximum refinement level of $j_{\max} = 15$. Lowering the wavelet tolerance improves the resolution of dynamically important regions and reduces the overall constraint violation.




Figure E.8 The sum of the volume-weighted Hamiltonian and momentum constraint violations over the first $20M$ of a head-on evolution of two equal-mass charged black holes with charge-to-mass ratio $q/m = 0.1$ and dimensionless spin $\chi = 0.4$. The simulations compare maximum refinement levels $j_{\max} \in \{14, 15, 16, 17\}$, corresponding to finest grid spacings $h \in \{0.0325, 0.0162, 0.00813, 0.00407\}$, respectively. The wavelet tolerance is fixed at $\varepsilon = 1 \times 10^{-5}$. Increasing the maximum refinement level reduces the overall constraint violation by allowing additional grid refinement in under-resolved regions.

structures. The resulting decrease in the constraints provides further evidence that the simulations improve with increased resolution.

E.3.2 Inspiral Constraint Quantities

We also monitor the constraint violations for an inspiraling binary. Figure E.9 shows the Hamiltonian and momentum constraint violations for the $q = 1$, $\chi = 0.4$, $\lambda_+^+ = 0.3$ case, while Fig. E.10 shows the




Figure E.9 The Hamiltonian constraint $\mathcal{H}$ and the summed momentum constraint violation measured over the full inspiral simulation. The L_2 norm is normalized by the volume of the grid. This representative case has $q = 1$, $\chi = 0.4$, and $\lambda_+^+ = 0.3$.

corresponding electromagnetic constraints. These indicators provide a representative measure of the constraint behavior during the inspiral and merger for a charged binary configuration.




Figure E.10 The volume-weighted L_2 norms of the electromagnetic constraints measured over the full inspiral simulation. These include the magnetic no-monopole constraint, $\nabla \cdot B$, and the electric Gauss constraint, $\nabla \cdot E$. This representative case has $q = 1$, $\chi = 0.4$, and $\lambda_+^+ = 0.3$.

Appendix F

CFD Notes

F.0.1 Stability tests

Appendix G

Running the EMDA Code on MaryLou

G.1 Building emda-gr

To build emda-gr, the following external packages are required:

- **C/C++11 or higher** (tested with GNU and Intel compilers)
- **MPI implementation** (tested with MVAPICH2, OpenMPI, and Intel MPI)
- **BLAS/LAPACK** (tested with OpenBLAS and Intel MKL)
- **GSL** (GNU Scientific Library); set `GSL_ROOT_DIR` in CMake to enable automatic detection

On MaryLou, the recommended module environment is:

```
andrewc0@login03:$ ml
```

Currently Loaded Modules:

1. gcc/14.1
2. python/3.12

3. `cmake/3.28`
4. `gcc-runtime/13.2.0-g4hc7ev`
5. `eigen/3.4.0-6kqviac`
6. `openmpi5-GR`
7. `gsl-GR`
8. `mkl-GR`
9. `rclone/1.65.2-tyg4lvv`

To ensure that these dependencies are always available, it is convenient to load them automatically in your shell configuration. Open your `~/.bash_profile`:

```
vim ~/.bash_profile
```

An example `.bash_profile` is shown below. Additional customizations may be added as needed.

```
# --- Modules ---
```

```
module purge
```

```
module use /grp/home/grp_GR/.modulefiles
```

```
module load gcc/14.1 python/3.12 cmake/3.28 eigen/3.4.0-6kqviac
```

```
module load openmpi5-GR gsl-GR mkl-GR rclone
```

```
export HCOLL_DIR=/vapps/rhel9/x86_64/HPC-X/hpcx-v2.19-gcc-inbox-redhat9-cuda12-x86_64/
```

```
export LD_LIBRARY_PATH="${HCOLL_DIR}:${LD_LIBRARY_PATH:-}"
```

```
# --- GSL ---

export GSL_DIR=/grphome/grp_GR/apps/gsl-2.8
export GSL_ROOT_DIR=/grphome/grp_GR/apps/gsl-2.8
export LD_LIBRARY_PATH="${GSL_DIR}/lib:${LD_LIBRARY_PATH:-}"

# --- MKL ---

if [ -n "${MKLROOT:-}" ] && [ -d "${MKLROOT}/lib/intel64" ]; then
export LD_LIBRARY_PATH="${MKLROOT}/lib/intel64:${LD_LIBRARY_PATH:-}"
fi

# --- GCC 14 runtime: libstdc++ and libgcc_s ---

# Required to avoid GLIBCXX and CXXABI runtime errors.

if command -v g++ >/dev/null 2>&1; then
GCC_LIBDIR="$(dirname "$(g++ -print-file-name=libstdc++.so.6)")"
GCC_GCCLIBDIR="$(dirname "$(g++ -print-file-name=libgcc_s.so.1)")"

if [ -d "$GCC_LIBDIR" ]; then
export LD_LIBRARY_PATH="$GCC_LIBDIR:${LD_LIBRARY_PATH:-}"
fi

if [ -d "$GCC_GCCLIBDIR" ]; then
export LD_LIBRARY_PATH="$GCC_GCCLIBDIR:${LD_LIBRARY_PATH:-}"
```

```
fi
fi

# --- History settings ---

HISTSIZE=1000000
HISTFILESIZE=2000000
HISTCONTROL=ignoredups:erasedups
HISTIGNORE="ls:bg:fg:history"
shopt -s histappend

# Save history after every command.

PROMPT_COMMAND="history -a; history -c; history -r; $PROMPT_COMMAND"
```

After reloading the shell, verify that the modules have been loaded correctly:

```
ml
```

The output should match the module list shown above.

G.2 Building the Solver

First, clone the EMDA repository. Create a project directory if desired, then run:

```
git clone [git@github.com](mailto:git@github.com):dfvankomen/emda-gr.git
cd emda-gr
```

CMake will automatically fetch `dendrolib` during configuration.

Create and enter a build directory:

```
mkdir build
```

```
cd build
```

Configure the project using:

```
ccmake ..
```

CMake will locate the required libraries and open an interactive configuration interface. The main settings that may need to be adjusted are the system of equations and the target CPU architecture.

The available equation systems include EMDA, EMD, and EMGR. For a first test run, choose the EMDA system.

The CPU architecture should match the compute nodes on which the job will run. On MaryLou, the `physics2` and `m12` partitions use 64-core AMD EPYC 7763 processors. These are Zen 3 CPUs, so set

```
CPU_ARCH = znver3
```

After configuration is complete, generate the build files from within `ccmake`. Then compile the code using:

```
make -j
```

G.2.1 Build Outputs

Once the build completes, the `emdaSolver` and `tpid` binaries should be located in the solver build directory, for example:

```
emda-gr/build/emdaSolver/
```

At this point, the solver is ready to be used for simulations.

G.3 Runs and Job Scripts

After building the solver, move from the home directory to the scratch directory. The location of the scratch directory depends on the supercomputer. On MaryLou, the scratch area is located under `/nobackup/autodelete/usr/`.

First, change into your scratch area:

```
cd /nobackup/autodelete/usr/YOURUSERNAME/
```

Create a directory for the EMDA runs and enter it:

```
mkdir EMDA_runs
```

```
cd EMDA_runs
```

Next, copy the solver binaries and a parameter file into the run directory. In the commands below, replace `~/path/to/` with the location of your local `emda-gr` checkout.

Copy `emdaSolver`:

```
cp ~/path/to/emda-gr/build/emdaSolver/emdaSolver ./
```

Copy `tpid`:

```
cp ~/path/to/emda-gr/build/emdaSolver/tpid ./
```

Copy the parameter file:

```
cp ~/path/to/emda-gr/emda_simplified.param.toml ./
```

Verify that everything copied correctly:

```
ls
```


You should see:

```
emda_simplified.param.toml  emdaSolver  tpid
```

Now create the standard output directories for the run:

```
mkdir cp prof vtu bah
```

These directories store the main simulation outputs:

- **cp**: checkpoint files, which allow the run to be restarted
- **prof**: waveform extraction, ADM quantities, charges, and constraint outputs
- **vtu**: visualization files that can be opened in ParaView
- **bah**: apparent-horizon data

A later section will describe the most important parameter-file settings needed to control these outputs.

G.3.1 Submitting a Run with Slurm

To run the simulation on the cluster, create a Slurm job script. For example:

```
vim Job.sb
```

Paste the following into the file:

```
#!/bin/bash
#SBATCH --time=72:00:00
#SBATCH --ntasks=256
#SBATCH --nodes=2
```

```
#SBATCH --mem-per-cpu=4096M
#SBATCH -J EMDA_RUN
#SBATCH --mail-user=[youremail@email.com] (mailto:youremail@email.com)
#SBATCH --mail-type=END,FAIL
#SBATCH --partition=m12

echo "Starting run"

source ~/.bash_profile

./tpid emda_simplified.param.toml 256 &> TPID.txt
mpirun -np 256 ./emdaSolver emda_simplified.param.toml &>> Inspiral.txt

echo "Run completed"
```

The `tpid` command generates the initial data, while `emdaSolver` evolves that data forward in time. Since these commands are written on separate lines and the `tpid` command is not placed in the background, the script waits for `tpid` to finish before starting `emdaSolver`. This ordering is necessary because `tpid` produces the binary initial-data file, `rit_q2_tpid_sol.bin`, which is then read by `emdaSolver`.

On the `m12` partition, each node has 128 CPU cores. Therefore, a run using 256 MPI tasks typically uses 2 nodes. If the number of MPI tasks is changed, the number of requested nodes should be adjusted accordingly.

Submit the job with:

```
sbatch Job.sb
```

Once submitted, the job will enter the queue and begin running when resources become available. You can check the status of your jobs with:

```
squeue | grep yourusername
```

After the job starts, you can monitor the output by inspecting the text files generated during execution. For example:

```
cat TPID.txt
```

or

```
tail -f Inspiral.txt
```

At this point, the simulation is running.

G.4 Parameter-File Explanation

The parameter file is read by both `tpid` and `emdaSolver`. It allows many run settings to be changed without rebuilding the code, which makes it easy to adjust output frequencies, grid settings, checkpointing behavior, and initial-data parameters.

This section summarizes some of the most important parameters. Parameters not discussed here can usually be left at their default values.

```
# Whether or not the solver should attempt to restore from a checkpoint
```

```
EMDA_RESTORE_SOLVER = false
```

This option determines whether `emdaSolver` should attempt to restart from an existing checkpoint. For a fresh run, this should be set to `false`. To continue a previous run from checkpoint data, set this to `true`.

`EMDA_IO_OUTPUT_FREQ = 400`

This controls how often VTU output is written. Smaller values produce output more frequently, but they also increase storage usage. This parameter should be chosen based on how much visualization data is needed.

`EMDA_REMESH_TEST_FREQ = 200`

This sets how often the solver invokes the adaptive mesh refinement algorithm to determine whether the grid should be refined or coarsened. If this value is too small, remeshing can occur too frequently and may introduce unnecessary grid noise. If it is too large, the grid may not adapt quickly enough to resolve the dynamics. A value around 200 has worked well in typical EMDA runs.

`EMDA_GW_EXTRACT_FREQ = 25`

This controls how often gravitational-wave quantities are extracted. The same output step also includes other useful quantities, such as black-hole positions and masses. This output is not especially expensive compared with VTU output, so values in the range 25–100 are usually reasonable.

`EMDA_CHECKPT_FREQ = 100`

This specifies the number of time steps between checkpoints. Checkpointing is relatively inexpensive compared with full visualization output, so this value can be kept fairly small. More frequent checkpoints make it easier to restart a run after a crash or wall-time limit.

```
# VTU file prefix
```

```
EMDA_VTU_FILE_PREFIX = "vtu/emda_gr"
```

```
# Checkpoint file prefix
```

```
EMDA_CHKPT_FILE_PREFIX = "cp/emda_cp"
```

```
# Profiling file prefix
```

```
EMDA_PROFILE_FILE_PREFIX = "prof/emda_prof"
```

These options control where different types of output data are written. The `vtu` directory stores visualization output, the `cp` directory stores checkpoint files, and the `prof` directory stores extracted waveforms and other reduced data products.

```
# Which evolution variables to output
```

```
EMDA_VTU_OUTPUT_EVOL_INDICES = [0, 1, 2, 3, 4, 5, 6, 7, 8, 9,  
10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,  
25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35]
```

```
# Number of evolution variables to output
```

```
EMDA_NUM_EVOL_VARS_VTU_OUTPUT = 36
```

These settings determine which evolution variables are written to the VTU files. These variables include quantities such as the metric components, extrinsic-curvature components, gauge variables, and matter fields. If storage usage becomes a concern, some variables may be omitted from the VTU output.

```
# Which constraint variables to output
```

```
EMDA_VTU_OUTPUT_CONST_INDICES = [0, 1, 2, 3, 4, 5, 6, 7,  
8, 9, 10, 11, 12, 13]
```

```
# Number of constraint variables to output
```

```
EMDA_NUM_CONST_VARS_VTU_OUTPUT = 14
```

These options are analogous to the evolution-variable output settings, but they apply to constraint quantities. These include quantities such as the Hamiltonian constraint, the momentum constraints, and additional constraints associated with the electromagnetic sector.

```
# Whether to isolate an X slice
```

```
EMDA_VTU_X_SLICE = false
```

```
# Whether to isolate a Y slice
```

```
EMDA_VTU_Y_SLICE = false
```

```
# Whether to isolate a Z slice
```

```
EMDA_VTU_Z_SLICE = true
```

```
# Any combination may be set to true
```

Writing VTU output for the full 3-D grid can be expensive, so it is often preferable to output a 2-D slice instead. For most visualization purposes, a single slice is sufficient. In typical runs, the Z slice is a good default choice.

Although multiple slice options can be enabled at the same time, doing so may produce output that is less straightforward to interpret. For simple visualization, enable only one slice direction at a time.

```
EMDA_DENDRO_GRAIN_SZ = 50
```

This controls the approximate number of grid blocks assigned to each processor. In practice, this parameter should usually be kept between 25 and 200. Very small or very large values may reduce parallel efficiency.

```
EMDA_DENDRO_AMR_FAC = 0.2
```

This controls the sensitivity of the adaptive mesh refinement algorithm. Smaller values generally make the grid more sensitive to refinement, while larger values make refinement less aggressive. This parameter should be adjusted with care, since it affects both accuracy and computational cost.

```
EMDA_ELE_ORDER = 6
```

This sets the finite-element order used by the Dendro infrastructure. Loosely speaking, it controls how much information each block must exchange with its neighbors. Higher values increase communication costs and may slow the code. For the current EMDA implementation, this should generally be left at 6.

```
# Minimum depth of the mesh/octree
```

```
EMDA_MINDEPTH = 4
```

```
# Maximum depth of the mesh/octree
```

```
EMDA_MAXDEPTH = 15
```

These parameters control the minimum and maximum refinement levels of the octree mesh. The minimum depth sets the coarsest allowed grid resolution, while the maximum depth sets the finest allowed grid resolution. Increasing `EMDA_MAXDEPTH` improves the maximum available resolution, but it also increases memory usage and computational cost.

The CFL condition is set by the finest level of the grid, so increasing `EMDA_MAXDEPTH` can significantly increase the runtime. Each increase in `EMDA_MAXDEPTH` halves the grid spacing on the finest level. As a rule of thumb, each black hole should be resolved by at least roughly 50 points across the horizon. For an equal-mass binary with total mass $M = 1$ on a ± 400 grid, `EMDA_MAXDEPTH = 15` is usually adequate. For larger mass ratios, the smaller black hole requires additional refinement. For example, a $q = 2$ binary typically requires `EMDA_MAXDEPTH = 16`.

```
#####
```



```
# REFINEMENT STRATEGY
```

```
# Which refinement mode to use
```

```
# 0 - WAMR
```

```
# 1 - EH
```

```
# 2 - EH_WAMR
```

```
# 3 - BH_LOC
```

```
# 4 - BH_WAMR
```

```
EMDA_REFINEMENT_MODE = 4
```

This parameter controls the refinement strategy used by the code. The recommended choice is currently `EMDA_REFINEMENT_MODE = 4`, which uses black-hole-centered refinement together with wavelet adaptive mesh refinement.

```
EMDA_AMR_R_RATIO = 1.618033988749895
```

This parameter determines how quickly the forced refinement around each black hole falls off with radius. The default value is reasonable and should usually be left unchanged for initial runs.

```
# Wavelet Tolerance Function to use (see grUtils.cpp for adding more)
```

```
# 1,2 - deprecated

# 3 - center highly refined; low refinement elsewhere eases over time

# 4 - start with refinement centered at BHs, then widen to  $r=0$  focused

# 5 - focus refinement in self-similar ways about each BH

# 6 - refine well in  $r \leq t$  and coarsely in  $r > t$ 

EMDA_USE_WAVELET_TOL_FUNCTION = 6
```

This parameter controls how the wavelet refinement tolerance varies across the computational domain. Function 6, developed by William, has worked well in practice and is the recommended choice for current EMDA runs. More details can be found in Ref. [27].

```
# Minimum wavelet tolerance

EMDA_WAVELET_TOL = 0.00001

# Gravitational wave refinement tolerance

EMDA_GW_REFINE_WTOL = 0.0001

# Maximum wavelet tolerance
```

```
EMDA_WAVELET_TOL_MAX = 0.001
```

```
# Wavelet tolerance radii (for shell-based tolerance)
```

```
EMDA_WAVELET_TOL_FUNCTION_R0 = 3.0
```

```
EMDA_WAVELET_TOL_FUNCTION_R1 = 12.0
```

The minimum wavelet tolerance, `EMDA_WAVELET_TOL`, is the most important of these parameters because it sets the strictest refinement criterion used by the grid. Smaller values force more refinement and generally improve accuracy, but they also increase memory usage and runtime. The remaining parameters depend on the chosen wavelet tolerance function. The default values are reasonable starting points, but they may be adjusted for higher-resolution runs or for runs in which the wave-extraction region requires additional refinement.

```
# Indices of variables used for refinement (see grDef.h for definitions)
```

```
EMDA_REFINE_VARIABLE_INDICES = [0, 1, 2, 3, 4, 5, 6, 7, 8, 9,  
10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,  
25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35]
```

```
# Number of variables used for refinement
```

```
EMDA_NUM_REFINE_VARS = 36
```

These parameters specify which evolution variables are used when deciding whether the grid should be refined. It is not always necessary to use every evolution variable as a refinement variable, but the full list is a safe default for general-purpose EMDA runs.

```
# CFL (Courant--Friedrichs--Lewy) factor
```

```
EMDA_CFL_FACTOR = 0.25
```

```
# Simulation start time
```

```
EMDA_RK_TIME_BEGIN = 0
```

```
# Simulation end time
```

```
EMDA_RK_TIME_END = 1000
```

These parameters control the CFL factor and the total coordinate time of the simulation. The CFL factor sets the time step relative to the grid spacing on the finest refinement level. The parameters EMDA_RK_TIME_BEGIN and EMDA_RK_TIME_END determine the start and end times of the evolution. The remaining Runge–Kutta settings can usually be left at their default values.

```
#####
```

```
# EMDA THEORY PARAMETERS
```

```
#####
```

```
EMDA_LAMBDA = [1, 1, 1, 1]
```

```
EMDA_ALPHA_THEORY = [1.0, 1.0]
```

```
EMDA_LF = [1.0, 0.0]
```

```
EMDA_P_EXPO = -1.0
```

```
EMDA_ALPHA_POS_NEG_FLOOR = false
```

```
EMDA_ETA = [0.4, 0.4]
```

```
EMDA_XI = [0, 0, 0]
```

```
CHI_FLOOR = 0.00001
```

```
EMDA_TRK0 = 0.0
```

```
EMDA_ETA_R0 = 1.31
```

```
EMDA_ETA_POWER = [2.0, 2.0]
```

```
EMDA_ETADAMP = 1.0
```

```
ETA_CONST = 2.0
```

```
ETA_R0 = 30.0
```

```
ETA_DAMPING = 1.0
```

```
ETA_DAMPING_EXP = 1.0
```

These parameters define the theory and gauge choices used in the evolution. The main parameters that are typically changed are the two theory couplings α_0 and α_1 , stored in EMDA_ALPHA_THEORY. The choices

$$(\alpha_0, \alpha_1) = (1, 1)$$

correspond to EMDA,

$$(\alpha_0, \alpha_1) = (1, 0)$$

correspond to EMD, and

$$(\alpha_0, \alpha_1) = (0, 0)$$

correspond to Einstein–Maxwell. The remaining gauge parameters are generally safe to leave at their default values.

```
# Kreiss--Oliger dissipation parameter
```

```
KO_DISS_SIGMA = 0.4
```

This parameter controls the strength of Kreiss–Oliger numerical dissipation. Some dissipation is needed for stability, and `KO_DISS_SIGMA = 0.4` is a reasonable starting value.

```
#####
```

```
# Select initial data type
```

```
# See the bottom of the file for reference
```

```
EMDA_ID_TYPE = 0
```

This parameter selects the type of initial data used by the evolution. The choice `EMDA_ID_TYPE = 0` uses the TPID solver. Other black-hole initial-data options are available, but TPID is the standard choice for the binary black-hole simulations described here.

```
#####
```

```
# Set up EMDA grid extents
```

```
EMDA_GRID_MIN_X = -400.0
```

```

EMDA_GRID_MAX_X = 400.0
EMDA_GRID_MIN_Y = -400.0
EMDA_GRID_MAX_Y = 400.0
EMDA_GRID_MIN_Z = -400.0
EMDA_GRID_MAX_Z = 400.0

```

These parameters define the physical extent of the computational domain. The values shown here are reasonable defaults for many binary simulations. The outer boundary should be placed far enough from the wave-extraction region that boundary reflections do not contaminate the extracted waveforms during the time interval of interest.

```
#####
```

```
# Black hole parameters
```

```

EMDA_BH1_AMR_R = 1.0
EMDA_BH1_CONSTRAINT_R = 2.0
EMDA_BH1_MAX_LEV = 15
EMDA_BH1 = {MASS = 0.5, X = 5.0, Y = 0.0, Z = 0.0,
V_X = -0.00100282, V_Y = 0.095775311, V_Z = 0.0,
SPIN = 0.0, SPIN_THETA = 0.0, SPIN_PHI = 0.0,
ELECTRIC_CHARGE = 0.05, MAGNETIC_CHARGE = 0.0}

```

```

EMDA_BH2_AMR_R = 1.0
EMDA_BH2_CONSTRAINT_R = 2.0
EMDA_BH2_MAX_LEV = 15
EMDA_BH2 = {MASS = 0.5, X = -5.0, Y = 0.0, Z = 0.0,

```

```
V_X = 0.00100282, V_Y = -0.095775311, V_Z = 0.0,
SPIN = 0.0, SPIN_THETA = 0.0, SPIN_PHI = 0.0,
ELECTRIC_CHARGE = 0.05, MAGNETIC_CHARGE = 0.0}
```

These parameters define the black-hole properties used by the evolution and by the refinement infrastructure. The parameter `EMDA_BH1_AMR_R` sets the radius within which refinement is forced around the first black hole, while `EMDA_BH1_CONSTRAINT_R` sets the radius excluded when computing constraint norms. The corresponding BH2 parameters play the same role for the second black hole. The parameters `EMDA_BH1_MAX_LEV` and `EMDA_BH2_MAX_LEV` set the maximum refinement level allowed near each black hole.

The X, Y, and Z entries do not directly determine the black holes' initial positions in the TPID solve. Instead, they specify where the evolution code begins searching for the black-hole locations and where it centers the black-hole refinement regions. For equal-mass binaries, these values should match the corresponding $\pm b$ positions used in the TPID parameters.

The electric charge should remain below the black-hole mass in order to avoid exceeding the extremal limit. Magnetic charge is still under development and should not be used for standard production runs. The spin parameter corresponds to the dimensionless spin χ , with angular momentum $S = \chi m^2$.

TPID reads the charge and spin from these black-hole parameter blocks, but it does not use the listed mass or position values in the same way as the evolution code. The TPID mass and separation are set separately by the TPID parameters described below.

Finally, note that `V_X`, `V_Y`, and `V_Z` represent the black-hole momenta, not their coordinate velocities. This distinction is important when setting up binary initial data.

```
#####
```

```
# GRAVITATIONAL WAVE EXTRACTION PARAMETERS
```



```
EMDA_GW_NUM_RADAI = 6
EMDA_GW_RADAI = [50.0, 60.0, 70.0, 80.0, 90.0, 100.0]
```

```
EMDA_GW_NUM_LMODES = 7
EMDA_GW_L_MODES = [2, 3, 4, 5, 6, 7, 8]
```

```
# Radius at which conserved quantities are extracted
```

```
EMDA_GW_EXTRACTION_RADIUS = 300.0
```

```
# Modes for EM, dilaton, and axion radiation
```

```
EMDA_RAD_NUM_LMODES = 6
EMDA_RAD_L_MODES = [1, 2, 3, 4, 5, 6]
```

```
#####
```

These parameters control the extraction of gravitational, electromagnetic, dilaton, and axion radiation. Radiation is extracted on spherical surfaces at the radii listed in `EMDA_GW_RADAI`. The gravitational-wave modes are set by `EMDA_GW_L_MODES`, while the modes for the electromagnetic, dilaton, and axion fields are set by `EMDA_RAD_L_MODES`.

The mode numbers correspond to the spherical-harmonic multipoles being extracted. Although modes can be extracted up to roughly $\ell \approx 10$, the dominant contributions are usually contained in the lower modes, especially $\ell \leq 4$.

```
# Time steps between apparent-horizon (AEH) solves
AEH_SOLVER_FREQ = 50
```

This parameter controls how often the apparent-horizon solver is called. The remaining AEH-related settings should generally be left at their default values unless the apparent-horizon finder fails or additional horizon information is needed.

TPID_PAR_B = 5.0

TPID_TARGET_M_PLUS = 0.48594

TPID_TARGET_M_MINUS = 0.48594

The parameter TPID_PAR_B sets the coordinate half-separation used by TPID. For example, TPID_PAR_B = 5.0 corresponds to punctures located near $x = \pm 5$. The target masses are the values used internally by TPID when constructing the initial data.

TPID_CENTER_OFFSET = {X = 0.0, Y = 0.0, Z = 0.0}

For equal-mass binaries, this should usually be left at zero. For unequal-mass binaries, the system may need to be recentered so that the center of mass is located at the origin.

TPID_NPOINTS_A = 40

TPID_NPOINTS_B = 40

TPID_NPOINTS_PHI = 20

These parameters control the resolution of the TPID grid. The values shown here are reasonable defaults for a $q=1$ black hole, but higher values may be needed for more accurate initial data and higher mass ratios.

TPID_GIVE_BARE_MASS = 1

This parameter determines whether the puncture masses are fixed or solved for. When TPID_GIVE_BARE_MASS = 0, TPID solves for the bare masses. When TPID_GIVE_BARE_MASS = 1, TPID uses the specified target masses.

A useful procedure is to first estimate the desired masses, run TPID with mass solving enabled, then update the target masses and rerun with `TPID_GIVE_BARE_MASS = 1` to finalize the initial data.

```
TPID_REPLACE_LAPSE_WITH_SQRT_CHI = true
```

This option should be enabled when using the slow-start lapse.

```
# Include dilatonic contributions in rho during TPID solve
```

```
TPID_USE_DILATON = true
```

```
# Interpolated initial data should not be used with TPID_USE_DILATON
```

```
# You should also be using the EMD equations in this case
```

```
TPID_USE_INTERPOLATED_SOLUTION = false
```

These parameters control how scalar-field contributions are included in the initial-data solve. If both options are set to `false`, TPID generates Einstein–Maxwell initial data. Setting `TPID_USE_INTERPOLATED_SOLUTION = true` produces EMD-type initial data. Setting `TPID_USE_DILATON = true` includes dilatonic contributions directly in the TPID solve.

The initial-data choice should be consistent with both the theory parameters in `EMDA_ALPHA_THEORY` and the system of equations selected when the code was configured with CMake.

Appendix H

ParaView

This chapter provides a brief introduction to several basic ParaView workflows used to inspect Dendro simulation output. We begin by loading simulation files, then describe how to create slices, visualize fields, use the Calculator filter, adjust color maps, export data, and locate regions of interest.

H.1 Opening Simulation Data

To load data into ParaView, click the **Open File** button in the top-left toolbar. This button is highlighted in Fig. H.1.

After clicking the **Open File** button, navigate to the directory where the simulation output is saved. Dendro commonly writes visualization data in `.vtu` format, which stands for *VTK Unstructured Grid*. For parallel runs, the output is often accompanied by a `.pvtu` file. The `.pvtu` file acts as a wrapper that tells ParaView how the individual `.vtu` pieces fit together.

In most cases, one should open the `.pvtu` file rather than opening the individual `.vtu` files by hand. This allows ParaView to load the full parallel dataset as a single object.




Figure H.1 The ParaView interface with the **Open File** icon highlighted.

H.2 Creating a Slice

Once the data has been loaded, a common operation is to create a two-dimensional slice through the three-dimensional dataset. In this example, the **Slice** tool is used to create a slice through the data, as shown in Fig. H.2.




Figure H.2 The ParaView interface with the **Slice** and **Calculator** tools highlighted.

After selecting the **Slice** tool, ParaView displays the slice settings in the Properties panel. These settings determine the location and orientation of the slicing plane.

H.3 Adjusting the Slice and Viewing Data

ParaView allows the user to choose how the data should be sliced. Often, we want to create z -slices, since this is the default orientation used for many Dendro visualizations. To do this, set the slice normal to $(0, 0, 1)$. We also set the origin to $(0, 0, 10^{-5})$. The small nonzero z -value is used because setting the slice exactly at $z = 0$ can sometimes cause ParaView to miss cells lying on the symmetry plane.

The **Show Plane** option should be turned off so that only the data slice is displayed. After changing these settings, click **Apply**.

Once the slice is displayed, the visualized field can be changed using the field-selection menu. The available time steps can be explored using the time controls, and the representation menu can be used to switch between viewing modes such as **Surface** and **Wireframe**. These controls are highlighted in Fig. H.3.

H.4 Using the Calculator

The **Calculator** filter can be used to construct derived quantities from the fields stored in the simulation output. After selecting the loaded dataset or slice, click the **Calculator** button. In the Calculator panel, choose the appropriate scalar or vector fields from the available data arrays. Mathematical operations can then be applied to these fields to define a new quantity for visualization or export.




Figure H.3 ParaView controls for stepping through time, selecting the displayed field or color option, and switching between representations such as **Surface** and **Wireframe**.

For example, the Calculator can be used to rescale a field, combine multiple fields, or construct diagnostic quantities that are not stored directly in the output files. After entering the desired expression, click **Apply** to create the new field.

H.5 Warping by Scalar

The **Warp By Scalar** filter can be used to displace a surface according to the value of a scalar field. This is useful for visualizing the relative amplitude of a field on a slice.

To use the filter, click **Filters** and search for **Warp By Scalar**. After selecting the filter, its settings appear in the Properties panel, in the same general location as the slice settings.

In the **Scalars** menu, select the field that should determine the warp. The **Scale Factor** controls the strength of the displacement. Large scale factors can make small-amplitude fields easier to see, but they can also distort the visualization if chosen too large.




Figure H.4 The **Warp By Scalar** settings in the Properties panel. The **Scalars** menu selects the field used for the warping, and the **Scale Factor** controls the strength of the warp.

H.6 Using the Color Map Editor

The **Color Map Editor** controls how ParaView maps data values to colors. It can be used to change the color scale, adjust the visible data range, modify opacity, and choose a different color preset.

By default, the color mapping may not automatically update when stepping through time. The easiest way to ensure that the colors match the data range at each time step is to rescale the color range when the time step changes. This prevents the visualization from using an outdated color range from a previous time step.

H.7 Macros, Animations, CSV Output, and Finding Data

Often, we want to repeat the same visualization or post-processing steps for many datasets. ParaView can automate these repetitive tasks using macros. To create a macro, click **Tools** and select **Start Trace**. Then perform the sequence of actions that should be automated. When finished, click **Stop**




Figure H.5 The **Color Map Editor** panel in ParaView. This panel controls the color scale, opacity, and transfer function used to visualize the selected field.

Trace. ParaView will then allow the recorded trace to be saved as a Python script or macro, which can later be reused.

ParaView can also save the currently loaded data as a CSV file. To do this, click **File** and select **Save Data**. This saves the data from the currently active object, such as the current slice. The resulting CSV file can then be loaded into Python, MATLAB, Mathematica, or another analysis tool.

Animations can be saved by clicking **File** and selecting **Save Animation**. This is useful for producing movies that show the time evolution of a field.

Finally, ParaView can be used to find data that satisfies certain conditions. To do this, click **View** and select **Find Data**. In the **Find Data** panel, choose the dataset to search. The element type can usually be left as **Point**. Then choose the field to search, set the desired condition and value, and click **Find Data**. The matching region will be highlighted in the render view.

Appendix I

Waveform Catalog

I.1 Gravitational Waveform Catalog

The following figures show the extracted $r\Psi_4^{(2,2)}$ waveforms at $r \simeq 100M$. For each mass ratio and spin, the neutral case is shown above the charged configurations. The charged cases are ordered by charge magnitude and charge configuration.

I.2 Φ_1 Waveform Catalog

The following figures show the extracted Φ_1 waveforms at $r \simeq 100M$. Since there is no radiation in the neutral case, only charged configurations are shown. The charged cases are ordered by charge magnitude and charge configuration. For the $+-$ cases, we show the $(\ell, m) = (1, 1)$ mode, while for the $+0$ and $++$ cases we show the $(\ell, m) = (2, 2)$ mode.

I.3 Φ_2 Waveform Catalog

The following figures show the extracted Φ_2 waveforms at $r \simeq 100M$. Since there is no radiation in the neutral case, only charged configurations are shown. The charged cases are ordered by charge magnitude and charge configuration. For the $+-$ cases, we show the $(\ell, m) = (1, 1)$ mode, while for the $+0$ and $++$ cases we show the $(\ell, m) = (2, 2)$ mode.




Figure I.1 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.




Figure I.2 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.




Figure I.3 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 1$ and $\chi = 0.8$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.




Figure I.4 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 2$ and $\chi = 0$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.




Figure I.5 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 2$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations ordered by increasing charge magnitude and charge configuration.




Figure I.6 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 4$ and $\chi = 0.4$. The neutral case is shown at the top, followed by charged configurations with $|\lambda_i| = 0.05$ and $|\lambda_i| = 0.10$.




Figure I.7 Extracted $r\Psi_4^{(2,2)}$ waveforms for $q = 8$ and $\chi = 0.2$.




Figure I.8 Extracted Φ_1 waveforms for $q=1$ and $\chi=0$.





Figure I.9 Extracted Φ_1 waveforms for $q = 1$ and $\chi = 0.4$.

**Figure I.10** Extracted Φ_1 waveforms for $q = 1$ and $\chi = 0.8$.




Figure I.11 Extracted Φ_1 waveforms for $q = 2$ and $\chi = 0$.




Figure I.12 Extracted Φ_1 waveforms for $q = 2$ and $\chi = 0.4$.




Figure I.13 Extracted Φ_1 waveforms for $q = 4$ and $\chi = 0.4$.









Figure I.14 Extracted Φ_1 waveforms for $q = 8$ and $\chi = 0.2$.

**Figure I.15** Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0$.

**Figure I.16** Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0.4$.

Figure I.17 Extracted Φ_2 waveforms for $q = 1$ and $\chi = 0.8$.

**Figure I.18** Extracted Φ_2 waveforms for $q = 2$ and $\chi = 0$.**Figure I.19** Extracted Φ_2 waveforms for $q = 2$ and $\chi = 0.4$.




Figure I.20 Extracted Φ_2 waveforms for $q = 4$ and $\chi = 0.4$.




Figure I.21 Extracted Φ_2 waveforms for $q = 8$ and $\chi = 0.2$.

Bibliography

- [1] A. Einstein, “Die Feldgleichungen der Gravitation,” Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin) pp. 844–847 (1915).
- [2] C. M. Will, “The Confrontation between General Relativity and Experiment,” *Living Reviews in Relativity* **17**, 4 (2014).
- [3] N. Yunes, X. Siemens, and K. Yagi, “Gravitational-wave tests of general relativity with ground-based detectors and pulsar-timing arrays,” *Living Reviews in Relativity* **28**, 3 (2025).
- [4] N. Ashby, “Relativity in the Global Positioning System,” *Living Reviews in Relativity* **6**, 1 (2003).
- [5] S. M. Carroll, *Spacetime and Geometry: An Introduction to General Relativity* (Addison-Wesley, 2004).
- [6] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, 1973).
- [7] A. Einstein, “Näherungsweise Integration der Feldgleichungen der Gravitation,” Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin) pp. 688–696 (1916).
- [8] B. P. Abbott *et al.*, “Observation of Gravitational Waves from a Binary Black Hole Merger,” *Phys. Rev. Lett.* **116**, 061102 (2016).

- [9] LIGO Laboratory, “LIGO Facts,” <https://www.ligo.caltech.edu/page/facts>, n.d., accessed: 2026-04-27.
- [10] A. G. Abac *et al.*, “Black Hole Spectroscopy and Tests of General Relativity with GW250114,” *Phys. Rev. Lett.* **136**, 041403 (2026).
- [11] J. Veitch *et al.*, “Robust parameter estimation for compact binaries with ground-based gravitational-wave observations using the LALInference software library,” *Phys. Rev. D* **91**, 042003 (2015).
- [12] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Pürrer, X. J. Forteza, and A. Bohé, “Frequency-domain gravitational waves from nonprecessing black-hole binaries. II. A phenomenological model for the advanced detector era,” *Phys. Rev. D* **93**, 044007 (2016).
- [13] F. Löffler *et al.*, “The Einstein Toolkit: A Community Computational Infrastructure for Relativistic Astrophysics,” *Classical and Quantum Gravity* **29**, 115001 (2012).
- [14] M. Boyle *et al.*, “The SXS Collaboration catalog of binary black hole simulations,” *Classical and Quantum Gravity* **36**, 195006 (2019).
- [15] B. Brügmann, J. A. González, M. Hannam, S. Husa, U. Sperhake, and W. Tichy, “Calibration of Moving Puncture Simulations,” *Physical Review D* **77**, 024027 (2008).
- [16] K. Clough, E. A. Lim, D. G. Figueroa, and P. G. Ferreira, “GRChombo: Numerical Relativity with Adaptive Mesh Refinement,” *Classical and Quantum Gravity* **32**, 245011 (2015).
- [17] B. Daszuta, F. Zappa, W. Cook, D. Radice, S. Bernuzzi, and V. Morozova, “GR-Athena++: puncture evolutions on vertex-centered oct-tree AMR,” *Astrophys. J. Suppl.* **257**, 25 (2021).

- [18] M. Fernando, D. Neilsen, Y. Zlochower, E. W. Hirschmann, and H. Sundar, “Massively parallel simulations of binary black holes with adaptive wavelet multiresolution,” *Phys. Rev. D* **107**, 064035 (2023).
- [19] T. W. Baumgarte and S. L. Shapiro, *Numerical Relativity: Solving Einstein’s Equations on the Computer* (Cambridge University Press, 2010).
- [20] T. W. Baumgarte and S. L. Shapiro, *Numerical Relativity: Starting from Scratch* (Cambridge University Press, 2021).
- [21] M. Fernando, D. Neilsen, E. Hirschmann, and H. Sundar, “A Scalable Framework for Adaptive Computational General Relativity on Heterogeneous Clusters,” In *Proceedings of the ACM International Conference on Supercomputing*, ICS ’19 (Association for Computing Machinery, New York, NY, USA, 2019).
- [22] M. Fernando, D. Neilsen, H. Lim, E. Hirschmann, and H. Sundar, “Massively Parallel Simulations of Binary Black Hole Intermediate-Mass-Ratio Inspirals,” *SIAM Journal on Scientific Computing* **41**, C97–C138 (2019).
- [23] B. P. Abbott *et al.*, “GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs,” *Phys. Rev. X* **9**, 031040 (2019).
- [24] R. Abbott *et al.*, “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo during the Second Part of the Third Observing Run,” *Phys. Rev. X* **13**, 041039 (2023).
- [25] A. G. Abac *et al.*, “GWTC-4.0: Updating the Gravitational-Wave Transient Catalog with Observations from the First Part of the Fourth LIGO-Virgo-KAGRA Observing Run,” 2025.
- [26] M. Pürrer and C.-J. Haster, “Gravitational waveform accuracy requirements for future ground-based detectors,” *Physical Review Research* **2**, 023151 (2020).

- [27] W. K. Black, D. Neilsen, E. W. Hirschmann, D. F. Van Komen, and M. Fernando, “Nyquist-resolving gravitational waves via orbital frequency-based refinement,” *Phys. Rev. D* **111**, 124001 (2025).
- [28] S. K. Lele, “Compact Finite Difference Schemes with Spectral-like Resolution,” *Journal of Computational Physics* **103**, 16–42 (1992).
- [29] B. Daszuta, “Spectrally-tuned compact finite-difference schemes with domain decomposition and applications to numerical relativity,” *Phys. Rev. D* **107**, 104036 (2023).
- [30] B. P. Abbott, others (LIGO Scientific Collaboration, and V. Collaboration), “Tests of General Relativity with GW150914,” *Phys. Rev. Lett.* **116**, 221101 (2016).
- [31] R. Abbott, others (LIGO Scientific Collaboration, V. Collaboration, and K. Collaboration), “Tests of General Relativity with GWTC-3,” *Physical Review D* **112**, 084080 (2025).
- [32] N. Yunes and F. Pretorius, “Fundamental Theoretical Bias in Gravitational Wave Astrophysics and the Parameterized Post-Einsteinian Framework,” *Phys. Rev. D* **80**, 122003 (2009).
- [33] R. Nair, S. Perkins, H. O. Silva, and N. Yunes, “Fundamental Physics Implications on Higher-Curvature Theories from the Binary Black Hole Signals in the LIGO-Virgo Catalog GWTC-1,” *Phys. Rev. Lett.* **123**, 191101 (2019).
- [34] E. Berti *et al.*, “Testing General Relativity with Present and Future Astrophysical Observations,” *Class. Quant. Grav.* **32**, 243001 (2015).
- [35] S. Alexander and N. Yunes, “Chern–Simons modified general relativity,” *Physics Reports* **480**, 1–55 (2009).
- [36] M. Okounkova, L. C. Stein, M. A. Scheel, and S. A. Teukolsky, “Numerical binary black hole collisions in dynamical Chern-Simons gravity,” *Phys. Rev. D* **100**, 104026 (2019).

- [37] E. Barausse, C. Palenzuela, M. Ponce, and L. Lehner, “Neutron-star mergers in scalar–tensor theories of gravity,” *Phys. Rev. D* **87**, 081506 (2013).
- [38] M. Shibata, K. Taniguchi, H. Okawa, and A. Buonanno, “Coalescence of binary neutron stars in a scalar–tensor theory of gravity,” *Phys. Rev. D* **89**, 084005 (2014).
- [39] E. Berti, V. Cardoso, L. Gualtieri, M. Horbatsch, and U. Sperhake, “Numerical simulations of single and binary black holes in scalar–tensor theories: circumventing the no-hair theorem,” *Phys. Rev. D* **87**, 124020 (2013).
- [40] C. Palenzuela, E. Barausse, M. Ponce, and L. Lehner, “Dynamical scalarization of neutron stars in scalar–tensor gravity theories,” *Phys. Rev. D* **89**, 044024 (2014).
- [41] H. Witek, L. Gualtieri, P. Pani, and T. P. Sotiriou, “Black holes and binary mergers in scalar Gauss-Bonnet gravity: Scalar field dynamics,” *Phys. Rev. D* **99**, 064035 (2019).
- [42] J. L. Ripley and F. Pretorius, “Scalarized black hole dynamics in Einstein-dilaton-Gauss-Bonnet gravity,” *Phys. Rev. D* **101**, 044015 (2020).
- [43] W. E. East and J. L. Ripley, “Evolution of Einstein-scalar-Gauss-Bonnet gravity using a modified harmonic formulation,” *Phys. Rev. D* **103**, 044040 (2021).
- [44] M. Okounkova, “Numerical relativity simulation of GW150914 in Einstein dilaton Gauss-Bonnet gravity,” *Phys. Rev. D* **102**, 084046 (2020).
- [45] M. Elley, H. O. Silva, H. Witek, and N. Yunes, “Spin-induced dynamical scalarization, descalarization, and stealthness in scalar-Gauss-Bonnet gravity during a black hole coalescence,” *Phys. Rev. D* **106**, 044018 (2022).

- [46] D. D. Doneva, L. Aresté Saló, K. Clough, P. Figueras, and S. S. Yazadjiev, “Testing the limits of scalar-Gauss-Bonnet gravity through nonlinear evolutions of spin-induced scalarization,” *Physical Review D* **108**, 084017 (2023).
- [47] E. W. Hirschmann, L. Lehner, S. L. Liebling, and C. Palenzuela, “Black hole dynamics in Einstein-Maxwell-dilaton theory,” *Phys. Rev. D* **97**, 064032 (2018).
- [48] F.-L. Julié, “On the motion of hairy black holes in Einstein-Maxwell-dilaton theories,” *JCAP* **01**, 026 (2018).
- [49] V. Cardoso *et al.*, “NR/HEP: roadmap for the future,” *Classical and Quantum Gravity* **29**, 244001 (2012).
- [50] M. Corman, L. Lehner, W. E. East, and G. Dideron, “Nonlinear studies of modifications to general relativity: Comparing different approaches,” *Physical Review D* **110**, 084048 (2024).
- [51] A. Held and H. Lim, “Black-hole binaries and waveforms in Quadratic Gravity,” 2025.
- [52] R. M. Wald, *General Relativity* (University of Chicago Press, 1984).
- [53] R. D’Inverno, *Introducing Einstein’s Relativity* (Oxford University Press, 1992).
- [54] E. Newman and R. Penrose, “An Approach to Gravitational Radiation by a Method of Spin Coefficients,” *J. Math. Phys.* **3**, 566–578 (1962).
- [55] S. Giri, U. Danielsson, L. Lehner, and F. Pretorius, “Exploring Black Hole Mimickers: Electromagnetic and Gravitational Signatures of AdS Black Shells,” *Physical Review D* **111**, 024007 (2025).
- [56] K. Clough, T. Dietrich, and S. Khan, “What no one has seen before: gravitational waveforms from warp drive collapse,” *The Open Journal of Astrophysics* **7** (2024).

- [57] C. Bambi *et al.*, “Black hole mimickers: from theory to observation,” arXiv preprint arXiv:2505.09014 (2025).
- [58] N. Yunes and X. Siemens, “Gravitational-wave tests of general relativity with ground-based detectors and pulsar-timing arrays,” *Living Rev. Relativ.* **16**, 9 (2013).
- [59] K. Yagi and L. C. Stein, “Black hole based tests of general relativity,” *Class. Quant. Grav.* **33**, 054001 (2016).
- [60] J. Meidam, M. Agathos, C. Van Den Broeck, J. Veitch, and B. S. Sathyaprakash, “Testing the no-hair theorem with black hole ringdowns using TIGER,” *Phys. Rev. D* **90**, 064009 (2014).
- [61] H. Yang, K. Yagi, J. Blackman, L. Lehner, V. Paschalidis, F. Pretorius, and N. Yunes, “Black hole spectroscopy with coherent mode stacking,” *Phys. Rev. Lett.* **118**, 161101 (2017).
- [62] R. Abbott *et al.*, “GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo during the First Half of the Third Observing Run,” *Phys. Rev. X* **11**, 021053 (2021).
- [63] N. Afshordi *et al.*, “Black Holes Inside and Out 2024: visions for the future of black hole physics,” 2024.
- [64] C. Bambi *et al.*, “Black hole mimickers: from theory to observation,” 2025.
- [65] L. Bernard, S. Giri, L. Lehner, and R. Sturani, “Generic EFT-motivated beyond General Relativity gravitational wave tests and their curvature dependence: from observation to interpretation,” *Physical Review D* **112**, 124013 (2025).
- [66] M. Okounkova, L. C. Stein, J. Moxon, M. A. Scheel, and S. A. Teukolsky, “Numerical relativity simulation of GW150914 beyond general relativity,” *Phys. Rev. D* **101**, 104016 (2020).

- [67] J. L. Ripley and F. Pretorius, “Dynamics of a $\{\mathbb{Z}\}_2$ symmetric EdGB gravity in spherical symmetry,” *Classical and Quantum Gravity* **37**, 155003 (2020).
- [68] R. Cayuso and L. Lehner, “Nonlinear, noniterative treatment of EFT-motivated gravity,” *Phys. Rev. D* **102**, 084008 (2020).
- [69] A. Held and H. Lim, “Nonlinear dynamics of quadratic gravity in spherical symmetry,” *Phys. Rev. D* **104**, 084075 (2021).
- [70] A. Held and H. Lim, “Nonlinear evolution of quadratic gravity in 3 + 1 dimensions,” *Phys. Rev. D* **108**, 104025 (2023).
- [71] W. E. East and N. Siemonsen, “Instability and backreaction of massive spin-2 fields around black holes,” *Phys. Rev. D* **108**, 124048 (2023).
- [72] T. May and W. E. East, “Nonlinear evolution of the spin-2 black hole superradiant instability,” 2025.
- [73] M. Corman, L. Lehner, W. E. East, and G. Dideron, “Nonlinear studies of modifications to general relativity: Comparing different approaches,” *Phys. Rev. D* **110**, 084048 (2024).
- [74] A. Sen, “Rotating Charged Black Hole Solution in Heterotic String Theory,” *Phys. Rev. Lett.* **69**, 1006–1009 (1992).
- [75] A. Sen, “Black hole solutions in heterotic string theory on a torus,” *Nucl. Phys. B* **440**, 421–440 (1995).
- [76] W. Israel, “Event Horizons in Static Vacuum Space-Times,” *Phys. Rev.* **164**, 1776–1779 (1967).
- [77] B. Carter, “Axisymmetric Black Hole Has Only Two Degrees of Freedom,” *Phys. Rev. Lett.* **26**, 331–333 (1971).

- [78] H. O. Silva, H. Witek, M. Elley, and N. Yunes, “Dynamical Descalarization in Binary Black Hole Mergers,” *Phys. Rev. Lett.* **127**, 031101 (2021).
- [79] C. N. Pope, D. O. Rohrer, and B. F. Whiting, “Perturbations of Black Holes in Einstein-Maxwell-Dilaton-Axion (EMDA) Theories,” 2025.
- [80] S. K. Sahoo, N. Yadav, and I. Banerjee, “Imprints of Einstein–Maxwell–dilaton–axion gravity in the observed shadows of Sgr A* and M87*,” *Phys. Rev. D* **109**, 044008 (2024).
- [81] S. K. Sahoo and I. Banerjee, “Deciphering signatures of Kerr–Sen black holes in presence of plasma from the Event Horizon Telescope data,” *JCAP* **10**, 100 (2025).
- [82] S. Gasparotto, J. Zosso, L. Aresté Saló, D. D. Doneva, and S. S. Yazadjiev, “Gravitational Memory from Hairy Binary Black Hole Mergers,” 2026.
- [83] M. Agathos, W. Del Pozzo, T. G. F. Li, C. Van Den Broeck, J. Veitch, and S. Vitale, “TIGER: A data analysis pipeline for testing the strong-field dynamics of general relativity with gravitational wave signals from coalescing compact binaries,” *Phys. Rev. D* **89**, 082001 (2014).
- [84] M. Shibata and T. Nakamura, “Evolution of three-dimensional gravitational waves: Harmonic slicing case,” *Phys. Rev. D* **52**, 5428–5444 (1995).
- [85] T. W. Baumgarte and S. L. Shapiro, “Numerical integration of Einstein’s field equations,” *Phys. Rev. D* **59**, 024007 (1998).
- [86] T. W. Baumgarte and S. L. Shapiro, *Numerical Relativity: Solving Einstein’s Equations on the Computer* (Cambridge University Press, Cambridge, UK, 2010).
- [87] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower, “Accurate evolutions of orbiting black-hole binaries without excision,” *Phys. Rev. Lett.* **96**, 111101 (2006).

- [88] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, “Gravitational wave extraction from an inspiraling configuration of merging black holes,” *Phys. Rev. Lett.* **96**, 111102 (2006).
- [89] J. DeBuhr, B. Zhang, M. Anderson, D. Neilsen, E. W. Hirschmann, T. Grenga, and S. Paolucci, “Relativistic Hydrodynamics with Wavelets,” *The Astrophysical Journal* **867**, 112 (2018).
- [90] M. Anderson, E. W. Hirschmann, S. L. Liebling, and D. Neilsen, “Relativistic MHD with adaptive mesh refinement,” *Classical and Quantum Gravity* **23**, 6503–6524 (2006).
- [91] H.-O. Kreiss and J. Oliger, *Methods for the Approximate Solution of Time Dependent Problems*, Vol. 10 of *Global Atmospheric Research Programme (GARP): GARP Publication Series* (GARP Publication, 1973).
- [92] Z. B. Etienne, T. Assumpção, L. R. Werneck, and S. D. Tootle, “BHaHAHA: A Fast, Robust Apparent Horizon Finder Library for Numerical Relativity,” *Classical and Quantum Gravity* **43**, 075007 (2026).
- [93] M. Alcubierre, J. C. Degollado, and M. Salgado, “The Einstein–Maxwell system in 3+1 form and initial data for multiple charged black holes,” *Phys. Rev. D* **80**, 104022 (2009).
- [94] S. Mukherjee, N. K. Johnson-McDaniel, W. Tichy, and S. L. Liebling, “Conformally curved initial data for charged, spinning black hole binaries on arbitrary orbits,” 2022.
- [95] M. Zilhão, V. Cardoso, C. Herdeiro, L. Lehner, and U. Sperhake, “Collisions of oppositely charged black holes,” *Phys. Rev. D* **89**, 044008 (2014).
- [96] M. Zilhão, V. Cardoso, C. Herdeiro, L. Lehner, and U. Sperhake, “Collisions of charged black holes,” *Phys. Rev. D* **85**, 124062 (2012).

- [97] G. Bozzola and V. Paschalidis, “Initial data for general relativistic simulations of multiple electrically charged black holes with linear and angular momenta,” *Phys. Rev. D* **99**, 104044 (2019).
- [98] A. G. Abac *et al.*, “GW250114: Testing Hawking’s Area Law and the Kerr Nature of Black Holes,” *Physical Review Letters* **135**, 111403 (2025).
- [99] A. Carroll, E. W. Hirschmann, H. Lim, D. Neilsen, D. F. Van Komen, and S. Vander Ploeg Fallon, “Nonlinear Stability of Kerr–Sen Black Holes in Merging Binaries,” (2026).
- [100] G. Bozzola and V. Paschalidis, “General Relativistic Simulations of the Quasi-Circular Inspiral and Merger of Charged Black Holes: GW150914 and Fundamental Physics Implications,” *Physical Review Letters* **126**, 041103 (2021).
- [101] G. Bozzola and V. Paschalidis, “Can Quasi-Circular Mergers of Charged Black Holes Produce Extremal Black Holes?,” *Physical Review D* **108**, 064010 (2023).
- [102] R. Luna, G. Bozzola, V. Cardoso, V. Paschalidis, and M. Zilhão, “Kicks in Charged Black Hole Binaries,” *Physical Review D* **106**, 084017 (2022).
- [103] G. W. Gibbons, “Vacuum Polarization and the Spontaneous Loss of Charge by Black Holes,” *Communications in Mathematical Physics* **44**, 245–264 (1975).
- [104] Z. Pan and H. Yang, “Black Hole Discharge: Very-High-Energy Gamma Rays from Black Hole-Neutron Star Mergers,” *Physical Review D* **100**, 043025 (2019).
- [105] J. H. Horne and G. T. Horowitz, “Rotating Dilaton Black Holes,” *Phys. Rev. D* **46**, 1340–1346 (1992).
- [106] S. Mignemi and N. R. Stewart, “Charged black holes in effective string theory,” *Phys. Rev. D* **47**, 5259–5269 (1993).

- [107] M. Khalil, N. Sennett, J. Steinhoff, J. Vines, and A. Buonanno, “Hairy binary black holes in Einstein-Maxwell-dilaton theory and their effective-one-body description,” *Physical Review D* **98**, 104010 (2018).
- [108] K. Yagi, L. C. Stein, N. Yunes, and T. Tanaka, “Post-Newtonian, Quasi-Circular Binary Inspirals in Quadratic Modified Gravity,” *Physical Review D* **85**, 064022 (2012).
- [109] F.-L. Julié and E. Berti, “Post-Newtonian dynamics and black hole thermodynamics in Einstein-scalar-Gauss-Bonnet gravity,” *Physical Review D* **100**, 104061 (2019).
- [110] P. K. Gupta, “Binary dynamics from Einstein-Maxwell theory at second post-Newtonian order using effective field theory,” *Physical Review D* **112**, 104047 (2025).
- [111] G. Bozzola and V. Paschalidis, “General relativistic simulations of the quasi-circular inspiral and merger of charged black holes: GW150914 and fundamental physics implications,” *Phys. Rev. Lett.* **126**, 041103 (2021).
- [112] Z. B. Etienne, “NRPyPN: Validated Post-Newtonian Expressions for Binary Black Hole Initial Data,” 2023.
- [113] J. Healy, C. O. Lousto, H. Nakano, and Y. Zlochower, “Post-Newtonian Quasicircular Initial Orbits for Numerical Relativity,” *Classical and Quantum Gravity* **34**, 145011 (2017).
- [114] A. Ramos-Buades, S. Husa, and G. Pratten, “Simple procedures to reduce eccentricity of binary black hole simulations,” *Phys. Rev. D* **99**, 023003 (2019).
- [115] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, “Improved filters for gravitational waves from inspiraling compact binaries,” *Physical Review D* **57**, 885–907 (1998).
- [116] I. W. Harry and S. Fairhurst, “Targeted coherent search for gravitational waves from compact binary coalescences,” *Physical Review D* **83**, 084002 (2011).

- [117] B. P. Abbott *et al.*, “Effects of waveform model systematics on the interpretation of GW150914,” *Classical and Quantum Gravity* **34**, 104002 (2017).
- [118] G. Bozzola, “kuibit: Analyzing Einstein Toolkit simulations with Python,” *The Journal of Open Source Software* **6**, 3099 (2021).
- [119] G. Papallo and H. S. Reall, “On the local well-posedness of Lovelock and Horndeski theories,” *Phys. Rev. D* **96**, 044019 (2017).
- [120] A. D. Kovacs and H. S. Reall, “Well-posed formulation of scalar-tensor effective field theory,” *Phys. Rev. Lett.* **124**, 221101 (2020).
- [121] R. P. Woodard, “The Theorem of Ostrogradsky,” *Scholarpedia* **10**, 32243 (2015).
- [122] S. E. Perkins, R. Nair, H. O. Silva, and N. Yunes, “Improved gravitational-wave constraints on higher-order curvature theories of gravity,” *Phys. Rev. D* **104**, 024060 (2021).
- [123] J. Ferreira, G. Bozzola, C. A. R. Herdeiro, V. Paschalidis, and M. Zilhão, “Electromagnetic duality degeneracy in dynamical black hole mergers,” 2026.
- [124] C. Palenzuela, L. Lehner, O. Reula, and L. Rezzolla, “Beyond ideal MHD: towards a more realistic modeling of relativistic astrophysical plasmas,” *Monthly Notices of the Royal Astronomical Society* **394**, 1727–1740 (2009).
- [125] J. M. Bowen and J. York, James W., “Time-asymmetric initial data for black holes and black-hole collisions,” *Phys. Rev. D* **21**, 2047–2055 (1980).
- [126] Z. B. Etienne, “Improved moving-puncture techniques for compact binary simulations,” *Phys. Rev. D* **110**, 064045 (2024).
- [127] J. A. Valiente Kroon, “Nonexistence of Conformally Flat Slices in Kerr and Other Stationary Spacetimes,” *Phys. Rev. Lett.* **92**, 041101 (2004).

-
- [128] A. Garat and R. H. Price, “Nonexistence of conformally flat slices of the Kerr spacetime,” *Phys. Rev. D* **61**, 124011 (2000).
- [129] S. Dain, C. O. Lousto, and R. Takahashi, “New conformally flat initial data for spinning black holes,” *Phys. Rev. D* **65**, 104038 (2002).
- [130] M. Ansorg, B. Brügmann, and W. Tichy, “A single-domain spectral method for black hole puncture data,” *Physical Review D* **70**, 064011 (2004).