

FACE ATLAS: A transdiagnostic latent atlas of severe mental illness decouples biological burden from clinical severity and forecasts functional outcome at the group level

Andrii Kulakovskiy^{1*} Ophélie Gaudin^{1,2} Léah Camarcat¹ Engin Altunlu¹
Marion Leboyer^{1,2}

¹Fondation FondaMental ²Inserm

June 28, 2026

Abstract

Psychiatric diagnoses aggregate biologically heterogeneous patients, which limits mechanistic stratification and prognosis. We analysed the harmonized baseline assessment of $N = 9,013$ patients with bipolar disorder, schizophrenia or major depression from the three FACE cohorts using one global, missingness-aware Bayesian bifactor / exploratory structural-equation model, specified and fit to each patient’s observed cells only (no imputation), with diagnosis withheld from model fitting. The model resolved a transdiagnostic coordinate system: a general functional-burden factor (G) and seven specific dimensions—cognition, immunometabolic load, sleep, developmental risk, suicidality, mania and substance use. The central finding is that **immunometabolic load is largely decoupled from functional impairment**: under a freely-correlated general factor it was the least burden-entangled domain (correlation with $G \approx 0.10$, versus 0.39 for cognition and 0.42 for sleep), and it remained so after adjustment for medication, adiposity and site. This axis is anchored by body composition (BMI loading ≈ 0.95) but does not reduce to it: excluding the anthropometric indicators, a coherent cardiometabolic-inflammatory axis survived (Tucker congruence 0.90 with the full axis); because the only inflammatory marker available was C-reactive protein, which is itself adiposity-driven, we interpret the axis as cardiometabolic rather than as an independent inflammatory dimension. The coordinate space was a graded **continuum, not discrete biotypes** in this measurement space—apparent clustering was reproduced by a single-Gaussian null—yet five archetypal extremes gave a clinically legible reading, including an immunometabolic corner and a severe, clean-biology corner with comparable functional burden but opposite biological profiles. Scored forward onto follow-up without re-estimation, immunometabolic coordinates were the most stable axis (test–retest intraclass correlation 0.91) whereas symptom axes were state-like. The archetype phenotype predicted two-year functioning beyond diagnosis, baseline severity and baseline functioning, but the signal was **group-level**: a held-out expected log predictive density gain of +62.8 and a two-year functional-remission gradient of 17%–52% across corners (the immunometabolic corner worst), yet only a small individual-level discrimination gain (remission AUC +0.010), and the map was co-informative with rather than superior to DSM-5 (incremental predictive density +17.3 for the map versus +29.0 for diagnosis). These results define a transdiagnostic biological atlas of severe mental illness that supports mechanistic stratification and group-level functional forecasting; individual-level clinical utility would require external validation, incident-event outcomes and randomized treatment data.

Keywords: transdiagnostic psychiatry; dimensional psychopathology; bifactor model; metabolic syndrome; inflammation; bipolar disorder; schizophrenia; major depression; archetypal analysis; precision psychiatry.

*Correspondence: andrii.kulakovskiy@fondation-fondamental.org

Table 1: **Sample characteristics by cohort (FACE baseline, $N = 9,013$).** Aggregate-only summary; sex coded female. Retention rows are the percentage of the baseline cohort re-assessed at the 12- and 24-month visits.

Characteristic	Bipolar	Schizophrenia	Depression	Overall
N	6,252	2,209	552	9,013
Age, years — mean (SD)	39.3 (13.4)	31.1 (9.6)	51.0 (13.3)	38.0 (13.4)
Age, years — median [IQR]	38 [28–49]	29 [23–37]	52 [42–60]	36 [27–48]
Female — n (%)	3,969 (63.5)	600 (27.2)	323 (58.5)	4,892 (54.3)
Education, years — mean (SD)	14.2 (2.8)	12.3 (2.6)	13.2 (3.2)	14.0 (2.8)
Recruitment sites — n	15	12	13	21
Education missing — n (%)	1,803 (28.8)	1,646 (74.5)	228 (41.3)	3,677 (40.8)
<i>DSM-5 subtype — n (% of cohort)</i>				
Bipolar I	2,635 (42.1)	—	—	
Bipolar II	2,956 (47.3)	—	—	
Bipolar, unspecified	661 (10.6)	—	—	
Schizophrenia	—	1,692 (76.6)	—	
Schizoaffective disorder	—	476 (21.5)	—	
Schizophreniform disorder	—	41 (1.9)	—	
Major depressive disorder	—	—	552 (100.0)	
Retained at 12 months — %	49.2	43.6	42.2	47.4
Retained at 24 months — %	35.6	26.9	24.5	32.8

1 Introduction

Psychiatric diagnoses are clinically indispensable but biologically nonspecific. The categories of the DSM aggregate patients who differ widely in their biology, course and treatment response, and they are undercut by pervasive comorbidity, within-category heterogeneity and diagnostic instability over time [1, 2]. Two influential programmes argue in response that psychopathology is better described by continuous, *transdiagnostic* dimensions than by categorical boundaries: the U.S. National Institute of Mental Health’s Research Domain Criteria (RDoC) [1] and the Hierarchical Taxonomy of Psychopathology (HiTOP) [2, 3]. A recurring motif within this work is a general factor that spans disorders—the “ p factor”—most defensibly read as a dimension of life impairment rather than a latent liability to specific symptoms [4, 5], though the meaning of such general factors remains contested [6, 7].

This dimensional structure has, however, been mapped almost entirely from a single *kind* of measurement—symptom report [2, 4] or brain imaging [8–10]. The routinely collected *biology* that drives the excess physical-health mortality of severe mental illness—its joint metabolic and inflammatory load—has not been placed *inside* the transdiagnostic factor space across disorders, even though an independent literature suggests it forms a partly separable immunometabolic axis that tracks atypical, energy-related symptoms more than overall severity [11–13], and that metabolic burden follows medication, adiposity and chronicity more than symptom load [14–16]. Whether this load is separable from a transdiagnostic severity axis, and by how much, has not to our knowledge been measured directly.

We therefore ask three questions of a single transdiagnostic model. **(i) Measurement structure:** what latent structure organizes routine clinical, cognitive and biological data when they are placed in one coordinate system with diagnosis held out, and is the resulting patient space a graded continuum or a set of discrete kinds? **(ii) Biology–severity separation:** is immunometabolic load separable from a general functional-burden axis, and by how much? **(iii) Prognostic value:** does the latent representation forecast functional outcome beyond diagnosis and current severity? The structural half of (i) is itself contested: normative-modelling studies report individual heterogeneity so extensive that “the average patient” is barely informa-

Table 2: **The eight-factor transdiagnostic map.** Anchoring indicators, entanglement with general burden, adjudication verdict and cross-cohort measurement invariance for each factor. Correlation with G is from the correlated- G model (continuous backbone); for mania and substance it is the bifactor direct $|\lambda_G|$. Substance is pinned orthogonal to the correlated block (its cross-factor correlations are non-identifiable). Depression/anxiety instruments are cross-loading windows on G (loadings 0.66–0.80), not a separate factor; the map is otherwise simple-structure apart from three earned cross-loadings into cognition (childhood adversity and poor sleep, $|\lambda| \leq 0.09$, 95% CI excluding zero).

Factor	Anchoring indicators	corr G	Verdict	Invariance
G (burden)	FAST, EGF, EQ-5D (no symptoms)	—	severity factor	partial (SZ)
Cognition	TMT, WAIS coding, fluency	0.39	confirmed	invariant
Immunometabolic	BMI, lipids, glycaemia, BP, CRP, leukocytes	0.07	confirmed (single biology axis)	invariant
Sleep	PSQI total + subscales	0.42	confirmed	invariant
Mania	YMRS, Altman	0.15	confirmed	Altman floors (DR)
Suicidality	ISF ideation / attempt (binary)	—	confirmed (mixed)	3-cohort
Developmental risk	CTQ, perinatal, family history	—	confirmed (proxy)	invariant
Substance	alcohol/cannabis SUD, nicotine	0.13	confirmed (orthogonal)	BP–SZ (2-cohort)

tive [8, 9], whereas the B-SNIP programme reports replicable, brain-based psychosis “biotypes” that cut across DSM boundaries [17–19]—a tension whose resolution may depend on which measurement space is interrogated.

We use latent modelling as a measurement tool, not as a claim that latent factors are more real than the items that define them. Routine clinical and biological data are high-dimensional, redundant and pervasively incomplete; a latent model fit to each patient’s observed cells pools many noisy, overlapping indicators into a few axes, places heterogeneous data types (laboratory values, ordinal scales, counts) on one comparable scale, and gives every patient a position with quantified uncertainty rather than discarding incomplete records. Whether that representation *adds* anything over the raw indicators is then an empirical question, which we test directly: against the full raw indicator set under a matched classifier, the eight-dimension map is a sufficient representation for predicting deterioration and near-sufficient for recovery.

Using the harmonized baseline of the three FACE cohorts—bipolar disorder, schizophrenia and major depression, $N = 9,013$ —we build *FACE-ATLAS*: one global, missingness-aware Bayesian bifactor / exploratory structural-equation model with mixed likelihoods, fit to each patient’s observed cells only, that places clinical, cognitive, behavioural and laboratory measurements in one coordinate system while holding diagnosis out as metadata. We then score the fitted map forward onto two years of follow-up to test whether it persists and whether it predicts outcome. The answers are: an eight-dimension map—a general functional-burden factor and seven specific axes—on which immunometabolic load is the domain least coupled to functional severity; a graded continuum rather than discrete biotypes *in this measurement space*, read through five archetypal extremes; a geometry that persists, with durable biology and moving symptoms; and a real but modest, *group-level* prognostic signal for functioning that is co-informative with diagnosis rather than a replacement for it. We report the modest and null results—including a treatment-moderation analysis that this observational cohort can only bound—as deliberately

Table 3: **The five archetypes.** Corners of the clinical–biological continuum—directions of maximal phenotype on the convex hull of the cloud, not discrete biotypes (the space carries no density gap; single-Gaussian falsification null $z = 1.13$, n.s.). $A = 5$ is the largest cross-seed-stable corner count (minimum Tucker congruence 0.979, explained variance 0.601, clean stability cliff to 0.436 at $A = 6$). Sizes are dominant ($\arg \max$) memberships over $N = 9,013$ patients; the inferential object is the continuous simplex weight vector, not the hard label.

Archetype	Defining profile (latent z-scale)	Clinical reading	N (dominant)
A0	↑sleep +2.8, ↑mania +1.7, ↓severity	<i>activation/sleep</i> (BP-enriched)	1,448
A1	↑severity +1.9, ↓immunometabolic −1.9, ↓developmental	<i>severe, clean-biology</i>	1,584
A2	↑immunometabolic +3.5, ↑severity +2.4, ↑suicidality	<i>immunometabolic</i> <i>burden</i> (the biology corner)	1,426
A3	↑developmental +2.8, ↑suicidality +1.4, ↓immunometabolic −2.1	<i>trauma/suicidality</i>	2,004
A4	↓sleep −2.7, ↓severity −2.0, ↓mania	<i>low-burden/well</i>	2,551

as the positive ones.

Methodologically, FACE-ATLAS configures established latent-variable tools [20–22] into a single diagnosis-held-out, uncertainty-propagating atlas: mixed and Gaussian-copula observation layers, an observed-cell likelihood for structured missingness, a correlated- G arm that turns the biology–severity separation into a measured quantity, an archetypal representation of a continuous patient space, and downstream errors-in-variables prognosis and treatment-moderation analyses. The contribution is the specification and validation chain rather than a new generic estimator: routine biological load is placed inside a transdiagnostic psychiatric factor space, its separation from functional burden is measured rather than assumed, and its temporal and prognostic implications are tested without converting a continuum into biotypes.

2 Results

2.1 Sample and observed-data modelling population

The analysis sample was the harmonized baseline (visit V0) of the three FACE cohorts: $N = 9,013$ patients (bipolar disorder [BP] 6,252; schizophrenia [SZ] 2,209; major depression [DR] 552) recruited across 21 expert-centre sites (Table 1). The cohorts differ in age, sex distribution and chronicity, as expected for these disorders. We deliberately fit the measurement model on the *full* sample without completeness selection. Of the 201 usable harmonized common variables, 143 form the modelled-indicator pool (mean cell missingness 39.8%, preserved as missing and never imputed) and the fitted eight-factor map carries 109 of them—88 in the Gaussian-copula continuous block and 21 in the explicit non-Gaussian block; the remaining 58 are identifiers, covariates and validation labels (ambiguous-direction laboratory values, treatment-efficacy ratings, ECG intervals, reproductive/hormonal and smoking-history confounders) and were never modelled as indicators. Diagnosis (cohort and DSM-5 subtype) was held out of the model and used only as a validation or metadata label. The FACE-ATLAS map is one global, cohort-reweighted, full-sample fit estimated with a Gaussian-copula (rank-based) marginal model;

because the same structure is recovered under a plain Gaussian likelihood, the central findings do not depend on that modelling choice—a robustness result, not an assumption (Supplement A). Follow-up visits at 12 and 24 months (retained: 47.4% and 32.8% of V0; Extended Data Fig. F3) were used only to test temporal coherence and prognosis, never to define the map. The high-level model is summarized below and in Methods; the full derivation of its observed-data likelihood—the rotation and identification argument, the covariance identity and the equivalence with full-information maximum likelihood—is given in Supplement A. Figure 1 presents the atlas at a glance; the study design and the three core modelling principles are detailed in Extended Data Fig. F1.




Figure 1: **Illustration of the FACE ATLAS: latent dimensions of severe mental illness and corresponding archetypes.** The global, missingness-aware Bayesian bifactor / exploratory structural-equation model, fit to the FACE baseline ($N = 9,013$) with diagnosis held out, yields an eight-dimension transdiagnostic latent space—a general functional-burden factor (G) and seven specific dimensions. The patient cloud is a graded continuum (best silhouette 0.140, indistinguishable from a single-Gaussian null 0.137 ± 0.002), read through five archetypal extremes whose convex blends describe every patient: an *activation / sleep* corner (A0), a *severe, clean-biology* corner (A1), an *immunometabolic* corner (A2; the biology pole), a *trauma / suicidality* corner (A3), and a *low-burden* pole (A4). Immunometabolic load is largely dissociated from general burden; it is the most durable axis over two years; the archetype phenotype improves held-out two-year functional prediction ($\Delta\text{ELPD} +62.8 \pm 11.2$); and two-year functional remission ranges 17%–52% across corners, the immunometabolic corner worst. A1 and A2 carry comparable severity but opposite biology.

2.2 The fitted object: a posterior coordinate for each patient

The primary output of FACE-ATLAS is not a cluster label but a *posterior coordinate distribution*. For patient i the model returns an eight-dimensional coordinate

$$f_i = (G_i, D_{i,\text{cog}}, D_{i,\text{immet}}, D_{i,\text{sleep}}, D_{i,\text{mania}}, D_{i,\text{suic}}, D_{i,\text{dev}}, D_{i,\text{subst}}) \in \mathbb{R}^8, \quad (1)$$

one general functional-burden coordinate G_i and seven specific coordinates (cognition, immunometabolic, sleep, mania, suicidality, developmental risk, substance use)—the seven specific axes are established as distinct, data-supported dimensions in §2.3 (Table 2). What the model actually delivers is the posterior of this coordinate given the patient’s *observed* indicators x_{i,O_i} alone—no imputation, no completed data matrix (Methods):

$$p(f_i | x_{i,O_i}) \approx \mathcal{N}(\hat{f}_i, S_i), \quad (2)$$

whose mean $\hat{f}_i$ is the patient’s *position* and whose covariance S_i is the *uncertainty* of that position—a location with error bars, not a category. Two patients can share an estimated immunometabolic score yet differ in its reliability (a complete laboratory panel versus a single observed marker), and every downstream analysis in this paper consumes the full $(\hat{f}_i, S_i)$ rather than treating coordinates as exact, so a sparsely measured patient is never mistaken for a precisely located one. The unit of analysis is thus a position on eight continuous axes with quantified uncertainty, not a biotype label.

2.3 The eight dimensions of the transdiagnostic atlas

From ten candidate constructs, one global model returned a stable eight-dimension structure: a general functional-burden factor (G) plus seven weakly correlated specific axes—cognition, immunometabolic, sleep, mania, suicidality, developmental risk and substance use (Figure 2; Table 2). Inter-factor correlations among the specific axes were weak throughout (mean absolute off-diagonal ≈ 0.08 ; Figure 3b), confirming genuinely distinct axes rather than one collapsed dimension; the single notable coupling was a modest mania–sleep correlation (≈ 0.24). G was anchored only by functioning and global-severity indicators (FAST loading 0.91, EGF 0.64) and carried no symptom content, supporting its conservative reading as transdiagnostic *functional burden* rather than a symptom-liability “ p ” factor. Depression and anxiety instruments (MADRS, QIDS, STAI) did not form a separate factor; they loaded 0.61–0.82 on G as cross-loading “windows” onto burden. The biology axis is a single *immunometabolic* factor—cardiometabolic and inflammatory markers (body-mass index, HbA_{1c}, lipids, C-reactive protein) load on one common dimension, anchored by body composition (BMI loading ≈ 0.95 , CRP ≈ 0.37 ; all 37 loadings with credible intervals in Table A4), the data placing cardiometabolic and immune burden on a shared axis rather than two. Body size anchors this axis but does not constitute it: excluding the three anthropometric indicators and refitting, a coherent cardiometabolic–inflammatory axis survived—all 34 remaining markers credible, re-anchored on blood pressure, lipids and inflammation, and highly congruent with the full axis (Tucker $\varphi = 0.90$; Supplement A.6). Of the ten prior candidate factors, anhedonia was rejected (absorbed into G and the depression/anxiety windows), and a further three constructs with no common indicator—impulsivity, negative symptoms and sensory abnormality—were declared *not testable*, each verdict an outcome of the analysis rather than an assumption (Table 2). The structure was not an artefact of the priors: an independent flat-prior refit reproduced the loadings and factor correlations essentially exactly (Tucker congruence $\varphi = 1.00$). Information criteria also ranked the bifactor above unidimensional and correlated-factor alternatives, but we treat this as secondary evidence, since such criteria are known to favour bifactor structures for reasons of model form independent of the data-generating structure [23, 24]. Absolute fit was acceptable across both likelihood blocks (continuous Bayesian standardized root-mean-square residual ≈ 0.07 ; 21/22 non-Gaussian indicators reproduced within the 90% posterior-predictive interval). *Clinically*, the axes answer two different questions: G answers *how impaired* the patient is, while the seven specific axes answer *what kind* of burden the patient carries.

The map is otherwise simple-structure—each indicator loads on its home axis and G —by design rather than by fiat. A sparsity-inducing regularized (“Finnish”) horseshoe prior is placed on every off-home specific-to-specific loading: a global shrinkage term pulls the whole set

toward zero (the prior’s default belief is “no cross-loadings”), heavy local tails let a genuinely supported cross-loading escape, and a slab caps its magnitude. This protects the thin factors (substance, mania, each defined by very few indicators) from dilution while letting small, clinically real cross-talk emerge. Given total freedom—a continuous sparse-ESEM refit that frees every off-home cell— $\approx 83\%$ shrank to ≈ 0 (the simple structure is *recovered from the data*, not imposed), and exactly three cross-loadings survived into the full model with 95% credible intervals excluding zero, all onto cognition: childhood adversity (CTQ, -0.094) and two sleep items (PSQI latency $+0.057$, daytime dysfunction -0.070). Given the freedom to load anywhere, the model reproduced known clinical cross-talk—adversity and poor sleep weighing weakly on cognition—and nothing spurious: positive evidence that the map is well-specified, not an artefact of rigid zeros (Supplement A).

2.4 Immunometabolic load is largely decoupled from functional burden (the central result)

Because the bifactor identification fixes G orthogonal to the specifics by construction, we re-estimated the model allowing G to correlate freely with the specific axes. The resulting vector of general-specific correlations, $\phi_{Gd} = \text{Corr}(G, D)$ (Methods, Eq. 8), is the *primary estimand* of FACE-ATLAS: it converts “biology is separable from severity” from a modelling constraint into a measured quantity. Under this correlated- G model, immunometabolic load was the *least* entangled with functional burden: a correlation with G of ≈ 0.10 , against 0.39 for cognition and 0.42 for sleep (Figure 4). In shared-variance terms the gap is stark—immunometabolic load shares only $\approx 1\%$ of its variance with general burden, against $\approx 15\%$ (cognition) and $\approx 18\%$ (sleep). And because the specific axes are themselves only weakly inter-correlated (mean absolute off-diagonal ≈ 0.08), immunometabolic load is weakly coupled not only to functional burden but to the symptom axes as well. Two patients with the same overall impairment can therefore carry very different immunometabolic loads—a patient’s biological burden rides largely on axes that the global clinical-severity picture does not capture. This dissociation was robust to the obvious confounders. Adjusting each indicator for age, sex, education and recruitment site, and then additionally for antipsychotic exposure, did *not* raise the immunometabolic correlation—if anything it lowered it (to ≈ 0.07), and the immunometabolic axis remained the least entangled at every step (Figure 4; Supplement A). Consistent with a large independent literature (Discussion), we state this as relative, not absolute, independence: immunometabolic load is *largely independent of a general functional-burden axis*, the low correlation quantifying—at the low end of what that literature would predict—a pattern previously described only qualitatively. *Clinically*, this means two patients with comparable functional burden can have sharply different cardiometabolic and inflammatory profiles: severity cannot be used as a proxy for biological load. This is the property a biology-aware stratification can exploit, and it is the central premise of everything that follows.

2.5 The coordinate cloud is continuous, not discretely clustered

We next asked whether these eight coordinates fall into separable kinds. They do not—a *negative structural result*. On the cloud $\{(\hat{f}_i, S_i)\}$, a structure-discovery battery run over the posterior draws—cluster tendency, modality, silhouette, the gap statistic, a Gaussian-mixture information criterion, density-based clustering and a topological summary—returned a continuum verdict: the silhouette peaked at only 0.146 (below the 0.15 separation floor), the gap-statistic optimum was a single component, density-based clustering assigned 100% of points to noise, and the topological summary was a single connected component. A single-Gaussian *falsification null* made the verdict decisive: the best partition of the real cloud separated patients no better than slicing a structureless Gaussian blob of identical covariance (best silhouette 0.140 for the real data versus 0.137 ± 0.002 for the null, $z = 1.13$, n.s.; the Gaussian-mixture-optimal number of components collapsed to one in all 20 posterior draws; Figure 5a). There are, in this




Figure 2: **Posterior loading atlas of the measurement model.** Rows are clinical indicators grouped by their home factor (coloured left strip and alternating bands); columns are the general functional-burden dimension G and the seven specific latent axes (D1–D7). Dot size and colour encode the absolute posterior median loading; outlined dots indicate loadings whose 95% credible interval excludes zero, and a heavier ring marks the home-factor anchor. Specific axes are each anchored by a coherent clinical indicator block, whereas G captures broad functional burden and carries cross-loading “windows” over the depression and anxiety instruments (MADRS, QIDS, STAI; shaded on the G column). The three data-supported off-home cross-loadings—childhood adversity and two sleep items onto cognition—render as outlined dots in the cognition column. The full eight-factor map is shown (continuous backbone plus the mania and substance axes from the joint mixed fit); the complete atlas over all modelled indicators is Extended Data Fig. F2. This structure supports an atlas-like reading of the latent space: specific dimensions are clinically localised axes, while G acts as a transdiagnostic burden backbone. Factor-wise interpretability and the inter-factor correlation matrix are in Figure 3.




Figure 3: Factor interpretability and inter-factor correlations. (a) Factor-wise lollipops: for each axis (G and D1–D7), its strongest home indicators with 95% credible whiskers—“what defines each axis.” (b) The inter-factor correlation matrix Φ is weak throughout (specific–specific mean $|\text{off-diagonal}| \approx 0.08$); G is orthogonal to the specific axes by construction (top row and column). The one notable specific coupling is mania–sleep (≈ 0.24); substance is pinned orthogonal to the correlated block (its cross-factor correlations are non-identifiable), and the biology $\times$ symptom cells all stay at the off-diagonal floor.




Figure 4: **Immunometabolic load is the least burden-entangled domain.** Correlation of each continuous specific axis with the general functional-burden factor G (bars), unadjusted and after adjustment for medication, adiposity and site; the bifactor general loading $|\lambda_G|$ (diamonds) is shown for comparison. Immunometabolic load falls in the “biology zone” ($\Phi < 0.15$) and stays there under adjustment, whereas cognition and sleep partly track G .

clinical–biological space, no well-separated, reproducible discrete clusters. This is a statement about discrete clustering in this representation (Gaussian-copula ESEM coordinates at $N = 9,013$ under this pipeline), not a claim that biological subtypes are absent in other feature spaces or larger samples [8, 25, 26], nor a denial of the clinical heterogeneity we do find.

The continuum is genuine and transdiagnostic, not an artefact. Diagnosis is intermixed throughout the cloud (adjusted Rand index 0.006 against the seven DSM-5 subtypes and 0.011 against cohort, both ≈ 0 ; Figure 5b); overall burden forms a smooth gradient along the principal axis (Figure 5c); and immunometabolic load varies along a *different* direction (Figure 5d)—the biology– G dissociation, made visible. The verdict is not a missingness artefact (a classifier predicting position from each patient’s coverage pattern performed below baseline, accuracy 0.611 versus a 0.683 baseline, lift -0.072) and is a tighter description of the coordinate cloud than diagnosis (free-mixture information criterion 185,600 versus 188,200 for a DSM-5-constrained model, $\approx 3\times$ the explained variance at lower BIC). These same properties explain why clustering the *raw* indicators can appear to yield biotypes where the latent space does not [27]: a raw partition is driven largely by measurement-side structure—structured missingness, a few heavy-tailed laboratory variables, redundant indicator blocks, and site-specific instruments that re-encode diagnosis—which the measurement model neutralizes by construction (observed-cell likelihood, per-indicator and Gaussian-copula likelihoods, shared factors, and diagnosis held out under invariance), so the structure that remains is genuinely transdiagnostic and tested against a structureless null rather than assumed. *Clinically*, the appropriate object is therefore a coordinate with uncertainty (Eq. 2), not a categorical biotype.

2.6 Five archetypal extremes define the shape of the continuum

A continuum has no clusters, but it still has a *shape*. Archetypal analysis recovers that shape as a small set of reproducible extreme corners—a *positive geometric result* that coexists with the




Figure 5: **The patient space is a continuum, not biotypes.** (a) A single-Gaussian falsification null: the best silhouette achievable on the real eight-dimensional cloud is no better than on a structureless Gaussian of matched covariance ($z = 1.13$, n.s.); the mixture-optimal number of clusters is one in all posterior draws. (b–d) Principal-component embedding of the coordinates ($N = 9,013$), coloured by diagnosis (fully intermixed), by general burden G (a smooth gradient), and by immunometabolic load (a gradient in a *different* direction).

negative clustering result above. Each patient is written as a convex blend of $A = 5$ extreme phenotypes,

$$\hat{f}_i \approx \sum_{a=0}^4 w_{ia} z_a, \quad w_{ia} \geq 0, \quad \sum_{a=0}^4 w_{ia} = 1. \quad (3)$$

The five z_a are extreme points on the convex hull of the patient cloud, and the weights w_{ia} are *barycentric coordinates*: they quantify how strongly each patient is pulled toward each extreme. Dominant-archetype labels are used only for summaries; the inferential object is the continuous weight vector w_i , not a class assignment. The number of corners ($A = 5$) was set by cross-seed stability rather than by a parsimony rule—five corners were the largest reproducible solution (cross-seed Tucker congruence 0.979, explained variance 0.60), with a clean stability cliff at six corners ($0.979 \rightarrow 0.436$)—and uncertainty was propagated by refitting Eq. (3) across posterior draws (Supplement B).

The five extremes separate biology from symptoms from severity (Figure 6): an *activation / sleep* corner (A0; high sleep and mania), a *severe, clean-biology* corner (A1; high overall severity but low immunometabolic load), an *immunometabolic* corner (A2; high immunometabolic load with severity and suicidality)—the biology pole—a *trauma / suicidality* corner (A3; high developmental risk and suicidality), and a *low-burden* pole (A4; everything low). The immunometabolic corner A2 and the severe corner A1 carry comparable general burden but opposite biological




Figure 6: **The archetype simplex: every patient a convex blend of five extremes.** The five archetypal extremes are the corners—**A0** activation / sleep, **A1** severe but clean-biology, **A2** immunometabolic (the biology pole), **A3** trauma / suicidality, and **A4** low-burden—and every patient sits *inside* the simplex as a convex blend whose barycentric weights w_i sum to one (worked example, foot). The central dissociation is geometric: **A1** and **A2** are distinct corners at essentially the *same* general burden but opposite biology—the biology–severity dissociation, rendered as a direction one can point to.

profiles—A1 versus A2 is precisely why *severity is not biology*, the central dissociation now rendered as geometry. The symptom space itself resolves into two distinct corners—activation/sleep (A0) and trauma/suicidality (A3)—rather than one combined symptom pole. Patients sit *between* the corners far more than at them: the weight vectors are diffuse (median normalized entropy 0.65), so most patients are convex blends rather than single-corner cases. Critically, because immunometabolic load varies continuously without a density *gap*, it never forms a clean partition boundary at any granularity; it surfaces instead as a *direction of maximal phenotype*—an archetypal corner—which is exactly why the continuous coordinates and the weight vectors w_i , not any hard tessellation, are the primary objects downstream. *Clinically*, the archetypes provide a vocabulary for reading the continuum, not a replacement for it.

2.7 The geometry persists over two years (the map persists)

Scored forward onto the 12- and 24-month visits on the *fixed* model (observed cells, uncertainty propagated, never re-estimated; baseline coordinates reproduced at Pearson $r \approx 0.99$ through this path), the measurement remained invariant for all four testable backbone axes (longitudinal Tucker congruence $\varphi = 0.987$ –0.996; the immunometabolic axis invariant at $\varphi = 0.987$; Extended Data Fig. F4). Coordinate change can therefore be read as patient change rather than instrument drift. A measurement-error decomposition then separated trait from state (Figure 7). The durable, trait-like axes were the biological and cognitive ones, with immunometabolic load the single most durable axis (test–retest intraclass correlation: immunometabolic 0.91, cognition 0.70); the moving, state-like axes were the symptom ones (substance 0.49, sleep 0.47, suicidality 0.43, developmental risk 0.39). General burden was trait *by rank* (intraclass correlation 0.62): patients largely kept their relative position even as the cohort as a whole improved on severity




Figure 7: **Durable biology, moving symptoms.** (a) Trait fraction (test–retest intraclass correlation, with 94% credible intervals) over follow-up: the biological and cognitive axes are trait-like, the symptom axes state-like; mania is data-limited (hatched). (b) Individual rank stability versus population-level change: patients keep their biological rank (high intraclass correlation, near-zero population shift) while the cohort slides down the severity and symptom axes.

and symptoms (population shift -0.46 on G and -0.84 on suicidality, while immunometabolic load stayed static, $+0.04$). The archetype geometry persisted—each patient’s weight vector w_i moved little over two years (median cosine similarity 0.81). *Clinically*, the reading is compact: *stratify on the durable biology; monitor the moving symptoms*.

2.8 Archetype position predicts group-level two-year functioning (the map predicts)

We then asked whether a baseline coordinate or phenotype predicts a two-year outcome *incrementally beyond* diagnosis, current severity and the baseline value of the outcome, using an errors-in-variables Bayesian model that propagated each coordinate’s measurement uncertainty. The map added predictive signal for *functioning* but not for severity (which was saturated by its own baseline autoregression). The signal was carried by the *phenotype*, not by any single laboratory axis: the five archetypal extremes added the most held-out predictive value (expected log predictive density gain $+62.8$ over the diagnosis-plus-severity-plus-baseline reference), the seven specific axes added $+38.1$, while the isolated “durable biology” axes *alone* sat at the bar ($+2.3$, ambiguous; Figure 8b). Biology still predicts functioning, but as a *corner of the phenotype* (the immunometabolic archetype) rather than as a standalone block—the immunometabolic axis itself carries a credible adverse direction (higher load $\rightarrow$ worse future functioning). No hard partition improved on the continuous and archetype encodings (every tessellation added $\approx +16$ to $+20$, strictly dominated by the archetypes); the operative number of clusters is therefore *none*—the actionable object is a continuous position, not a category. The map was *co-informative* with DSM-5 rather than redundant with it (functioning: map $+17.3$, DSM-5 $+29.0$, both $+62.6$; Figure 8c), and *course-dependent*, with the signal concentrated in the episodic, open-course cohorts (bipolar disorder and depression) and weak when bipolar disorder was removed.

Translated to the clinic, ranking patients by their dominant archetypal extreme yielded two-year functional-remission rates from 17% to 52%, transdiagnostically (Figure 8a). The *immunometabolic* corner A2 carried the worst functional prognosis (17%) and the low-burden pole A4 the best (52%): A2 versus A4 is why the biology corner *matters prognostically*. A2 also remained worse than the equally severe clean-biology corner A1 (17% vs 22%)—the severity-is-not-




Figure 8: Prognostic reach for functioning. (a) Archetype prognostic atlas: two-year functional-remission rate by baseline dominant archetype; the immunometabolic corner is worst (17%) and the low-burden pole best (52%), a transdiagnostic 17%–52% gradient. (b) Held-out incremental value for functioning: the continuous archetype encoding dominates the seven specific axes, any hard tessellation, and the isolated biology axes (operative number of clusters = none). (c) The map and DSM-5 are co-informative—each adds beyond the other, and both together add most—but the individual-level remission AUC gain is only +0.010.

biology dissociation, now translated into outcome. This gradient was a genuine within-diagnosis effect, not a cohort-composition artefact: the rank held inside every cohort (immunometabolic corner → well pole: bipolar 27% → 73%, depression 31% → 72%, schizophrenia 9% → 25%; a cohort-adjusted best-versus-worst odds ratio ≈ 6.3), standardizing to a common cohort mix moved it by only $\approx 4\%$, and the corner-by-cohort interaction was non-significant. The individual-level discrimination gain was nonetheless small (functional-remission AUC +0.010 over a strong diagnosis-plus-severity reference). These are not in tension: a wide spread in *group* remission rates with a small *individual* AUC gain is the expected signature of groups that differ in average risk but overlap substantially at the patient level, layered on an already-strong autoregressive baseline—the map enriches and forecasts at the group level without sharply separating individuals. The functioning signal survived inverse-probability weighting for attrition (+54.4), restriction to either two of the three cohorts (+56.4, +61.8), and a permutation null (which correctly vanished). Finally, against the full set of raw indicators under a matched gradient-boosting classifier, the eight-dimension map was a *sufficient* representation for predicting deterioration (equal AUC) and *near-sufficient* for recovery (raw indicators +0.04 AUC), with 92–97% of the raw recovery signal already contained within the eight factors (Extended Data Fig. F5)—the compression is faithful, not lossy in any axis the map omits. *Clinically*, the atlas is useful for group-level enrichment and monitoring, not as an individual remission calculator.

2.9 A variational re-estimation reproduces the atlas at a fraction of the cost

Because the audit-grade NUTS fit is expensive, we asked whether the same measurement model can be recovered by a much cheaper estimator. Re-estimating the eight-factor map as a generalized linear latent-variable model fit by stochastic variational inference (Methods; Supplement H) reproduces the NUTS atlas on its two load-bearing objects at roughly 100× less cost (≈ 4.5 minutes on a CPU versus hours). The loading map is the same map: loading columns




Figure 9: **A variational re-estimation reproduces the NUTS atlas at $\sim 100\times$ less cost.** (a) Per-factor loading congruence (Tucker φ) between the variational GLLVM and the NUTS copula fit; all eight axes clear the major (0.95) and thin-factor (0.85) bars. (b) Per-patient coordinate agreement (Pearson r); all eight axes exceed 0.90. The full loading scatter ($r = 0.993$), the Φ comparison, and the model and inference details are in Supplement H.

are congruent on all eight factors (Tucker φ 0.96–0.999; G 0.957) and the full item $\times$ factor loading scatter is $r = 0.993$ with no sign flips (Figure 9a). Patients land in the same place: per-patient coordinates correlate with the NUTS coordinates at $r \geq 0.90$ on all eight axes (≥ 0.99 for the G and immunometabolic axes; Figure 9b). The map is also a faithful generative model—9,013 synthetic patients drawn from it match the observed marginals (Kolmogorov–Smirnov median 0.04) and covariance (correlation SRMR 0.078, as for NUTS). The one quantity the variational engine does *not* reproduce is the inter-factor correlation Φ : its structure is correct (the biology $\perp G$ zeros are exact) but the off-diagonals are systematically attenuated by $\sim 21\%$ under the mean-field posterior—a structural property of variational inference, not a tunable defect, which is why NUTS is retained for correlation magnitudes and all credible intervals (Supplement H). The variational engine is therefore a fast operational and generative stand-in for the map and coordinates; a fuller development is left to future work.

3 Discussion

Across three cohorts of severe mental illness, a single missingness-aware Bayesian model places clinical and biological measurements in one transdiagnostic coordinate system and yields an eight-dimension atlas whose most consequential property is a dissociation: immunometabolic load is the domain least entangled with overall functional severity. The cardiometabolic and inflammatory markers resolve as a single immunometabolic axis anchored by body composition (BMI loading ≈ 0.95 , weight 0.88, waist circumference 0.84), with smaller metabolic, inflammatory and cardiovascular contributions (C-reactive protein ≈ 0.37); that axis sits nearly independent of G . Because the one inflammatory marker available here, CRP, is itself substantially adiposity-driven [28, 29], we read this as an adiposity-anchored cardiometabolic axis and do not interpret it as an inflammatory dimension independent of body composition—the IL-6-type signal implicated in mood pathology [30, 31] was not measured here. The axis is nonetheless more than body habitus: excluding the anthropometric indicators and refitting, a coherent cardiometabolic–inflammatory axis survived, highly congruent with the full axis (Tucker $\varphi = 0.90$; Supplement A.6). If biological burden simply tracked how impaired a patient is, it would add little to a clinician’s global impression. The data say the opposite—biological risk rides on an axis the severity picture does not see—and the five archetypal extremes make the consequence concrete. The immunometabolic corner and the severe, clean-biology corner carry comparable functional burden but opposite biological profiles, and they diverge in prognosis (the immunometabolic corner is the worst-prognosis pole and the well pole the best, two-year functional remission spanning 17% to 52% across the five corners). This is the precise property

a biology-aware stratification can exploit: it separates patients who look equally ill but are biologically opposite, and that separation is durable and prognostically meaningful.

This headline is corroborated, in direction, by a large and independent literature, which is why we state it as relative rather than absolute independence. Metabolic burden in severe mental illness is driven by medication, adiposity, age and chronicity more than by diagnosis or symptom severity, is present before chronic severity accrues, and is comparable across schizophrenia, bipolar disorder and depression [14–16, 32, 33]; inflammation marks a subgroup and specific symptoms rather than the severity continuum [34–36]; and the immuno-metabolic-depression programme has independently arrived at a separable biological dimension mapping onto atypical, energy-related symptoms [11–13, 37, 38]. That a completely different method, in different cohorts, converges on a separable biology axis is strong convergent validity. We also state the boundary conditions plainly: case–control inflammatory elevations are real [39–41], a modest severity or duration dose–response for metabolic burden exists [42, 43], metabolic dysfunction does track cognition and functioning in mood disorders [44], and much of the inflammation–symptom association is carried by adiposity rather than inflammation *per se* [28, 45]. Our small immunometabolic–*G* couplings are best read as a quantification of a known qualitative pattern, sitting at its low end because *G* indexes *functional burden*—from which biology is most decoupled—rather than the case–control contrast of being ill. That they survive, and even fall slightly under, adjustment for medication, adiposity and site is the strongest available evidence that the dissociation is not a confound.

The second contribution is structural. On these clinical–biological coordinates the patient space is a graded continuum, not a set of discrete biotypes, and it cuts almost entirely across DSM-5. This aligns with normative-modelling work reporting extensive individual heterogeneity in the same disorders [8–10], and it stands in apparent tension with the brain-based B-SNIP biotypes [17, 18]. The tension is more apparent than real: the two programmes interrogate different measurement spaces, B-SNIP itself found DSM diagnoses to be a severity continuum, and recent syntheses endorse both categorical and dimensional views [19]; clustering choices can also resurface apparent biotypes from continuous data [27]. We therefore claim no biotypes *in this eight-dimensional clinical–biological space*—a claim we make rigorous with a single-Gaussian falsification null (best-partition silhouette 0.140 against a structureless-Gaussian null of 0.137 ± 0.002 , $z = 1.13$) rather than by the absence of an elbow—not that biotypes are absent from all spaces. The five archetypes are corners of this continuum, not clusters within it, and the partition they induce is driven by activation and suicidality rather than by overall severity, which is why it cuts almost orthogonally across DSM-5 (ARI 0.006) while describing the space more tightly than diagnosis does. Anchoring *G* on functioning alone also sidesteps the well-known instability in what a symptom-only general factor means [6, 7, 46], while remaining consistent with the impairment reading of the *p* factor [4, 5].

The remaining contributions are deliberately calibrated. The map persists over two years—durable biology, moving symptoms—which both validates the measurement longitudinally and motivates a clinical division of labour between traits to stratify on and states to monitor. The immunometabolic axis is the single most durable dimension (test–retest ICC 0.91, fully invariant across visits; it remains trait-like, ICC 0.76, even with the anthropometric indicators removed, so the durability is not merely body mass’s stability—Supplement A.6), cognition is also trait-like, while severity and symptoms slide as the population improves: the durable part of the map is precisely its biological part.

This division is itself a design principle—*stratify on what stays put; monitor what moves*. Because the immunometabolic axis is both the durable trait and the dimension least entangled with functional burden, it is the stable axis on which to fix a patient’s position—for instance at the immunometabolic corner (**A2**); general burden is durable only *by rank* (ICC 0.62), with patients largely keeping their relative position even as the cohort improves in absolute terms. The symptom axes, by contrast, slide—population shifts of -0.46 on severity and -0.84 on

suicidality from baseline to two years, against an essentially static +0.04 on biology—and are therefore quantities to track over time rather than to fix a stratum on. We frame this as an interpretive design principle, read off the baseline structure and its forward-scored longitudinal coherence rather than from any validated protocol: what the durability buys is a principled separation of the axes worth stratifying on from those worth monitoring, not an individual treatment rule.

The map predicts future functioning incrementally beyond diagnosis and severity (archetypes $\Delta\text{ELPD} +62.8$ held-out, robust to inverse-probability weighting and permutation), but the gain is modest and group-level—the individual-binary lift is small (AUC +0.010)—it is co-informative with rather than superior to DSM-5, and it is carried by the biological *phenotype* (the immunometabolic archetype corner) rather than by any isolated laboratory axis. “Biology predicts functioning” is therefore a phenotype-level claim: the immunometabolic axis carries a credible adverse direction, but the predictive object is the fuller archetype representation in which it is embedded. That the eight-dimension map is nonetheless a *sufficient* representation of the full raw V0 indicator block (143 items, the representation benchmark’s raw arm) for predicting deterioration, and near-sufficient for recovery (the residual is within-factor item-level compression, not a missing axis), shows the parsimony is faithful rather than lossy. On observational, expert-centre treatment-as-usual the map did not reliably moderate treatment response; with confounding by indication the dominant threat, this analysis can only *bound* moderation rather than establish it, and we report it as an exploratory negative control (Supplement E). It is informative in one direction that matters here: the prognostic signal survives adjustment for the drug classes patients actually received (the immunometabolic carrier attenuates by under 8%), so the forecast is not a treatment proxy. We report these modest and null findings as deliberately as the positive ones; an atlas explicit about the gap between demonstrated scientific validity and undemonstrated individual-level clinical utility is more useful than an overstated one.

In sum, this atlas reframes a clinically actionable hypothesis with measured coordinates: not “which biotype is this patient?” but “where does this patient sit on continuous, partly biological axes that diagnosis and severity do not capture, and how durable is that position?” The immunometabolic axis a patient keeps over time forecasts the functional state they move toward—a group-level, transdiagnostic signal that complements diagnosis, identifies the immunometabolic corner as a concrete enrichment and monitoring target, and points to a specific, testable immunometabolic $\times$ treatment hypothesis for prospective study. Separating biological burden from clinical severity, and showing that the biological part is stable and prognostic, is the step that makes mechanistic stratification in severe mental illness a measurable rather than a rhetorical goal.

Limitations

Four classes of limitation bound these claims. *Internal validity only*: the map is discovered and validated within FACE, with no external-cohort replication, and the invariance partials that exist are documented rather than hidden (G re-anchors in schizophrenia, which lacks the FAST functioning anchor; self-rated mania floors in depression). *Outcomes are scale trajectories, not recorded events*: prognostic endpoints are state transitions defined from repeated functioning and severity scales, not registered relapses or hospitalizations—a deliberate de-scope from event-level prognosis. *Observational treatment*: confounding by indication is the dominant threat (E-values 1.1–1.8), exposures are drug-class rather than protocol, channelled treatments are non-estimable, and the cohort is patients in usual care at expert centres rather than a randomized sample; the treatment analysis can therefore only *bound* moderation, not establish selection. *Measurement coverage and horizon*: two axes are thin (mania has two indicators and is reported as data-limited; substance use is a two-cohort axis), education is 40.8% missing overall (and 74.5% in schizophrenia)—a structured missingness the observed-likelihood design handles but that limits that covariate—developmental risk scores as “state” partly because of recall

noise in childhood-adversity reports rather than true change, attrition is informative (addressed by inverse-probability weighting), and the follow-up window is two years across three visits. Establishing individual-level clinical utility or treatment guidance would require incident-event outcomes, randomized or trial-arm treatment data, and external validation beyond this baseline observational cohort.

4 Methods

This section gives a high-level account of the model and the analysis. The complete mathematical development—the generative model, the observed-data likelihood, the rotation and identification argument, the closed-form covariance identity, the equivalence between the marginalized Bayesian model and full-information maximum likelihood, and the stratification, temporal, prognostic and treatment estimators—is given with proofs in the Supplement (A–E).

4.1 Cohorts, harmonization and the no-imputation data layer

Data were the baseline (visit V0) of three FACE cohorts (Fondation FondaMental expert centres): bipolar disorder, schizophrenia and major depression ($N = 9,013$; BP 6,252, SZ 2,209, DR 552; 21 recruitment sites). A harmonized data dictionary maps each modelled variable to its per-cohort source column, a harmonization rule and per-variable plausibility (“sanity”) bounds; out-of-range values are set to missing (never clipped, never imputed). Deterministic skip-logic decoding sets structural zeros only where a gate is explicitly negative (for example, attempt count = 0 when ever-attempted = 0); this is decoding, not imputation. Each variable carries a likelihood family and a missingness type. Identifiers (patient, cohort, DSM-5 subtype, visit, site) are never modelled as indicators; cohort and subtype enter only as covariates, invariance-grouping variables or validation labels. Of the 201 usable harmonized common variables, 143 form the modelled-indicator pool (the comparator for the raw-indicator prognostic benchmark); the remaining 58 are identifiers, covariates or validation labels and are never factor indicators. The fitted factor map carries 109 of these indicators—88 in the Gaussian-copula continuous block and 21 in the explicit non-Gaussian block—across its eight axes.

4.2 Candidate ontology and soft priors

Ten candidate constructs defined a *soft-prior* ontology rather than a fixed output. Mapping the candidates against the common-variable coverage fixes the eligible set and is itself a primary result: a construct with no common indicator is reported *not testable*, never back-filled with an invented proxy. Each (indicator, factor) cell receives a prior *role*—primary, plausible-cross, unlikely or forbidden—whose default is shrinkage rather than exclusion, so the model can confirm, split, merge, downgrade, reject or declare not-testable any candidate. The formal prior is given in Supplement A.

4.3 The measurement model

The measurement model rests on five core equations; their full derivation, with proofs, is in Supplement A. We posit that a few latent factors generate the many measured quantities, so each patient is summarized by a low-dimensional coordinate. Patient i carries a factor vector of one general functional-burden axis G and $K = 7$ specific axes, given a standardized multivariate-normal prior,

$$f_i = (G_i, D_{i1}, \dots, D_{iK})^\top \sim \mathcal{N}_{K+1}(\mathbf{0}, \Phi), \quad \text{diag}(\Phi) = \mathbf{1}. \quad (4)$$

The unit diagonal of the factor correlation matrix Φ fixes the latent scale; G is the burden axis every indicator may share, while $D_{i1}, \dots, D_{iK}$ are the seven domain-specific axes (cognition,

immunometabolic, sleep, mania/activation, suicidality, developmental risk and substance use), so the map is eight-dimensional: one general axis plus seven specifics. The immunometabolic axis is a single biology factor carrying the cardiometabolic and inflammatory markers together (body-mass index, glycated haemoglobin, lipids, C-reactive protein), with $\text{bmi} \rightarrow \text{immunometabolic} \approx 0.95$ and $\text{crp} \rightarrow \text{immunometabolic} \approx 0.37$. Equation (4) is the formal content of the idea that a patient is *one general burden axis plus specific axes*.

Each indicator j is a linear reflection of those factors through a single predictor,

$$\eta_{ij} = \alpha_j + \lambda_{jG} G_i + \sum_{k=1}^K \lambda_{jk} D_{ik} + \beta_j^\top c_i, \quad (5)$$

with item intercept α_j , a general loading λ_{jG} onto G , specific loadings λ_{jk} , and item-level covariates c_i that shift an indicator’s mean but are never themselves factor indicators. Equation (5) is the bifactor exploratory structural-equation model: bifactor because every indicator may load on G and on its home specific factor; exploratory because small off-home cross-loadings λ_{jk} are admitted under a sparsity-inducing prior rather than forbidden outright. The predictor feeds a per-type observation density—Gaussian or log-Gaussian for continuous and laboratory measures, ordered-logit for ordinal, Bernoulli for binary, negative-binomial for counts—and continuous indicators additionally enter through a rank-based *Gaussian-copula* marginal, so that forcing heterogeneous variables onto a shared scale never manufactures covariance. The likelihood is invariant to rotation of the factors; identification comes from the loading priors, so a prior-free refit that recovers the same structure shows the map is recovered from the data, not imposed (Supplement A).

The off-home cross-loadings are governed by a designed sparsity mechanism that keeps the map mostly simple-structure while letting a few evidence-backed cross-loadings emerge. On each off-home specific $\leftrightarrow$ specific cell we place a regularized (“Finnish”) horseshoe prior [47, 48]: a global shrinkage scale $\tau \sim \text{HalfNormal}(\tau_0)$ pulls the whole set toward zero (the prior’s default belief is *no* cross-loadings), a heavy-tailed local scale lets a single cross-loading the data insist on escape that shrinkage, and a slab c caps the escaped magnitude so a cross-loading stays small (it is not a home loading). The prior is therefore *default-off*, *evidence-on*, *magnitude-capped*. This is exactly what protects the thin specific factors—substance and mania/activation are each defined by very few indicators—from dilution: spurious cells from other factors are crushed to zero by the global shrinkage, so a thin column stays defined only by its sign-anchored home indicators (home loadings are not horseshoed), while a clinically real weak cross-loading can still surface with a credible interval. Substance is additionally pinned orthogonal to the correlated block, its cross-factor correlations being non-identifiable. A continuous sparse-ESEM validation arm that frees every off-home cell confirms the structure is *recovered from the data, not imposed*: roughly 83% of the freed cells shrink to ≈ 0 , and the credible remainder is folded back into the full mixed map, which retains three small cross-loadings—childhood adversity (CTQ-37, -0.094), sleep latency (PSQI-latency, $+0.057$) and daytime dysfunction (PSQI-daytime, -0.070) onto cognition, each with a 95% interval excluding zero. Given total freedom the model reproduces known clinical cross-talk and nothing spurious, which is positive evidence the map is well specified (Supplement A).

Given the latent scores the indicators are conditionally independent (local independence: once the factors are fixed, only item-specific noise remains, a diagonal residual covariance Ψ). Equations (4)–(5) then generate a structured covariance among the indicators,

$$\Sigma = \Lambda \Phi \Lambda^\top + \Psi = \underbrace{\lambda_G \lambda_G^\top}_{\text{general: rank one}} + \underbrace{\Lambda_D \Phi_D \Lambda_D^\top}_{\text{specific factors}} + \underbrace{\Psi}_{\text{private noise}}, \quad (6)$$

where $\Lambda = [\lambda_G \mid \Lambda_D]$ collects the loadings and Φ_D is the specific-factor correlation block. Equation (6) is the generative claim in one line— G leaves a rank-one footprint that touches

every item, the specific factors contribute a block, and Ψ absorbs what no factor explains—and it is also why imputation is forbidden: a filled-in cell would inject covariance that (6) would misread as a factor.

Accordingly the model is fit to each patient’s *observed cells only*. With O_i the set of indicators observed for patient i and F_j the density of indicator j , the observed-data log-likelihood is

$$\log p(X_{\text{obs}} | \Theta) = \sum_{i=1}^N \sum_{j \in O_i} \log F_j(X_{ij} | \eta_{ij}, \theta_j). \quad (7)$$

A missing cell contributes no term and no completed $N \times J$ matrix is ever formed; deterministic skip-logic decodes structural zeros as *observed* values, not imputed ones. For the multivariate-normal block this observed-cell sum is exactly the full-information maximum-likelihood objective for incomplete data, obtained by *marginalizing* the missing coordinates rather than filling them (Supplement A). Equation (7) is the no-imputation invariant on which every downstream analysis rests.

Two factor covariances are estimated. The primary *bifactor* fit holds G orthogonal to the specific axes (Φ block-diagonal); this is the natural null for the biology question. A *correlated-G* arm then unzips that zero block,

$$\Phi = \begin{pmatrix} 1 & \phi_{Gd}^\top \\ \phi_{Gd} & \Phi_D \end{pmatrix}, \quad \phi_{Gd} = \text{Corr}(G, D) \in [-1, 1]^K, \quad (8)$$

turning “biology is separable from severity” from a built-in constraint into the *measured* vector ϕ_{Gd} : a domain is severity-entangled when its entry is large and separable when it is near zero (immunometabolic ≈ 0.10 , versus cognition 0.39 and sleep 0.42 in the same freely-correlated model—so the separation is a cross-domain ordering the model is free to violate, not a consequence of the bifactor zero block). Reporting the bifactor general loadings $|\lambda_{jG}|$ and the correlated- G correlations ϕ_{Gd} together—they agree on the ordering—is what makes the separation a measurable estimand rather than an artefact of the orthogonality constraint (Figure 4; Supplement A). The biology- G result is reported unadjusted and under a covariate-adjustment ladder (age, sex, education, site, antipsychotic exposure, adiposity) implemented by residualizing each indicator before the factor model.

4.4 Estimation, confirmation, invariance and scoring

The model is fit globally to the full sample ($N = 9,013$) under per-patient cohort weights $w_i = N/(K n_{c(i)})$, where $K = 3$ is the number of cohorts and $n_{c(i)}$ the size of patient i ’s cohort, so that each cohort contributes equal total weight N/K while every patient is retained and $\sum_i w_i = N$. Balancing by subsampling is rejected because matching the smallest cohort would discard $\sim 92\%$ of the bipolar sample; weighting equalizes cohort influence without that loss. This is a composite-likelihood weighting: the point estimate is the balanced transdiagnostic estimand, and because the resulting posterior standard deviations are order-correct rather than fully calibrated we validate them by cross-seed congruence. The map’s insensitivity to the weighting is quantified by the robustness battery below, one arm of which refits under inverse-cohort weighting against the unweighted full-sample fit (minimum Tucker $\varphi = 0.96$). The global posterior was reached by continuation: a well-conditioned sub-model was fit and then deformed toward the full target one source of difficulty at a time, each converged posterior warm-starting the next, and *only the global fit is interpreted*. The Gaussian block is marginalized analytically (Woodbury identity and matrix-determinant lemma, grouped by missingness pattern), which removes the per-patient latent funnel and makes full-sample estimation tractable on a workstation (Supplement A). Sampling used the No-U-Turn sampler [49] in PyMC [50] with the NumPyro/JAX backend [51]; a stage advanced only on standard convergence criteria (acceptable $\hat{R}$, adequate effective sample

size, zero divergences, no Heywood cases, acceptable posterior-predictive behaviour), with a cross-seed stability guard for the joint mixed fit. The global 8-factor fit converges cleanly ($\hat{R} = 1.03$, zero divergences). Identification does not rest on information criteria, which favour bifactor over correlated-factor structures for reasons of functional form that are independent of which model generated the data [23, 24, 52]. We rely instead on three prior-free checks performed in-engine. First, an independent flat-prior refit (identification constraints only, fresh seed, no warm start) reproduced the loadings and factor correlations to three decimals (Tucker congruence $\varphi = 1.00$), so the structure is driven by the data rather than by the soft loading priors. Second, the correlated- G arm (Eq. 8) tests the G -specific orthogonality directly rather than imposing it. Third, posterior-predictive checks gave a Bayesian standardized root-mean-square residual ≈ 0.07 for the continuous block and reproduced 21/22 non-Gaussian indicators within the 90% predictive interval. As a secondary check WAIC and PSIS-LOO [53, 54] also ranked the bifactor above unidimensional and correlated-factor alternatives, but for the reason above we treat this as confirmatory only. Because the marginalized Bayesian model and full-information maximum likelihood optimize the same observed-data objective (Supplement A), a standalone maximum-likelihood arm adds no independent evidence. Measurement invariance across BP/SZ/DR was tested by multi-group loadings and Bayesian differential item functioning, and robustness by leave-one-cohort-out congruence, diagnosis-balanced subsampling, site cluster-bootstrap and inverse-cohort weighting (minimum Tucker $\varphi = 0.96$). Each patient received, per axis, a posterior mean, standard deviation and credible interval, the number of observed indicators and a reliability tier, so that all downstream analyses propagate uncertainty rather than treating coordinates as equally measured.

4.5 A fast variational re-estimation

As an efficiency-oriented alternative to NUTS, we re-estimate the same measurement model—the bifactor/ESEM atlas of Eqs. (4)–(7), identical ontology and mixed likelihood—as a generalized linear latent-variable model [55] fit by stochastic variational inference [56, 57] in PyTorch [58]. Each patient’s coordinate receives a variational posterior $q_i(f_i) = \mathcal{N}(\mu_i, \Sigma_i)$, and the model is fit by minimizing the negative evidence lower bound (observed cells only; the soft loading prior enters as a penalty; reparameterized gradients [59]), with the global parameters as maximum-a-posteriori estimates. This trades NUTS’s exact posterior sampling for optimization within an approximate family: it is $\sim 100\times$ faster (≈ 4.5 minutes at full N on a CPU, versus hours) and reproduces the NUTS loading map and patient coordinates closely, at the cost of attenuated inter-factor correlations under a mean-field posterior—so it serves as a fast operational and generative engine while NUTS remains the authority for correlation magnitudes and credible intervals. The full model, inference and head-to-head comparison are in Supplement H.

4.6 Stratification, temporal coherence, prognosis and treatment

On the fitted coordinates $\{x_i, S_i\}$ —each patient’s posterior-mean position $x_i = \hat{f}_i \in \mathbb{R}^8$ together with its posterior covariance S_i —structure was tested *before* any clustering by a multi-statistic battery computed over the posterior draws, anchored by a *single-Gaussian falsification null*: synthetic data are drawn from one Gaussian with the empirical mean and covariance of the coordinates (structureless by construction), the identical pipeline is run on real and null data, and we ask whether the real cloud’s best partition separates patients any better than the null’s. It does not (best-partition silhouette 0.140 versus a null 0.137 ± 0.002 , $z = 1.13$, n.s.; Supplement B), so the verdict is a graded *continuum*, not discrete biotypes. The primary representations are therefore the continuous coordinates themselves and a soft archetype simplex, not any hard partition. Archetypal analysis [60] writes each patient as a convex blend of $A = 5$ extreme phenotypes pinned to the convex hull of the cloud (the barycentric representation reported in Eq. 3), each archetype itself a convex combination of patients; the simplex weights

$w_i \in \Delta^{A-1}$ are a patient’s soft archetype weighting, so the continuum is *read* through extremes rather than *cut* into kinds. The number of corners ($A = 5$) was set by cross-seed stability rather than parsimony—the largest A holding cross-seed Tucker congruence ≥ 0.8 , with a clean stability cliff at $A = 6$ ($0.979 \rightarrow 0.436$)—with the full optimization $z_a = \sum_j b_{ja} f_j$ and the draw-wise uncertainty propagation given in Supplement B. A measurement-error Gaussian mixture (extreme deconvolution [61], each patient carrying its own S_i) supplies a nested family of soft tessellations with no privileged number of regions, the operative granularity deferred to incremental predictive validity downstream (Supplement B). Each downstream analysis then treats the measurement model as *fixed*—reusing or re-scoring coordinates without re-discovering the map—and propagates each coordinate’s posterior uncertainty S_i rather than treating the scores as exact. Temporal coherence scored the follow-up visits on the *fixed* measurement model (observed cells, uncertainty propagated), tested longitudinal invariance, and decomposed trait from state with a measurement-error random-intercept model that plugs the known baseline measurement variance and includes visit fixed effects, yielding an individual rank-stability intraclass correlation and a population-level shift (Supplement C). Prognosis compared nested models—nuisance plus baseline outcome plus severity, then adding DSM-5 subtype, the map, or both—using an errors-in-variables Bayesian generalized linear model that propagates each coordinate’s posterior standard deviation, scored by held-out expected log predictive density and, for binary endpoints, by change in AUC; robustness used inverse-probability weighting, restriction to reliable axes and a permutation null, and a separate benchmark compared the map against all 143 raw indicators under a matched gradient-boosting classifier (Supplement D). Treatment exposures, harmonized to drug classes, were analysed as an exploratory negative control by an identification-first causal pipeline—common-support overlap gate, stabilized inverse-probability-of-treatment weighting on a propensity that includes the eight coordinates [62, 63], doubly-robust errors-in-variables moderation, an E-value sensitivity analysis [64] and a minimum-detectable-effect calculation that makes any null a *bounded* one; the full analysis is reported in the supplement (Supplement E).

4.7 Compute and reproducibility

The pipeline is configuration-first with fixed seeds. Raw per-cohort data are confidential and never leave their governed environment, while the code, the harmonized dictionary and the regenerated aggregate results are shareable; every reported number is reproducible from the numbered pipeline, and the sample-characteristics table is regenerated by an aggregate-only script. Estimation ran on a workstation (Apple M4 Pro, 24 GB; no GPU).

Data availability

The FACE database is the property of Fondation FondaMental; per-patient cohort data are confidential and governed by FondaMental’s data-access procedures, to which qualified researchers may apply. No per-patient data are reproduced in this article. Aggregate parameters, derived structure and all figure/table source values are available in the project’s regenerated aggregate outputs.

Code availability

The full analysis pipeline (data harmonization, the Bayesian measurement engine, stratification, temporal, prognosis and treatment modules), the harmonized variable dictionary, and configuration files are available at [\[TODO: repository URL / DOI\]](#). Every reported quantity is reproducible from the numbered pipeline; per-patient data are never required to regenerate the aggregate results, tables and figures.

Ethics

The FACE cohorts were approved by the relevant institutional review boards and all participants provided written informed consent. [TODO: Insert exact IRB/ethics committee names, approval numbers, and trial/cohort registration identifiers.]

Author contributions

[TODO: Complete per ICMJE / journal policy and FACE–FondaMental authorship conventions, including cohort investigators.]

Funding

[TODO: Fondation FondaMental and cohort funders to be specified.]

Competing interests

[TODO: Declare per journal policy.]

Acknowledgements

We thank the FACE–FondaMental Expert Centre network, participating clinicians and, above all, the participants. [TODO: Complete.]

Supplementary materials — derivations, proofs, diagnostics and reproducible tables

A The measurement model: full derivation

This annex develops the global measurement model from first principles: the generative factor model and its defining local-independence assumption (A.1); the mixed-likelihood observation model and the Gaussian-copula marginal (A.2); the observed-data likelihood and why missingness is handled by marginalization rather than imputation (A.3); the soft-prior loadings, the regularized-horseshoe sparse-ESEM that lets a few evidence-backed cross-loadings emerge, and the rotation problem they resolve (A.4); the closed-form covariance identity that carries the headline estimand (A.5); the Woodbury evaluation and its exact equivalence with full-information maximum likelihood (A.7); and the staged estimation and in-engine confirmation (A.8). Propositions are proved; everything reported in the main text is an evaluation of the objects defined here.

A.1 Generative model and local independence

Index patients $i = 1, \dots, N$, indicators $j = 1, \dots, J$, specific factors $k = 1, \dots, K$ and one general factor G . A few latent causes generate the many measured quantities: each patient carries a factor vector $f_i = (G_i, D_{i1}, \dots, D_{iK})^\top \in \mathbb{R}^{K+1}$ with

$$f_i \sim \mathcal{N}_{K+1}(\mathbf{0}, \Phi), \quad \text{diag}(\Phi) = \mathbf{1}, \quad (\text{A9})$$

the factor *correlation* matrix Φ carrying a unit diagonal to fix the latent scale. Each indicator is a linear reflection of the factors plus item-specific noise, through a single linear predictor

$$\eta_{ij} = \alpha_j + \lambda_{jG} G_i + \sum_{k=1}^K \lambda_{jk} D_{ik} + \beta_j^\top c_i, \quad (\text{A10})$$

with item intercept α_j , general loading λ_{jG} , specific loadings λ_{jk} (row j of the loading matrix $\Lambda = [\lambda_G \mid \Lambda_D] \in \mathbb{R}^{J \times (K+1)}$), and item-level covariates c_i that adjust an indicator’s *mean* but are never factor indicators. Stacking one patient’s items, with the general column separated from the specific block,

$$x_i = \mu_i + \underbrace{\lambda_G G_i}_{\text{general}} + \underbrace{\Lambda_D D_i}_{\text{specifics}} + \varepsilon_i, \quad \mu_i = \alpha + B c_i, \quad \varepsilon_i \sim \mathcal{N}(\mathbf{0}, \Psi), \quad (\text{A11})$$

with diagonal residual covariance $\Psi = \text{diag}(\sigma_1^2, \dots, \sigma_J^2)$.

Local independence (the defining assumption). Given a patient’s latent scores, the items are conditionally independent,

$$p(x_i \mid f_i, \Theta) = \prod_{j \in O_i} p(x_{ij} \mid f_i, \Theta), \quad (\text{A12})$$

i.e. the factors are the only reason items covary. In the Gaussian linear model this is *identical* to “ Ψ is diagonal”: local independence and diagonal residual covariance are two faces of one assumption—that the factors explain away all shared covariance. (It is also why filling missing cells is forbidden: an invented value injects covariance the model would mistake for a factor.) Because f_i is unobserved and the likelihood must depend on Θ alone, we integrate it out and multiply across independent patients,

$$L(\Theta) = \prod_{i=1}^N p(x_i \mid \Theta) = \prod_{i=1}^N \int \left[\prod_{j \in O_i} p(x_{ij} \mid f_i, \Theta) \right] \mathcal{N}(f_i; \mathbf{0}, \Phi) \, df_i. \quad (\text{A13})$$

This is the complete first-principles likelihood: a product over patients (independent units), a product over items (local independence), and an integral that removes the latent (the sum rule). Equation (A10) is a bifactor exploratory structural-equation model [20, 21]—bifactor because every indicator may load on G *and* on its home specific factor; exploratory because an indicator is *allowed* small cross-loadings on plausibly related factors, governed by a sparsity-inducing prior (A.4) that keeps the map mostly simple-structure yet lets a few evidence-backed cross-loadings emerge, rather than forbidding them outright as in confirmatory factor analysis. The fitted map has $K+1 = 8$ factors: the general factor G (overall burden) and seven specific axes—cognition, immunometabolic, sleep, mania/activation, suicidality, developmental-risk and substance.

A.2 Mixed-likelihood observation model and the Gaussian copula

Coercing variables onto a common scale would manufacture covariance and bias the very factors being estimated. Each indicator instead keeps the observation density appropriate to its type, all sharing the linear predictor (A10):

$$\text{continuous / } z\text{-score: } X_{ij} \sim \mathcal{N}(\eta_{ij}, \sigma_j^2), \quad (\text{A14})$$

$$\text{skewed laboratory: } \log X_{ij} \sim \mathcal{N}(\eta_{ij}, \sigma_j^2), \quad (\text{A15})$$

$$\text{ordinal } (C_j \text{ levels}): \Pr(X_{ij} \leq c) = \text{logit}^{-1}(\tau_{jc} - \eta_{ij}), \quad \tau_{j1} < \dots < \tau_{j,C_j-1}, \quad (\text{A16})$$

$$\text{binary: } X_{ij} \sim \text{Bernoulli}(\text{logit}^{-1}(\eta_{ij})), \quad (\text{A17})$$

$$\text{count: } X_{ij} \sim \text{NegBinom}(\mu_{ij} = e^{\eta_{ij}}, \phi_j). \quad (\text{A18})$$

The Gaussian-copula marginal (reported map). The reported map replaces the raw Gaussian likelihood on the continuous block with a semiparametric *Gaussian-copula* model: each continuous indicator j is mapped to a latent Gaussian score by its frozen empirical distribution function,

$$z_{ij} = \Phi_{\mathcal{N}}^{-1}(\hat{F}_j(X_{ij})), \quad (\text{A19})$$

and the factor model of A.1 is fit on z_{ij} . This makes the continuous block invariant to monotone re-scaling and robust to non-normal margins (heavy tails, skew, bounded scales) while leaving the latent dependence structure—the loadings and factor correlations that the scientific claims live in—unchanged. Because the rank transform (A19) is monotone, it alters none of the identification arguments below. Fitting the same structure under the plain Gaussian likelihood recovers the same loadings and correlations to within seed variability; that the central findings hold under both is reported in the main text as a robustness result. Writing $F_j(\cdot)$ for indicator j ’s density and $O_i \subseteq \{1, \dots, J\}$ for its *observed* indicators, the observed-data log-likelihood sums over observed cells only,

$$\log p(X_{\text{obs}} \mid \Theta) = \sum_{i=1}^N \sum_{j \in O_i} \log F_j(X_{ij} \mid \eta_{ij}, \theta_j), \quad (\text{A20})$$

so a missing cell contributes no term and no completed $N \times J$ matrix is ever formed; deterministic skip-logic decodes structural zeros as *observed* values.

A.3 Missing data is the marginal property, not imputation

Proposition 1 (Marginalization). *For patient i with observed continuous set $C_i \subseteq O_i$, the model-implied distribution of the observed sub-vector is obtained by deleting the unobserved coordinates from (μ, Σ) :*

$$X_{i,C_i} \sim \mathcal{N}(\mu_{i,C_i}, \Sigma_{C_i}), \quad \Sigma_{C_i} = \Lambda_{C_i} \Phi \Lambda_{C_i}^\top + \Psi_{C_i}, \quad (\text{A21})$$

with Λ_{C_i} the observed-row loadings and $\Psi_{C_i} = \text{diag}(\sigma_j^2)_{j \in C_i}$.

Proof. A foundational property of the multivariate normal is that marginals are read off by deleting coordinates: if $x \sim \mathcal{N}(\mu, \Sigma)$ then any sub-vector x_C is $\mathcal{N}(\mu_C, \Sigma_{CC})$. Apply this to $x_i \sim \mathcal{N}(\mu_i, \Sigma)$ with $\Sigma = \Lambda\Phi\Lambda^\top + \Psi$ (Proposition 3); the submatrix Σ_{CC} retains exactly the observed rows/columns of Λ and the observed diagonal of Ψ , giving (A21). $\square$

Hence “integrating out” the missing cells is not an extra step but the marginal property itself; summing $\log \mathcal{N}(\cdot)$ over each patient’s observed sub-vector is precisely the full-information maximum-likelihood (FIML) objective for incomplete multivariate-normal data. This is the formal justification of the no-imputation invariant: any completed matrix would inject between-cell covariance that (A12) would misattribute to a latent factor.

A.4 Soft-prior loadings, sparse-ESEM cross-loadings and the rotation problem

Theory enters through the prior on each loading. Each (indicator, factor) cell is assigned a role $\rho(j, k)$ —one of PRIMARY, plausible CROSS, UNLIKELY, or FORBIDDEN—and the prior on that loading is

$$\lambda_{jk} \mid \rho(j, k) \sim \begin{cases} \mathcal{N}_+(0.70, 0.25^2) & \text{PRIMARY (HOME CELL),} \\ \text{regularized horseshoe (A23)} & \text{PLAUSIBLE CROSS,} \\ \mathcal{N}(0, 0.05^2) & \text{UNLIKELY,} \\ \delta_0 & \text{FORBIDDEN,} \end{cases} \quad (\text{A22})$$

where $\mathcal{N}_+$ truncates to $(0, \infty)$ and δ_0 fixes the loading at zero. Remaining priors are weakly informative: $\alpha_j \sim \mathcal{N}(0, 1.5^2)$; $\sigma_j = 0.05 + \text{HalfNormal}(1)$ (the floor guards against a Heywood case); ordered-logit cutpoints τ_j . ordered; the negative-binomial concentration half-normal; and $\Phi \sim \text{LKJ}(\eta=2)$, whose density $\propto \det(\Phi)^{\eta-1}$ gently favours weakly correlated factors [65]. Latent scores use a non-centred parameterization to defuse the hierarchical funnel.

The cross-loading prior: a regularized horseshoe (default-off, evidence-on, magnitude-capped). A measurement map must choose how each indicator relates to factors other than its home factor. Both naive choices fail here. *Hard-zero* (rigid CFA) fixes every off-home specific $\leftrightarrow$ specific cell at exactly 0: clean and identified, but it forbids an instrument from loading weakly on a second axis even when the data support it. *Free / soft-Normal* (naive ESEM) frees every off-home cell at $\mathcal{N}(0, 0.05^2)$: this floods every factor’s column with weak cross-loadings, and a *thin* factor—one defined by very few indicators (substance, four; mania, two)—loses its identity as the flood of spurious cross-loadings into its column overwhelms its handful of genuine home loadings, going multimodal and poisoning convergence. The map needs the middle ground: let an instrument carry a small loading on several axes when the data genuinely support it, without collapsing a low-instrument factor. The classic tool for “mostly zero, but let real signal through” is a sparsity-inducing prior, and the right one is the regularized (“Finnish”) horseshoe [47, 48]. For each off-home specific $\leftrightarrow$ specific cross-loading λ_j (the PLAUSIBLE CROSS cells),

$$\lambda_j = z_j \tau \tilde{\eta}_j, \quad \tilde{\eta}_j = \sqrt{\frac{c^2 \eta_j^2}{c^2 + \tau^2 \eta_j^2}}, \quad z_j \sim \mathcal{N}(0, 1), \quad \eta_j \sim \text{HalfCauchy}(1), \quad \tau \sim \text{HalfNormal}(\tau_0), \quad (\text{A23})$$

with $\tau_0 = 0.05$ and slab $c = 0.30$, fit non-centred (z separate from the scales). Three pieces, each doing one job. The *global shrinkage* τ pulls the whole set of cross-loadings toward 0—the prior’s default belief is “no cross-loadings.” The *local shrinkage* η_j has heavy (Cauchy) tails, so a single cross-loading the data insist on can take a large η_j and escape the global shrinkage—the sharp spike at 0 with heavy tails that defines the horseshoe. The *slab* c caps the escape ($\tilde{\eta}_j \rightarrow c/\tau$ for

large η_j), so an escaped loading stays near $|\lambda| \approx c$ (cross-loadings remain *small*—they are not home loadings) and the funnel is tamed so the sampler mixes.

This prior is exactly matched to the scientific situation. Dilution happens when indicators from *other* factors acquire spurious cross-loadings *into* a thin factor’s column; under the horseshoe those spurious cells get small local η and are crushed to ≈ 0 by the global τ , so the thin column stays defined only by its home indicators (which keep their separate, sign-anchored positive prior and are *not* horseshoed). A genuine weak cross-loading—clinically real but carried by few instruments—can still emerge through the heavy tails and be reported with a credible interval. The prior is therefore default-off, evidence-on, magnitude-capped: the honest way to handle a weak signal in a factor with very few indicators. A map whose credible cross-loadings are few, small and clinically sensible is positive evidence the structure is well-specified, not an artefact of rigid zeros.

The data-supported cross-loadings. A continuous sparse-ESEM validator freed all off-home specific $\leftrightarrow$ specific cells simultaneously and let the horseshoe select: $\approx 83\%$ shrank to ≈ 0 (median 0.002), so the simple structure is *recovered from the data, not imposed*, while the heavy tails let a handful of small, well-converged, clinically sensible cells survive; the thin factors stayed intact (substance home $|\lambda|$ 0.585, mania 0.484). Folding the survivors into the full mixed map—which arbitrates them under the correlated factor block—keeps three credible cross-loadings, all childhood-trauma / sleep items loading on cognition, each with a 95% credible interval excluding zero:

cross-loading	median	95% CI	reading
CTQ-37 $\rightarrow$ cognition	−0.094	[−0.134, −0.053]	childhood adversity $\leftrightarrow$ cognition
PSQI latency $\rightarrow$ cognition	+0.057	[+0.011, +0.102]	sleep $\leftrightarrow$ cognition
PSQI daytime $\rightarrow$ cognition	−0.070	[−0.122, −0.020]	sleepiness $\leftrightarrow$ cognition

The candidate mania cross-loadings widened to non-credible and were dropped: a mania $\leftrightarrow$ sleep relation belongs in the factor correlation ($\phi_{\text{mania, sleep}} \approx 0.24$), not an item cross-loading on a two-indicator factor. The headline of this exercise is the validation: given total freedom the model reproduces known clinical cross-talk and nothing spurious—positive evidence the 8-factor map is well-specified.

Proposition 2 (Rotation invariance of the likelihood). *For any invertible $M \in \mathbb{R}^{(K+1) \times (K+1)}$, the reparameterization $\Lambda^* = \Lambda M$, $\Phi^* = M^{-1} \Phi M^{-\top}$, $f^* = M^{-1} f$ leaves the implied covariance, and hence the Gaussian likelihood, unchanged:*

$$\Lambda^* \Phi^* \Lambda^{*\top} + \Psi = \Lambda M (M^{-1} \Phi M^{-\top}) M^\top \Lambda^\top + \Psi = \Lambda \Phi \Lambda^\top + \Psi = \Sigma. \quad (\text{A24})$$

Proof. Immediate from $MM^{-1} = I$: the inner $M(M^{-1} \Phi M^{-\top}) M^\top$ telescopes to Φ . Since x_i is determined in distribution by (μ_i, Σ) and Σ is invariant, so is the likelihood. $\square$

A factor model is therefore not identified from the likelihood alone; confirmatory factor analysis breaks this by fixing loadings to exactly zero, whereas the soft-prior model breaks it with the prior. The likelihood is rotation-invariant but the prior is *not*: a rotation moves a loading the prior wanted near zero away from zero, lowering $p(\Lambda)$, so the posterior $p(\Lambda \mid X) \propto p(X \mid \Lambda) p(\Lambda)$ concentrates at the orientation where small loadings align with the soft-zero priors and home loadings are positive and substantial; the positive truncation removes the residual sign-flip.

Corollary 1 (Flat-prior confirmation). *Under a flat loading prior (identification constraints only) the posterior mode equals the maximum-likelihood (FIML) solution. Hence a prior-free refit landing on the same loadings demonstrates that the structure is recovered from the data, not manufactured by the priors. Empirically this refit reproduced the map to Tucker congruence $\varphi = 1.00$.*

A.5 The covariance identity and the headline estimand

Proposition 3 (Bifactor covariance). *Under (A11) with $\text{Var}(G_i) = 1$, $\text{Cov}(D_i) = \Phi_D$, $\mathbb{E}[\varepsilon_i \varepsilon_i^\top] = \Psi$ diagonal, and the bifactor orthogonality $\mathbb{E}[G_i D_i^\top] = \mathbf{0}$ together with $(G_i, D_i) \perp \varepsilon_i$,*

$$\Sigma = \text{Cov}(x_i) = \underbrace{\lambda_G \lambda_G^\top}_{\text{general: rank one}} + \underbrace{\Lambda_D \Phi_D \Lambda_D^\top}_{\text{specific factors}} + \underbrace{\Psi}_{\text{private noise}} = \Lambda \Phi \Lambda^\top + \Psi, \quad \Phi = \begin{pmatrix} 1 & \mathbf{0}^\top \\ \mathbf{0} & \Phi_D \end{pmatrix}. \quad (\text{A25})$$

Proof. Expand $\Sigma = \mathbb{E}[(x_i - \mu_i)(x_i - \mu_i)^\top]$ with $x_i - \mu_i = \lambda_G G_i + \Lambda_D D_i + \varepsilon_i$. The nine resulting terms include cross-terms $\lambda_G \mathbb{E}[G_i D_i^\top] \Lambda_D^\top$ and $\mathbb{E}[(\lambda_G G_i + \Lambda_D D_i) \varepsilon_i^\top]$; the first vanishes by $\mathbb{E}[G_i D_i^\top] = \mathbf{0}$ and the second by independence of the factors and the noise. The survivors are $\lambda_G \lambda_G^\top \text{Var}(G_i)$, $\Lambda_D \Phi_D \Lambda_D^\top$ and Ψ , giving (A25); the zero off-diagonal block of Φ encodes $G \perp \text{specific}$. $\square$

The general factor leaves a *rank-one* footprint: every pair of items covaries through G by exactly $\lambda_{jG} \lambda_{j'G}$ —one shared source touching all items.

The headline lives in the off-diagonal block. The correlated- G model *unzips* the zero block, freeing $\phi_{Gd} = \text{Corr}(G, D)$:

$$\Phi = \begin{pmatrix} 1 & \phi_{Gd}^\top \\ \phi_{Gd} & \Phi_D \end{pmatrix}, \quad \Sigma = \lambda_G \lambda_G^\top + \underbrace{\lambda_G \phi_{Gd}^\top \Lambda_D^\top + \Lambda_D \phi_{Gd} \lambda_G^\top}_{\text{the terms orthogonality removed}} + \Lambda_D \Phi_D \Lambda_D^\top + \Psi. \quad (\text{A26})$$

The vector ϕ_{Gd} —the correlation of each specific factor with general burden—is the headline estimand of the enterprise: “biology is the least severity-entangled domain” is the statement that the immunometabolic entry is near zero while cognition and sleep are far larger (0.39 and 0.42). The bifactor orthogonality is the *null*; the correlated- G arm measures the departure from it; and the bifactor direct loading $|\lambda_{jG}|$ and the correlation ϕ_{Gd} agree on the ordering—reporting both is what turns “biology is separable” from a constraint artefact into a finding.

The immunometabolic factor is anchored by body composition and graded across metabolic, inflammatory and cardiovascular markers. Table A4 lists all 37 primary loadings with 95% credible intervals, every one excluding zero: the three body-size indicators (BMI 0.95, weight 0.88, waist 0.84) are the strongest, while the metabolic, inflammatory (CRP 0.37) and cardiovascular markers contribute smaller but credible loadings. The axis is therefore adiposity-anchored rather than inflammation-led; a sensitivity refit excluding the three anthropometric indicators, reported below, tests whether a coherent metabolic-inflammatory-cardiovascular axis survives once the body-size anchors are removed.

A.6 Sensitivity: the axis without its anthropometric anchors

To test whether the immunometabolic factor is merely a body-habitus axis, we refit the full eight-factor map after *excluding* the three anthropometric indicators (BMI, weight, waist circumference), leaving the metabolic, inflammatory and cardiovascular markers to define the axis on their own (balanced $N = 2,000$, otherwise identical specification; 0 divergences, $\hat{R} = 1.09$, so magnitudes are read as approximate). The axis did not collapse. All 34 remaining indicators loaded with 95% credible intervals excluding zero, and the column re-anchored on blood pressure (systolic and diastolic, standing and supine, 0.63–0.67), heart rate (0.44–0.48), the total/HDL cholesterol ratio (0.44), triglycerides (0.40) and inflammatory markers (white-cell count 0.36, C-reactive protein 0.32, neutrophils 0.28)—a coherent cardiometabolic-inflammatory syndrome rather than a residue. Against the canonical axis the loading column was highly congruent on the shared items (Tucker $\varphi = 0.90$), and the factor remained nearly uncorrelated with G and the

symptom axes (specific-specific $|\phi| \leq 0.02$, apart from the known immunometabolic–substance coupling ≈ 0.20). The body-size indicators are therefore the strongest *anchors* of the axis, not its content: removing them redistributes the loading onto the cardiovascular and metabolic markers without dissolving the dimension. Because the one inflammatory marker available, CRP, is itself substantially adiposity-driven [28–30], we name the axis cardiometabolic and adiposity-anchored and do not claim an inflammatory dimension independent of body composition.

The axis is also durable without its anthropometric anchors. Scored forward onto the 12- and 24-month visits under the fixed excluded model (the temporal protocol of C), the immunometabolic coordinate remained trait-like: a measurement-error test–retest intraclass correlation of 0.76 (94% CI [0.74, 0.78]), with an essentially static population trajectory (shift ≈ 0). This is below the full-axis 0.91—body mass is indeed the most stable single component—but still comfortably in the trait range, so the 0.91 durability is not merely the test–retest stability of body size but a property of the broader cardiometabolic–inflammatory constellation.

A.7 Woodbury evaluation and equivalence with FIML

Evaluating (A21) naively costs $\mathcal{O}(|C_i|^3)$ per patient. The Woodbury identity and matrix-determinant lemma reduce it to the $(K+1)$ -dimensional factor space.

Lemma 1 (Woodbury reduction). *With Ψ_{C_i} diagonal and Λ_{C_i} the observed-row loadings,*

$$\Sigma_{C_i}^{-1} = \Psi_{C_i}^{-1} - \Psi_{C_i}^{-1} \Lambda_{C_i} (\Phi^{-1} + \Lambda_{C_i}^\top \Psi_{C_i}^{-1} \Lambda_{C_i})^{-1} \Lambda_{C_i}^\top \Psi_{C_i}^{-1}, \quad (\text{A27})$$

$$\log |\Sigma_{C_i}| = \log |\Phi^{-1} + \Lambda_{C_i}^\top \Psi_{C_i}^{-1} \Lambda_{C_i}| + \log |\Phi| + \sum_{j \in C_i} \log \sigma_j^2. \quad (\text{A28})$$

The inner solve is $\mathcal{O}((K+1)^3)$, independent of $|C_i|$; patients sharing a missingness pattern share the inner $(K+1) \times (K+1)$ factorization, computed once per pattern. This removes the per-patient latent funnel for the continuous block and makes full-sample ($N = 9,013$) estimation tractable on a workstation; the non-Gaussian block retains explicit per-patient latents.

Proposition 4 (Bayes/FIML equivalence). *The marginalized Bayesian observed-data model and FIML optimize the same observed-data likelihood and differ only by the priors (A22). In particular (Corollary 1) the flat-prior MAP equals the FIML estimate, so a standalone FIML estimator adds no independent evidence and confirmation is done in-engine.*

Proof. By Proposition 1 the per-patient marginal density is the incomplete-data multivariate-normal density on C_i , whose log summed over patients is the FIML objective. The posterior is this likelihood times the prior; with a flat prior the log-posterior equals the log-likelihood up to a constant, so its mode is the FIML maximizer. $\square$

Factor scores. Each patient’s coordinates are the expected a posteriori (EAP) factor scores from their observed cells; for an all-continuous pattern,

$$\hat{f}_i = \mathbb{E}[f_i | X_{i,C_i}] = \Phi \Lambda_{C_i}^\top \Sigma_{C_i}^{-1} (X_{i,C_i} - \mu_{i,C_i}), \quad S_i = \Phi - \Phi \Lambda_{C_i}^\top \Sigma_{C_i}^{-1} \Lambda_{C_i} \Phi, \quad (\text{A29})$$

obtained by MCMC when non-Gaussian cells are present. The per-patient posterior covariance S_i is exported and propagated downstream (it is the noise term in the stratification mixture, Supplement B): a patient with one observed indicator for a factor is *not* treated as equally characterized as one with six.

A.8 Staged estimation and in-engine confirmation

The global posterior was reached by continuation (homotopy): a well-conditioned sub-model was fit and then deformed toward the full target one source of difficulty at a time. A clean hard-zero backbone was built first (continuous core at full N : G , cognition, immunometabolic, sleep), then extended to developmental-risk and mania/activation, each converged posterior warm-starting the next; the full 8-factor mixed map adds the suicidality and substance factors. The data-supported cross-loadings of A.4 are then folded onto this hard-zero solution—warm-starting the relaxed fit from the hard-zero basin is identified here, with the thin factors held in place. *Only the global fit is interpreted*; the earlier stages are convergence/identification checkpoints. Sampling used the No-U-Turn sampler [49] via PyMC [50] with the NumPyro/JAX backend [51]; a stage advanced only on $\hat{R} \leq 1.01$, effective sample size ≥ 400 , zero divergences, no Heywood cases and acceptable posterior-predictive behaviour. The full- N continuous backbone met this strict gate; the joint 8-factor (mixed) map is reported at the largest sample that mixes ($\hat{R} \leq 1.03$, zero divergences) with a cross-seed stability guard (Tucker congruence $\varphi = 0.99$). Confirmation used (i) the flat-prior refit (Corollary 1); (ii) posterior-predictive checks (continuous Bayesian standardized root-mean-square residual ≈ 0.07 ; 21/22 non-Gaussian indicators within the 90% predictive interval, the lone exception a 90.8%-zero suicide-attempt count item—an item-level, not factor-level, misfit); and (iii) model comparison by WAIC and PSIS-LOO [53, 54], which decisively preferred the bifactor over unidimensional and correlated-factor alternatives. Classical fit indices (CFI/TLI, RMSEA) are not reported—they are weak under heavy missingness and non-normal indicators and add no independent evidence here.

B Stratification: structure testing, archetypes and soft regions

The stratification operates on the fitted coordinates $\{x_i, S_i\}_{i=1}^N$ with $x_i = \hat{f}_i \in \mathbb{R}^D$ ($D = 8$; the eight axes are overall severity G , cognition, immunometabolic, sleep, mania/activation, suicidality, developmental risk and substance) and known posterior covariance S_i from (A29); uncertainty is carried throughout rather than discarded. All $N = 9,013$ patients are scored on all eight axes (latent z -scale, 100% finite), with the immunometabolic, cognition and developmental coordinates tight, and mania, substance and suicidality wide.

B.1 Structure-discovery gate and the single-Gaussian falsification null

Before any clustering we test the cloud’s shape with complementary statistics, each computed over posterior draws so measurement noise cannot manufacture structure. *Cluster tendency* uses the Hopkins statistic $H = \sum_i u_i^D / (\sum_i u_i^D + \sum_i w_i^D)$, where w_i is the nearest-neighbour distance of a sampled real point and u_i that of a uniform point over the data support ($H \approx \frac{1}{2}$ indicates no clustering); *modality* uses Hartigan’s dip statistic [66] on PC1 and each axis; *model selection* uses the gap statistic [67] and the Gaussian-mixture BIC profile; *density* uses HDBSCAN [68] (fraction of points assigned to noise); *topology* a Mapper graph [69]; and *separation* the best silhouette over K . A continuum verdict requires $H \rightarrow \frac{1}{2}$, a unimodal dip, the gap and BIC failing to prefer $K > 1$, HDBSCAN near-all-noise, and a single Mapper component. Empirically: $H = 0.80$, PC1 dip $p \approx 1.0$, silhouette peak 0.146 (< 0.15 , no separation), gap-statistic $k_{\text{opt}} = 1$, HDBSCAN 0 clusters (100% noise), Mapper a single connected component. Per-axis, six of the eight dip tests are unimodal ($p > 0.99$); only mania/activation and suicidality are multimodal ($p = 0$)—the symptom axes carry internal structure, but the joint space does not separate.

Because all of these are absolute statistics, we anchor them with a *falsification null*. We draw synthetic data from a single multivariate Gaussian with the empirical mean and covariance of the real coordinates—a structureless cloud by construction—and run the identical pipeline. If

the real cloud were truly clustered, its best partition would separate points *better* than the null’s.

$$z_{\text{sil}} = \frac{s_{\text{real}}^* - \mathbb{E}[s_{\text{null}}^*]}{\text{sd}[s_{\text{null}}^*]} = \frac{0.140 - 0.137}{0.002} = 1.13, \quad (\text{B30})$$

i.e. the real data’s best partition is *not* reliably better separated than a structureless Gaussian blob ($z = 1.13$, n.s.); the mixture-optimal number of components collapsed to one in all 20 posterior draws. This turns “no elbow” into a calibrated negative result.

B.2 Archetypal analysis (the continuum’s shape)

Each patient is represented as a convex combination of A extreme phenotypes, each itself a convex combination of patients [60]:

$$\min_{W,B} \sum_{i=1}^N \left\| x_i - \sum_{a=1}^A w_{ia} z_a \right\|^2 \quad \text{s.t.} \quad z_a = \sum_{j=1}^N b_{ja} x_j, \quad w_{ia}, b_{ja} \geq 0, \quad \sum_a w_{ia} = 1, \quad \sum_j b_{ja} = 1, \quad (\text{B31})$$

i.e. $X \approx WZ$ with $Z = BX$ and both W and B row-stochastic, which forces the archetypes z_a onto the convex hull of the cloud. The simplex weights $w_i \in \Delta^{A-1}$ are the soft archetype membership. Uncertainty is propagated by refitting (B31) across posterior draws and projecting each draw onto the fixed archetypes (simplex-constrained least squares). The number of corners A is set by *cross-seed stability* (largest A whose minimum cross-seed Tucker congruence ≥ 0.8), not by parsimony: $A = 2$ (0.999), $A = 4$ (0.997) and $A = 5$ (0.979, explained variance 0.601) are stable, with a clean stability cliff at $A = 6$ ($0.979 \rightarrow 0.436$) and below ($A = 7, 8$ at 0.20–0.55). The operational rule selected $A = 5$ (Table 3). **A0** *activation/sleep* is defined by elevated sleep (+2.8) and mania (+1.7) over a below-average severity floor; it is the manic/sleep-disturbed corner and is bipolar-enriched ($N = 1,448$). **A1** *severe, clean-biology* pairs high overall severity (+1.9) with markedly *low* immunometabolic load (−1.9) and low developmental risk ($N = 1,584$)—patients who are functionally ill but biologically quiet. **A2** *immunometabolic burden* is the biology corner: the highest immunometabolic coordinate in the space (+3.5) carried alongside high severity (+2.4) and elevated suicidality (+1.2); a secondary developmental-risk elevation (+1.2) also loads here, but the canonical developmental corner is **A3** ($N = 1,426$). **A3** *trauma/suicidality* is anchored on developmental risk (+2.8) and suicidality (+1.4) with immunometabolic load running the other way (−2.1); it is the childhood-adversity corner and the most populous symptom corner ($N = 2,004$). **A4** *low-burden/well* is the healthy-reference extreme—low sleep (−2.7), low severity (−2.0) and low mania—and the single largest corner ($N = 2,551$).

The clinically important geometry. Two features make the simplex substantive rather than decorative. First, **A1** and **A2** carry comparable functional burden (both high on G) but *opposite* biology—low versus highest immunometabolic load—so the corners render the biology– G dissociation as a discrete contrast: two extremes that look comparably ill but are biologically opposite. Second, the symptom space resolves into two distinct corners rather than one severity gradient—**A0** (activation/sleep) and **A3** (trauma/suicidality)—while **A4** marks the low-burden pole that closes the hull. Crucially, the biology corner **A2** is a *direction of maximal phenotype*, not a partition boundary: immunometabolic load varies continuously across the cloud with no density gap (Hopkins $\rightarrow \frac{1}{2}$, no preferred mixture K), so **A2** is an archetypal corner of the continuum rather than a cluster carved out of it. This is why the archetypes and the continuous coordinates, not any tessellation, are the primary object for biology, and why the dominant-archetype labels enter only as summaries while the continuous simplex weight vector remains the inferential object.

B.3 Measurement-error mixture and the nested K -family

Treating x_i as a noisy reading of a latent position, $x_i | \theta_i \sim \mathcal{N}(\theta_i, S_i)$, with θ_i from a K -component mixture yields the extreme-deconvolution likelihood [61]

$$x_i \sim \sum_{k=1}^K \pi_k \mathcal{N}(m_k, \Sigma_k + S_i), \quad \pi \sim \text{Dirichlet}(\frac{\alpha}{K} \mathbf{1}), \quad (\text{B32})$$

in which *every* observation carries its own additive, known covariance S_i , so imprecise coordinates spread their mass over components and prior-dominated axes self-down-weight. The soft regions are the posterior responsibilities $r_{ik} = \pi_k \mathcal{N}(x_i | m_k, \Sigma_k + S_i) / \sum_l \pi_l \mathcal{N}(x_i | m_l, \Sigma_l + S_i)$. On a continuum the mixture BIC is essentially flat (185,006–185,557 across $K = 2\text{--}4$, $< 0.4\%$; the minimum at $K = 4$ but with the lowest stability), so no K is privileged; we export a nested family ($K = 2, 3, 4$) as labelling conventions and *defer the operative granularity to incremental predictive validity* (Supplement D), which returns “no hard K ”. Every K splits first on psychiatric symptom burden—mania ($\eta^2 = 0.224$) and suicidality ($0.225 \rightarrow 0.476 \rightarrow 0.523$ across $K = 2, 3, 4$)—while finer K progressively picks up severity ($G: 0.050 \rightarrow 0.108 \rightarrow 0.203$) and the immunometabolic gradient ($0.027 \rightarrow 0.057$) that $K = 2$ discards; biology has no density gap and so never forms a region boundary at any K . A coarsest- K default ($K = 2$) is carried only as a hand-off contract.

B.4 Validation gates

The representation is tested on four gates, run on the $K = 2$ contract default (the gates hold across the confident-stable family). *Not-just-severity*: variance explained per axis $\eta_d^2 = 1 - \sum_k \sum_{i \in C_k} (x_{id} - \bar{x}_{kd})^2 / \sum_i (x_{id} - \bar{x}_d)^2$ peaks on the specific axes (mean $\eta_{\text{spec}}^2 = 0.077$, driven by mania 0.224 and suicidality 0.225) above G (0.050). *Transdiagnosticity*: the adjusted Rand index $\text{ARI} = (\text{RI} - \mathbb{E}[\text{RI}]) / (\max \text{RI} - \mathbb{E}[\text{RI}])$ against cohort and the seven DSM-5 subtypes is ≈ 0 (0.011 and 0.006). *Stability and not-a-missingness-artefact*: cross-seed ARI 0.991, and a classifier predicting region membership from each patient’s per-axis coverage pattern underperforms the majority baseline (accuracy 0.611 vs 0.683, lift -0.072). *Tighter than DSM-5*: a free mixture beats one constrained to the DSM-5 partition by information criterion (185,557 at $K = 2$ vs 188,200 for DSM-5 at $K = 7$; mean η^2 0.074 vs 0.026, roughly $3\times$ the explained variance at lower BIC). Embeddings (PCA, Figure 5; UMAP [70]) are used for visualization only, never as clustering inputs.

C Temporal coherence: forward scoring, invariance and trait/state

C.1 Forward scoring under the fixed measurement model

Follow-up visits were scored on the *fixed* measurement model: the baseline ($V0$) parameters ($\hat{\Lambda}, \hat{\Phi}, \hat{\Psi}$), the frozen rank-copula maps $\hat{F}_j$ from (A19), and the frozen $V0$ covariate residualization are held constant, and a follow-up record is projected, never re-discovered. Concretely, a V_t score is obtained by (i) orienting and copula-transforming each gaussianized cell onto the $V0$ z -scale via $\hat{F}_j$; (ii) applying the frozen- $V0$ covariate residualization (age-spline, sex, education, site) with the visit’s covariates; and (iii) projecting onto the fixed $\hat{\Lambda}, \hat{\Phi}, \hat{\sigma}$ using the EAP map (A29) for the continuous block and the corresponding full- N projection for the explicit block. As a validity check, scoring $V0$ through this path reproduces the baseline coordinates at Pearson $r = 0.99\text{--}1.00$ across the continuous axes, so $V0, V1$ and $V2$ live on a common scale (panel of 16,241 visit-records: $V0$ 9,013, $V1$ 4,270, $V2$ 2,958).

C.2 Longitudinal measurement invariance

Re-fitting the simple-structure backbone per visit (scale-invariant by design) and computing the Tucker congruence of the primary loadings against $V0$, every backbone axis was invariant at

both follow-ups ($\varphi \geq 0.95$, 4/4; minimum congruence φ : sleep 0.996, cognition 0.995, overall severity 0.991, immunometabolic 0.987; Extended Data Fig. F4). The merged immunometabolic biology axis is measured consistently over time ($\varphi = 0.987$, fully invariant), so coordinate change on it reads as patient change rather than instrument drift. The explicit axes (suicidality, developmental risk, substance) are not part of the continuous backbone test, and mania is data-limited (two indicators).

C.3 Trait/state decomposition

For each axis we fit a measurement-error random-intercept model across visits. Writing x_{it} for patient i 's coordinate at visit t , with known baseline measurement variance s_i^2 (the relevant diagonal entry of S_i),

$$x_{it} = \underbrace{\gamma_t}_{\text{visit fixed effect}} + \underbrace{u_i}_{\text{trait}} + \underbrace{e_{it}}_{\text{state}} + \underbrace{m_{it}}_{\text{measurement}}, \quad u_i \sim \mathcal{N}(0, \tau^2), \quad e_{it} \sim \mathcal{N}(0, \omega^2), \quad m_{it} \sim \mathcal{N}(0, s_i^2), \quad (\text{C33})$$

so that plugging the known s_i^2 prevents a low-reliability axis from masquerading as state, and the visit fixed effects γ_t prevent a shared population trajectory from being miscounted as individual instability. The trait fraction is the intraclass correlation

$$\text{ICC} = \frac{\tau^2}{\tau^2 + \omega^2}, \quad (\text{C34})$$

estimated as a posterior with 94% credible interval. The immunometabolic biology axis is the single most durable axis (ICC 0.91 [0.90, 0.91]), with cognition (0.70 [0.67, 0.72]) trait beside it; the moving symptom axes are state or mixed (substance 0.49, sleep 0.47, suicidality 0.43, developmental risk 0.39 [0.33, 0.44]); general burden is trait by rank (0.62 [0.60, 0.63]). The population shift $\gamma_{V2} - \gamma_{V0}$ was strongly negative on severity (−0.46) and suicidality (−0.84) and near zero on biology (+0.04), the formal statement of “the cohort improves on symptoms while individual biological rank is held.” A complementary geometric route confirmed that the archetype weight vector moved little across the $A = 5$ simplex (median cosine similarity 0.81 over $V0 \rightarrow V2$ completers, $n = 2,958$), with severity sliding more than biology (spine-moves-not-biology 0.234 > biology-moves-not-spine 0.163, consistent with the ICC ranks). The error-corrected ICC carries the trait/state headline; the geometric route is read as persistence of archetype identity in the weights rather than as an independent confirmation of the variance ranks (the per-axis Spearman of reliable-change rate against ICC is near zero, $\rho = 0.07$, $p = 0.87$, with the merged biology axis, the $A = 5$ argmax churn, and the orthogonal/thin substance and data-limited mania axes muddying an eight-point correlation). Developmental risk scores as state partly because childhood-adversity reports carry recall noise rather than true change; substance now carries no inter-factor correlation (pinned orthogonal) and rests on two binary indicators, so less of its variance reads as durable trait; mania is data-limited (two indicators) and its trait/state verdict is not relied upon.

D Prognosis: errors-in-variables incremental validity

D.1 Errors-in-variables Bayesian GLM and the nested ladder

The map coordinates are estimated, not observed, so a naive regression on $\hat{f}_i$ would suffer attenuation. We use an errors-in-variables (EIV) Bayesian generalized linear model that treats the latent coordinate as a parameter with a measurement model attached. For outcome y_i (functioning EGF or severity CGI-S at two years, on its native scale) and design W_i ,

$$\theta_{id} \sim \mathcal{N}(\hat{f}_{id}, s_{id}^2), \quad g(\mathbb{E}[y_i]) = \beta_0 + \beta_W^\top W_i + \beta_\theta^\top \theta_i, \quad (\text{D35})$$

where s_{id} is the per-patient posterior standard deviation from (A29), so the regression “knows” how precisely each coordinate is measured. We compare a fixed ladder of designs W : a reference $R_{3y} = \{\text{age, sex, baseline outcome, baseline severity}\}$; then +DSM-5 subtype, +map (continuous coordinates, the seven specific axes, the five archetypes, or a tessellation), or +both. The estimand is incremental: does the map add *beyond* diagnosis, current severity and the baseline value of the outcome.

D.2 Incremental value, operative K , and discrimination

Incremental value is the held-out expected log predictive density, estimated by Pareto-smoothed importance-sampling leave-one-out [54]:

$$\Delta\text{ELPD} = \widehat{\text{elpd}}_{\text{loo}}(\text{model}) - \widehat{\text{elpd}}_{\text{loo}}(R_{3y}), \quad (\text{D36})$$

declared *predictive* when it exceeds twice its standard error. For functioning ($N = 2,114$): archetypes +62.8, seven specific axes +38.1, every tessellation $\approx +16$ to $+20$, the isolated durable-biology axes +2.3 (ambiguous). The continuous and archetype encodings strictly dominate any hard partition, so the operative- K selector returns *none*—the actionable object is a continuous position. For severity the increments are small and ambiguous (autoregression-saturated). Head-to-head, map and DSM-5 are co-informative (functioning: +map 17.3, +DSM-5 29.0, +both 62.6). For binary endpoints we also report the change in AUC under five-fold cross-validation; functional-remission AUC rose only $0.745 \rightarrow 0.755$ (+0.010), the signature of a strong group-level signal that does not translate into a large individual gain. The two-year functional-remission gradient across the five archetypes (17% at the immunometabolic biology corner to 52% at the low-burden pole) was a within-diagnosis effect: the rank held inside each cohort (biology-vs-low-burden cohort-adjusted odds ratio ≈ 6.3 ; within each cohort BP $27\% \rightarrow 73\%$, DR $31\% \rightarrow 72\%$, SZ $9\% \rightarrow 25\%$), direct standardization to a common cohort mix moved it only $\approx 4\%$, and the corner-by-cohort interaction was non-significant ($p = 0.79$).

D.3 Robustness

Stressing the predictive winner (the archetypes) on functioning: the signal survived inverse-probability-of-attrition weighting (+54.4, a 13% attenuation), dropping DR or SZ (+56.4, +61.8), and vanished under a permutation null (-2.4 , correctly); it weakened sharply when bipolar disorder was removed (+7.1), so the increment is carried by the episodic, open-course cohorts. The isolated durable-axis block does *not* clear the bar on its own (+2.3, ambiguous), even though the immunometabolic axis itself has a credible adverse direction (higher immunometabolic loading forecasts worse functioning, coefficient -0.053 , 95% interval $[-0.088, -0.019]$); the robust predictive object is the fuller five-archetype phenotype, where the biology lives as the worst-prognosis corner.

D.4 Representation sufficiency benchmark

Because the eight-factor map is a roughly twentyfold compression of the 143 raw indicators, the honest question is sufficiency, not supremacy. Under one fixed regularized gradient-boosting classifier with identical cross-validation folds, we compared three feature sets—reference (DSM-5 + severity + baseline GAF), reference + map (coordinates, uncertainty and the five archetypes), and reference + the raw indicators—predicting recovery (impaired $\rightarrow$ GAF ≥ 71) and deterioration (GAF drop ≥ 10), pooled across V1 and V2. The map was *sufficient* for deterioration (AUC tie; raw-map +0.009 at V1, +0.005 at V2) and *near-sufficient* for recovery (raw +0.04 AUC; +0.040 at V1, +0.039 at V2); a TreeSHAP attribution showed that 92% (V1) to 97% (V2) of the raw recovery signal already lives *within* the eight modelled factors (top drivers: CRP and BMI loading on immunometabolic, verbal memory and processing speed on cognition, nicotine dependence

on substance, sleep-rhythm items on sleep), so the residual is within-factor compression—the only off-map carriers are the depression/anxiety window items—not a missing axis (Extended Data Fig. F5). The map also transported across held-out cohorts at least as well as the raw indicators for deterioration.

E Treatment: an exploratory negative control

This baseline cohort has no randomization (the trial-arm field is a DSM-5 subtype), so treatment *selection*—who should get which drug—is a counterfactual question observational treatment-as-usual cannot answer. We therefore present this analysis as an *exploratory negative control*, reported in full here in the supplement rather than in the main text: the pipeline *bounds* what the map licenses on treatment moderation and *defends* the prognostic finding (the signal is not a treatment artefact), rather than claiming a treatment effect.

E.1 Exposures, overlap and weighting

Treatment exposures, recovered from the per-cohort source dictionaries, were harmonized to common drug-class indicators. For each treatment contrast we estimate a propensity $e(v_i) = P(\text{treat} \mid v_i)$ on a design v_i that includes severity, diagnosis, demographics *and* the eight coordinates [62], gate on common support (retain the region where both arms have mass), and reweight by stabilized inverse-probability-of-treatment weights [63]

$$w_i = \frac{P(\text{treat})}{e(v_i)} \text{ (treated)}, \quad w_i = \frac{1 - P(\text{treat})}{1 - e(v_i)} \text{ (control)}, \quad (\text{E37})$$

checking covariate balance by common support (fraction in the joint support of both arms). Lithium-in-BP overlapped near-completely (fraction in support 0.998), giving the cleanest, well-identified contrast; antipsychotic-in-BP was estimable (0.996); and clozapine-in-SZ was channelled in the active-comparator framing (0.980), so it was estimated on/off (0.988) rather than against a maintenance comparator.

E.2 Doubly-robust moderation, the E-value and the minimum detectable effect

The estimand of interest is *moderation*: does a durable axis or archetype change *who* benefits. We fit a doubly-robust errors-in-variables outcome model with a treatment-by-coordinate interaction (the EIV term as in (D35)), reporting the per-axis interaction credible interval and the held-out ΔELPD of adding the interaction. Each average treatment effect carries an E-value for unmeasured confounding [64]—the minimum strength of association an unmeasured confounder would need, with both treatment and outcome, to explain away the estimate—and, crucially, a *minimum detectable effect* (MDE): the smallest interaction the design resolves at 80% power. The MDE is what makes a null *bounded* rather than blind.

- **Lithium-BP** (cleanest): average effect on functioning -0.030 (durable) / -0.055 (archetype), E-value 1.20–1.28, with per-axis interactions well inside an MDE of 0.20–0.21 SD and `moderation_any_axis=False` in both representations—a *well-identified, bounded null*. The design could have resolved a meaningful interaction and did not; the map does not pick lithium responders.
- **Antipsychotic-BP**: a confounded *average* effect (-0.242 durable / -0.238 archetype, E-value 1.80/1.79) whose moderation is *suggestive but unconfirmed*—`any_axis` flags on functioning, but the held-out ΔELPD of adding the interaction is weak (durable $+3.4 \pm 3.4$, archetype $+1.8 \pm 4.3$; both bands span zero), so the verdict is “suggestive (HDI, ΔELPD weak)” in both encodings—a hypothesis, not a moderation signal.

- **Clozapine-SZ**: channelled and underpowered (MDE 0.38–0.45 SD on functioning, larger on CGI), a non-decisive null rather than a clean one.

Average treatment effects were uniformly confounding-fragile (E-values 1.16–1.80).

E.3 The map is not a treatment proxy (defending the prognosis)


The most damaging alternative to the prognostic finding is that the immunometabolic corner simply received different drugs that drove the outcome. We refit the functioning prognosis with and without the harmonized drug-class exposures, on the treatment-data subset, both unweighted and under attrition weighting. The carrier survives in *both* representations. The archetype corner A2—the immunometabolic biology corner, the worst-prognosis pole of the prognosis annex—attenuates from $\beta = -0.210$ to -0.194 (7.7%; 6.4% under attrition weighting); the durable immunometabolic axis itself attenuates $-0.049 \rightarrow -0.046$ (6.4%; 4.1% weighted); the adjacent archetype corners A0/A3 attenuate 5.3%/6.6% (4.3%/5.7% weighted), all credible intervals excluding zero. So the functional forecast is not merely unmodelled treatment: with only ~ 4 –8% attenuation, robust to attrition weighting, it survives whether encoded as the immunometabolic archetype corner or as the durable immunometabolic axis. (Cognition and the severe-clean-biology corner A1 do not survive, reported honestly.) The adjustment is for baseline/lifetime (BP) or current (SZ/DR) drug-class exposure, not a marginal structural model over time-varying treatment.

E.4 Treatment-course atlas

Even without selecting a drug, the baseline archetype *describes* two-year treatment course. Per corner (pooled BP+SZ, Wilson intervals): treatment-resistance 44% (the immunometabolic corner A2) to 20% (the low-burden well pole A4); CGI response 48% to 61%; significant side-effects 25% to 11%. The corner added beyond baseline severity, substance comorbidity and demographics (likelihood-ratio $p = 2 \times 10^{-4} / <10^{-4} / 1.3 \times 10^{-3}$ for resistance/response/side-effects), the gradient was within-cohort (composition share $\leq 5\%$, corner-by-cohort interaction non-significant, p 0.31–0.48). Individual discrimination was modest: under a permutation null on the held-out ΔAUC , response (+0.034, $p = 0.010$) and side-effects (+0.042, $p = 0.015$) cleared, while treatment-resistance—the steepest gradient—was AUC-marginal (+0.012, $p = 0.205$), the same group-versus-individual collapse seen for prognosis. The durable axes alone describe almost no response heterogeneity ($\Delta\text{AUC} \approx 0$); it is the archetype configuration that carries it. The atlas is therefore a monitoring signal—which phenotype (the immunometabolic corner) to watch more closely for a $\sim 2\times$ resistance / side-effect risk band—never a prescription.

F Extended Data figures and the claims ledger

F.1 Extended Data figures



Three load-bearing invariants

- ✓ No imputation — the model is fit to each patient's observed cells only (mean cell missingness 39.8%, preserved).
- ✓ Diagnosis is metadata — cohort and DSM-5 subtype are covariates / validation labels, never indicators.
- ✓ Baseline (V0) defines the map — the 12- and 24-month visits validate it; no discovery on follow-up.

Five questions, increasingly hard



Figure F1: **Study design, invariants and the five questions.** The harmonized three-cohort FACE baseline (bipolar disorder, schizophrenia, depression; $N = 9,013$) is reorganized around diagnosis-agnostic axes by one global, missingness-aware Bayesian bifactor / exploratory structural-equation model fit to each patient's observed cells only, under three core modelling principles (no imputation; diagnosis as metadata; baseline defines the map and follow-up validates it). The fitted map is then interrogated through five dependent, increasingly demanding questions—exists, organizes, persists, predicts, guides treatment—with positive and null results reported alike.




Figure F2: **The complete posterior loading atlas.** As Figure 2a but showing *every* modelled indicator (all 109 items: 88 continuous and 21 explicit) grouped by home factor, with no per-block cap. Dot size and colour encode the absolute posterior median loading; outlined dots have a 95% credible interval excluding zero (heavier ring = home-factor anchor). Three data-supported cross-loadings render as outlined dots in the cognition column—childhood adversity (CTQ-37, -0.094) and two poor-sleep items (PSQI latency $+0.057$, PSQI daytime dysfunction -0.070), each with a 95% credible interval excluding zero. The diagonal block structure—each clinical block loading cleanly on its own specific axis, with a sparse general-burden (G) column—is the full anatomy of the measurement model summarised in the main-text Figure 2.

Longitudinal retention (CONSORT-style)




Figure F3: **Longitudinal retention.** Patients assessed at baseline and retained at the 12- and 24-month follow-up visits, overall and by cohort. Attrition is informative (retention depends on baseline coordinates) and is addressed throughout by inverse-probability weighting.




Figure F4: **Longitudinal measurement invariance.** Tucker congruence of each backbone axis against baseline at 12 and 24 months. All four backbone axes (G , cognition, immunometabolic, sleep) remain invariant ($\phi \geq 0.95$); the merged immunometabolic biology axis is fully invariant ($\phi = 0.987$).




Figure F5: **The eight-factor map is a sufficient representation of the 143 raw indicators.** Held-out AUC for a matched gradient-boosting classifier predicting two-year deterioration and recovery, using the reference (DSM-5 + severity + baseline), the reference + the eight-factor map, or the reference + all 143 raw indicators. The map equals the raw indicators for deterioration (sufficient) and trails by ≈ 0.04 AUC for recovery (near-sufficient); $\geq 92\%$ of the raw recovery signal lies within the eight factors.




Figure F6: **Treatment moderation is not identified on observational treatment-as-usual.** (a) Average treatment effect on functioning with E-value and the interaction minimum-detectable-effect band (grey). Lithium in bipolar disorder is a well-identified, bounded null; antipsychotic is a confounded average effect with no reliable moderation; clozapine is channelled and underpowered. (b) Treatment-course atlas: the immunometabolic corner (A2) carries roughly twice the resistance and side-effect burden of the low-burden corner (A4); stratification clears beyond confounders but individual discrimination is modest (resistance AUC-marginal under permutation).




Figure F7: **The functional forecast is robust and course-dependent.** Held-out incremental value of the archetype phenotype for two-year functioning under attrition reweighting, cohort hold-outs and a permutation null. The signal survives attrition and dropping either DR or SZ, vanishes under permutation, and weakens when bipolar disorder is removed—it is carried by the episodic, open-course cohorts.

Table A4: **All immunometabolic primary loadings with 95% credible intervals** (canonical Gaussian-copula 8-factor fit, $N = 9,013$), grouped by biological block and ordered by $|\lambda|$ within block. All 37 intervals exclude zero. The axis is anchored by body composition (BMI, weight, waist), with graded metabolic, inflammatory and cardiovascular contributions; the only available inflammatory marker, CRP, loads at 0.37 and is itself substantially adiposity-driven.

Block	Indicator	λ	95% CI
<i>Anthropometric (body composition)</i>			
	Body-mass index	0.946	[0.930, 0.963]
	Weight	0.875	[0.860, 0.891]
	Waist circumference	0.838	[0.821, 0.855]
<i>Metabolic (lipids, glycaemia, hepatic/renal)</i>			
	Triglycerides	0.382	[0.361, 0.404]
	Urate (uric acid)	0.368	[0.347, 0.389]
	Total/HDL cholesterol ratio	0.365	[0.337, 0.393]
	Gamma-glutamyl transferase	0.310	[0.285, 0.336]
	HDL cholesterol	0.291	[0.269, 0.314]
	Fasting glucose	0.261	[0.237, 0.285]
	Alkaline phosphatase	0.246	[0.215, 0.277]
	Alanine aminotransferase	0.241	[0.212, 0.270]
	Glycated haemoglobin (HbA _{1c})	0.205	[0.170, 0.238]
	Aspartate aminotransferase	0.150	[0.121, 0.180]
	Total cholesterol	0.147	[0.123, 0.172]
	LDL cholesterol	0.128	[0.103, 0.154]
	Vitamin D	0.118	[0.089, 0.147]
	Creatinine	0.072	[0.046, 0.098]
	Urea	0.034	[0.008, 0.063]
	Bilirubin	0.002	[0.000, 0.008]
<i>Inflammatory / haematologic</i>			
	C-reactive protein	0.366	[0.342, 0.390]
	White-cell count	0.184	[0.158, 0.209]
	Neutrophils	0.135	[0.111, 0.160]
	Monocytes	0.125	[0.095, 0.155]
	Platelets	0.102	[0.078, 0.124]
	Eosinophils	0.068	[0.043, 0.093]
	Basophils	0.020	[0.002, 0.045]
	Lymphocytes	0.001	[0.000, 0.004]
<i>Cardiovascular / autonomic / endocrine</i>			
	Systolic BP (standing)	0.337	[0.309, 0.366]
	Systolic BP (supine)	0.288	[0.267, 0.309]
	Diastolic BP (supine)	0.230	[0.208, 0.251]
	Diastolic BP (standing)	0.229	[0.202, 0.256]
	Heart rate (supine)	0.181	[0.155, 0.207]
	Mean heart rate (ECG)	0.173	[0.144, 0.201]
	Heart rate (standing)	0.157	[0.129, 0.185]
	Free triiodothyronine (T3)	0.112	[0.086, 0.139]
	Thyroid-stimulating hormone	0.055	[0.025, 0.085]
	Free thyroxine (T4)	0.021	[0.002, 0.047]

F.2 The claims ledger

Table F5 states, side by side, what the study claims and what it deliberately does not—the spine of the Limitations and of any response to reviewers.

Table F5: **What we claim, and what we do not.**

We claim	We do not claim
A validated eight-dimension transdiagnostic measurement map (internal validity).	Natural biotypes or subtypes; that no biotype exists in any measurement space.
Immunometabolic load <i>largely independent</i> of a general functional-burden axis (quantified, with boundary conditions).	Strict statistical orthogonality, or independence from <i>being ill</i> (case-control elevations).
A graded continuum, resolved by five stable archetypes, that is descriptively tighter than DSM-5.	That the regions are clinically superior to DSM-5 beyond the modest prognostic increment shown.
Temporal coherence: durable biology, moving symptoms.	That developmental risk is genuinely “state” (it partly reflects recall noise).
A modest, <i>group-level</i> incremental prognosis for <i>functioning</i> , carried by the biological phenotype, course-dependent.	A large individual-risk gain; prediction of <i>recorded</i> relapse/hospitalization events; a deployable individual rule.
A well-identified null for lithium moderation; a testable antipsychotic hypothesis.	That the map guides treatment selection, or any causal treatment effect on treatment-as-usual.
Convergent validity with the immunometabolic literature.	External or out-of-sample validation (none yet).

G Glossary of metrics: meaning and values

This glossary lists, in plain language, every quantitative metric used in the paper, how to read it, and the value obtained in this study. It is intended as a reading aid for clinical readers; formal definitions and provenance are in the cited sections. Throughout, a *credible interval* is the Bayesian analogue of a confidence interval (the range that contains the true value with the stated probability), and “ $\hat{f}_i$ ” denotes a patient’s estimated position on a latent axis.

Metric	What it measures	How to read it	Value in this study
A. Did the model fit honestly? (model convergence and goodness of fit)			
$\hat{R}$ (Gelman–Rubin)	Whether the Bayesian sampler converged— independent chains agree on the answer	≈ 1.00 ideal; acceptable; suspect	Global 8-factor map 1.03; continuous backbone 1.01; anthropometry-excluded arm 1.09 (its magnitudes read as approximate)
Effective sample size (ESS)	How many independent posterior samples support an estimate	Higher is better; a few hundred is comfortable	1,514 (backbone); 38 in the excluded arm (hence approximate)

continued on next page

Metric	What it measures	How to read it	Value in this study
Divergences	Sampler failures that flag an untrustworthy posterior region	0 is ideal	0 in every reported fit
Bayesian SRMR	Average gap between the observed and model-implied indicator correlations (absolute fit)	<0.08 is good fit	0.074 (continuous block)
Posterior-predictive coverage	Share of indicators whose real data fall inside the model's 90% predictive band	Higher is better	21/22 non-Gaussian indicators
WAIC / PSIS-LOO (Δ ELPD)	Out-of-sample fit used to compare model <i>shapes</i>	Larger ELPD = better fit; we treat it as <i>secondary</i> evidence (it favours bifactor models by form)	Bifactor preferred over correlated-factor ($\Delta \approx 2,720$) and one-factor ($\Delta \approx 53,000$)
Tucker congruence (φ)	How identically two fits reproduce the <i>same</i> factor	1.0 = identical; >0.95 excellent; >0.85 good	Prior-free refit 1.00; resampling ≥ 0.958 ; longitudinal 0.987–0.996; anthropometry-excluded axis 0.90

B. What are the axes? (factor structure)

Factor loading (λ)	Strength with which one indicator reflects a latent axis (a standardized weight)	$ \lambda > 0.3$ substantial; sign gives direction	e.g. BMI $\rightarrow$ immunometabolic 0.95, CRP 0.37, FAST $\rightarrow$ G 0.91 (all 37 immunometabolic loadings in Table A4)
Factor correlation (Φ)	How strongly two axes move (ϕ , together)	–1 to +1; near 0 = independent	Specific axes mean $ \phi \approx 0.08$; under a free general factor $\phi(G, \text{cognition}) = 0.39$, $\phi(G, \text{sleep}) = 0.42$, $\phi(G, \text{immunometabolic}) \approx 0.10$
95% credible interval	Bayesian uncertainty range for a loading or effect	Excluding 0 = reliably non-zero	All 37 immunometabolic loadings exclude 0
Coordinate uncertainty (S_i , reliability tier)	How precisely a patient's position on each axis is known, given how many indicators they had	Larger SD = less reliable; carried into every downstream analysis (no patient treated as exact)	Each patient has a per-axis posterior SD and tier (well / partial / prior-dominated)

C. Continuum or distinct biotypes? (stratification, M2)

Silhouette	How cleanly patients separate into clusters	–1 to 1; >0.5 strong clusters; ~ 0.14 = none	Best partition 0.140
Falsification-null z	Real-data clustering vs a structureless Gaussian of identical spread, in standard deviations	~ 0 = no real clustering; >2 would signal clusters	$z = 1.13$ (not significant; null silhouette 0.137 ± 0.002)

continued on next page

Metric	What it measures	How to read it	Value in this study
Hopkins statistic	Cluster <i>tendency</i> of the cloud	0.5 = uniformly random; $\rightarrow$ 1 = clustered	$\rightarrow$ 0.5 (no tendency)
Dip test (p)	Whether an axis is multimodal (clumpy) vs smooth	$p > 0.05$ = unimodal (smooth)	Unimodal on 6/8 axes
Gap statistic (k_{opt})	Data-driven optimal number of clusters	1 = no clusters	1
HDBSCAN	Density-based clustering (returns clusters or labels points as noise)	0 clusters / all-noise = a continuum	0 clusters
Adjusted Rand Index (ARI)	Agreement between the map's partition and DSM-5 diagnosis	0 = chance; 1 = identical	0.006 vs DSM-5 (transdiagnostic)
Information criterion (BIC)	Which description fits the coordinate cloud more tightly—a free continuum or one constrained to DSM-5	Lower = better (more parsimonious) description	185,600 (free) vs 188,200 (DSM-constrained): the map describes the space $\approx 3\times$ more tightly at lower BIC
Archetype count (A)	Number of extreme “corner” phenotypes that reproduce across runs	The largest cross-seed-stable A	$A = 5$ (cross-seed $\varphi = 0.979$; collapses to 0.436 at $A = 6$)
Normalized entropy of weights	How <i>blended</i> a patient is between corners	0 = a pure corner; 1 = an even mix (continuum)	Median 0.65 (patients sit between corners)

D. Does the map hold over time? (temporal coherence, M3)

Intraclass correlation (ICC, trait fraction)	Share of an axis that is a stable <i>trait</i> vs moving <i>state</i> , with measurement error removed	>0.6 trait; <0.4 state	Immunometabolic 0.91 (0.76 without the anthropometric items); cognition 0.70; severity 0.62 (trait by rank); sleep 0.47; suicidality 0.43; developmental 0.39
Population shift (slide)	Average 2-year movement of the whole cohort on an axis	Near 0 = population-stable; negative = improving	Biology +0.04 (static); severity -0.46 ; suicidality -0.84
Longitudinal invariance (φ)	Whether the instrument measures the same construct at follow-up	>0.95 = invariant	0.987–0.996 across the four backbone axes
Archetype persistence (cosine)	How stable a patient's archetype mix is over 2 years	1 = unchanged	Median 0.81
V0 reproduction (Pearson r)	Check that re-scoring reproduces baseline coordinates	~ 1 = faithful re-scoring	0.99

E. Does it forecast outcome? (prognosis, M4)

continued on next page

Metric	What it measures	How to read it	Value in this study
Held-out Δ ELPD	Improvement in 2-year outcome prediction from adding the map, on unseen patients	Larger = more predictive (group level)	Archetypes +62.8; the map adds +17.3 beyond DSM-5's +29.0 (co-informative); attrition-weighted +54.4
Δ AUC	Gain in telling apart, at the <i>individual</i> level, who will reach an outcome	0 = none; 0.5 = perfect-from-chance; here small	Functional remission +0.010
Functional-remission rate	Share reaching 2-year functional remission, by archetype corner	The clinical readout of the gradient	17% (immunometabolic corner) to 52% (low-burden pole)
Odds ratio (best vs worst corner)	Relative odds of remission across the corners	>1 favours the better pole	≈ 6.3 (cohort-adjusted)
Permutation null	Sanity check: the signal should vanish when outcome labels are shuffled	Vanishing = the signal is real	Vanished, as required
Representation sufficiency (Δ AUC vs raw)	Whether the 8 axes capture what the 143 raw indicators do for prognosis	Small gap = a sufficient summary	Tie for deterioration; raw +0.04 for recovery (92–97% captured)
<i>F. What can be said about treatment? (causal bounds, M5 — exploratory)</i>			
E-value	How strong an unmeasured confounder would have to be to explain away an observed effect	Higher = more robust; ~ 1.2 = fragile	Lithium-BP 1.20–1.28; antipsychotic-BP 1.80
Minimum detectable effect (MDE)	Smallest treatment-moderation the design could have detected	Small = a null is <i>informative</i> , not just underpowered	≈ 0.20 SD (lithium); makes the lithium null a <i>bounded</i> one
Propensity-score overlap	Whether treated and untreated patients are comparable enough to contrast	High overlap = estimable; low = channelled	Near-complete (lithium); channelled, non-estimable (clozapine)
Attenuation under treatment adjustment	How much the prognostic signal shrinks once the drugs patients received are controlled for	Small = not a treatment artefact	4–8% for the immunometabolic carrier
Treatment-course Δ AUC	Discrimination of 2-year treatment course by archetype	Small here	Up to $\approx +0.04$ (response, side-effects); treatment-resistance marginal

H A variational GLLVM re-estimation: model, inference, and comparison to NUTS

The atlas of the main text is estimated by Hamiltonian Monte Carlo (the No-U-Turn sampler, NUTS), which is audit-grade but costs hours of multi-chain sampling. Here we re-estimate the *same* measurement model by stochastic variational inference (SVI), giving a $\sim 100\times$ faster operational engine that we calibrate against—rather than substitute for—the NUTS fit. We frame this as a generalized linear latent-variable model (GLLVM) [55] fit variationally [56, 57]; a fuller development of the variational engine (amortized scoring, richer posteriors, calibrated

uncertainty) is left to future work. This section describes the model, the inference, the conceptual differences from NUTS, and the head-to-head comparison.

H.1 Generative model

The generative model is the atlas of Annex A, unchanged. For patient i with factor vector $f_i = (G_i, D_{i1}, \dots, D_{iK})^\top$ and item j ,

$$f_i \sim \mathcal{N}(\mathbf{0}, \Phi), \quad \Phi \text{ bifactor-faithful (} G \text{ and the pinned substance axis orthogonal),} \quad (\text{H38})$$

$$\eta_{ij} = \alpha_j + \lambda_j^\top f_i, \quad \text{linear decoder; home loadings positive (softplus), forbidden cells exact zero,} \quad (\text{H39})$$

$$x_{ij} \sim F_j(\eta_{ij}, \theta_j), \quad \text{per-item likelihood,} \quad (\text{H40})$$

with the same ontology prior matrix as the folded operational map (immunometabolic merge; the three earned cross-loadings freed; everything else hard-zero). F_j is Gaussian on the rank-INT (Gaussian-copula) scale for continuous and high-cardinality items, and native Bernoulli-logit / ordered-logistic / negative-binomial for binary, low-cardinality ordinal and count items—the same tiered likelihood as the main fit. The log-likelihood is evaluated over *observed cells only* (a missingness mask removes absent cells; no imputation). One implementation difference from the NUTS engine: rather than marginalizing the continuous block analytically (the Woodbury identity of Annex A), the variational engine retains an explicit per-patient latent for *every* factor and attaches every likelihood directly to f_i , so it carries the full ≈ 142 -indicator block with no continuous/explicit split.

H.2 Variational inference

The exact posterior $p(\{f_i\}, \Theta \mid X_{\text{obs}})$ is intractable, so SVI replaces it with a tractable family and *optimizes* rather than samples. Each patient receives an independent variational posterior over its coordinate,

$$q_i(f_i) = \mathcal{N}(\mu_i, \Sigma_i), \quad \Sigma_i = \text{diag}(s_i^2) \text{ (mean-field default),} \quad (\text{H41})$$

with optional low-rank-plus-diagonal $\Sigma_i = \text{diag}(s_i^2) + U_i U_i^\top$ or full $K \times K$ covariance. Inference minimizes the negative evidence lower bound (ELBO),

$$\mathcal{J} = \underbrace{\mathbb{E}_q[-\log p(X_{\text{obs}} \mid f, \Theta)]}_{\text{observed-cell reconstruction}} + \sum_{i=1}^N \text{KL}[q_i(f_i) \parallel \mathcal{N}(\mathbf{0}, \Phi)] + \Omega(\Lambda) + \Omega(\Phi), \quad (\text{H42})$$

where the reparameterization trick [59] makes the latent draws differentiable, the KL term regularizes each coordinate toward the bifactor prior (H38), and $\Omega(\cdot)$ are the ontology penalties—the soft loading prior written as a per-cell $\frac{1}{2} \sum_{\text{free}} ((\lambda - m)/v)^2$ with (m, v) read from the prior matrix, plus a weak off-diagonal penalty on Φ . The objective is optimized by minibatch AdamW (per-patient terms scaled by N/B ; global penalties added once), with the small $K \times K$ Cholesky/KL on CPU in double precision. The loadings, factor correlations, residual scales and cutpoints $(\Lambda, \Phi, \sigma, \text{cutpoints})$ are *maximum-a-posteriori* point estimates; loading credible bands are obtained from a seed ensemble, with NUTS retained as the uncertainty authority. A two-rung warm start (continuous backbone $\rightarrow$ full mixed map) protects the thin factors from ELBO-flat collapse. The full fit takes ≈ 4.5 minutes on a workstation CPU (PyTorch [58]), against the NUTS fit’s hours.

H.3 NUTS versus variational inference

The two engines optimize the same model but differ in what they compute and what they cost (Table H7). NUTS draws asymptotically exact samples from the full joint posterior, delivering calibrated uncertainty for every quantity; SVI returns the best approximation *within a chosen family*, which is fast but biased by the family. The load-bearing consequence here is specific and predictable: a mean-field (diagonal) q_i cannot represent posterior covariance *between* factors, so to fit the data it shrinks the prior inter-factor correlations the likelihood would otherwise support. The result is faithful factor loadings and patient coordinates—which are governed by the posterior *means*—but systematically attenuated inter-factor correlations Φ and underestimated coordinate standard deviations, both governed by the posterior *covariance*. This is why we keep NUTS as the authority for Φ magnitudes and for any credible interval, and use the variational engine for the map, the coordinates, fast reruns and synthetic-cohort generation.

Table H7: NUTS versus variational inference for the FACE-ATLAS measurement model.

	NUTS (main fit)	Variational GLLVM
Inference	MCMC: asymptotically exact samples from the full joint posterior	Optimization: best approximation within a chosen posterior family
Latent treatment	Continuous block marginalized (Woodbury); latents only for discrete items	Explicit per-patient latent for every factor; all ≈ 142 items
Global parameters	Full posterior (loadings, Φ , residuals) with credible intervals	MAP point estimates; loading bands from a seed ensemble
Coordinate uncertainty	Calibrated posterior SD per axis	Relative reliability signal (mean-field underestimates SD)
Inter-factor Φ	Posterior magnitudes with uncertainty	Structure correct; off-diagonals attenuated (mean-field bias)
Cost	Hours (multi-chain)	≈ 4.5 min CPU ($\sim 100\times$ faster)

H.4 Comparison results

On the eight-factor operational map, the variational engine reproduces the NUTS fit on its two load-bearing objects—the measurement map and the patient coordinates—and matches it as a generative model.

The measurement map. Loading columns are congruent on all eight factors (Tucker φ : G 0.957, every specific axis 0.96–0.999; Table H8), and the full item $\times$ factor loading scatter is $r = 0.993$ over 208 cells with no sign flips (Figure H8). G is the hardest axis—it carries every item’s small bifactor loading—and is the only one near the major-axis bar; a richer posterior raises it to 0.974. *It is the same map.*

The patient coordinates. Per-patient coordinates correlate with the NUTS coordinates at $r \geq 0.90$ on all eight axes (Table H9: G 0.990, immunometabolic 0.998, cognition 0.992, down to substance 0.908, the orthogonal, instrument-poor axis). *Patients land in the same place.*

Generative fidelity. Drawing 9,013 synthetic patients from the fitted map and comparing to the observed data, the marginals match (Kolmogorov–Smirnov median 0.04; per-variable mean fidelity $r = 1.000$, SD fidelity $r = 0.998$) and so does the joint covariance (correlation SRMR 0.078, matching the NUTS fit’s ≈ 0.07)—the eight factors reproduce the data’s correlation structure, not merely its margins.

Table H8: Loading congruence (Tucker φ) vs NUTS.

Factor	cells	Tucker φ
G (overall severity)	106	0.957
cognition	16	0.998
immunometabolic	37	0.998
sleep	12	0.999
suicidality	12	0.998
developmental risk	21	0.998
mania/activation	2	0.984
substance	2	0.963

Table H9: Coordinate agreement (Pearson r) vs NUTS.

Factor	n (well)	r
G (overall severity)	8641	0.990
cognition	6451	0.992
immunometabolic	8825	0.998
sleep	7522	0.968
suicidality	8285	0.984
developmental risk	8767	0.986
mania/activation	6152	0.976
substance	4249	0.908

The one gap: inter-factor Φ . The correlation structure is reproduced— G and substance rows are exactly zero in both engines, and the sleep/suicidality/developmental-risk/mania block is positively correlated in both (Figure H9)—but the off-diagonals are systematically attenuated: correlated-block mean $|\phi|$ is 0.083 (variational) versus 0.105 (NUTS), a $\sim 21\%$ shrinkage concentrated in the small negative cognition correlations (the substantive immunometabolic $\leftrightarrow$ cognition coupling is preserved). Climbing the full posterior-richness ladder—mean-field $\rightarrow$ low-rank $\rightarrow$ full $K \times K$ covariance—reduces the shrinkage monotonically ($21\% \rightarrow 20\% \rightarrow 18\%$) but does not close it, establishing the residual as a *structural bias of variational inference on the inter-factor correlation hyperparameter* rather than a tunable defect (an exploratory arm removing the Φ penalty and loosening gradient clipping left Φ unchanged). This is precisely the quantity for which NUTS is retained as the authority; the variational Φ should be read as directionally correct but not as a magnitude.

Summary. The variational GLLVM is a calibrated, $\sim 100\times$ -faster stand-in for the NUTS measurement map and patient coordinates (loadings Tucker 8/8, scatter $r = 0.993$; coordinates $r \geq 0.90$ on 8/8 axes) and a faithful generative model of the cohort, with the inter-factor correlation Φ as the one structurally attenuated quantity for which NUTS remains the authority. It is suited to fast reruns, sensitivity sweeps, synthetic-cohort generation, and as an independent-estimator robustness check on the map; developing it into a fully calibrated Bayesian engine is future work.

References

- [1] Thomas Insel, Bruce Cuthbert, Marjorie Garvey, Robert Heinssen, Daniel S Pine, Kevin Quinn, Charles Sanislow, and Philip Wang. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiatry*, 167(7): 748–51, 2010. doi: 10.1176/appi.ajp.2010.09091379. PMID: 20595427.
- [2] Roman Kotov, Robert F Krueger, David Watson, Thomas M Achenbach, Robert R Althoff, R Michael Bagby, Timothy A Brown, William T Carpenter, Avshalom Caspi, Lee Anna Clark, Nicholas R Eaton, Miriam K Forbes, Kelsie T Forbush, David Goldberg, Deborah Hasin, Steven E Hyman, Masha Y Ivanova, Donald R Lynam, Kristian Markon, Joshua D Miller, Terrie E Moffitt, Leslie C Morey, Stephanie N Mullins-Sweatt, Johan Ormel, Christopher J Patrick, Darrel A Regier, Leslie Rescorla, Camilo J Ruggero, Douglas B Samuel, Martin Sellbom, Leonard J Simms, Andrew E Skodol, Tim Slade, Susan C South, Jennifer L Tackett, Irwin D Waldman, Monika A Waszczuk, Thomas A Widiger, Aidan G C Wright, and Mark Zimmerman. The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *J Abnorm Psychol*, 126(4):454–477, 2017. doi: 10.1037/abn0000258. PMID: 28333488.




Figure H8: **The variational map is the NUTS map.** Item $\times$ factor loadings from the variational GLLVM against the NUTS copula fit ($n = 208$ cells, $r = 0.993$), coloured by cell type. Large home and G -anchor loadings align on the diagonal; cross, window and small bifactor- G cells cluster tightly near zero.

- [3] Camilo J Ruggero, Roman Kotov, Christopher J Hopwood, Michael First, Lee Anna Clark, Andrew E Skodol, Stephanie N Mullins-Sweatt, Christopher J Patrick, Bo Bach, David C Cicero, Anna Docherty, Leonard J Simms, R Michael Bagby, Robert F Krueger, Jennifer L Callahan, Michael Chmielewski, Christopher C Conway, Barbara De Clercq, Allison Dornbach-Bender, Nicholas R Eaton, Miriam K Forbes, Kelsie T Forbush, John D Haltigan, Joshua D Miller, Leslie C Morey, Praveetha Patalay, Darrel A Regier, Ulrich Reininghaus, Alexander J Shackman, Monika A Waszczuk, David Watson, Aidan G C Wright, and Johannes Zimmermann. Integrating the Hierarchical Taxonomy of Psychopathology (HiTOP) into clinical practice. *J Consult Clin Psychol*, 87(12):1069–1084, 2019. doi: 10.1037/ccp0000452. PMID: 31724426.
- [4] Avshalom Caspi, Renate M Houts, Daniel W Belsky, Sidra J Goldman-Mellor, HonaLee Harrington, Salomon Israel, Madeline H Meier, Sandhya Ramrakha, Idan Shalev, Richie Poulton, and Terrie E Moffitt. The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders? *Clin Psychol Sci*, 2(2):119–137, 2014. doi: 10.1177/2167702613497473. PMID: 25360393.
- [5] Benjamin B Lahey, David H Zald, Scott F Perkins, Victoria Villalta-Gil, Katherine B Werts, Carol A Van Hulle, Paul J Rathouz, Brooks Applegate, Quetzal A Class, Holly E Poore, Ashley L Watts, and Irwin D Waldman. Measuring the hierarchical general factor model of psychopathology in young adults. *Int J Methods Psychiatr Res*, 27(1), 2017. doi: 10.1002/mpr.1593. PMID: 28990308.
- [6] Manuel Heinrich, Christian Geiser, Pavle Zagorscak, G Leonard Burns, Johannes Bohn, Stephen P Becker, Michael Eid, Theodore P Beauchaine, and Christine Knaevelsrud. On the Meaning of the “Factor” in Symmetrical Bifactor Models of Psychopathology:




Figure H9: **Inter-factor correlation Φ : structure reproduced, magnitudes attenuated.** Variational Φ (left), NUTS Φ (middle), and their difference (right). The block-diagonal orthogonality (G , substance) is exact in both and the correlated specific block is positive in both; the difference panel localizes the $\sim 21\%$ attenuation to the small cognition cross-correlations.

Recommendations for Future Research From the Bifactor-(-1) Perspective. *Assessment*, 30(3):487–507, 2021. doi: 10.1177/10731911211060298. PMID: 34861784.

- [7] Ashley L Watts, Bridget A Makol, Isabella M Palumbo, Andres De Los Reyes, Thomas M Olino, Robert D Latzman, Colin G DeYoung, Phillip K Wood, and Kenneth J Sher. How Robust Is the p Factor? Using Multitrait-Multimethod Modeling to Inform the Meaning of General Factors of Youth Psychopathology. *Clin Psychol Sci*, 10(4):640–661, 2021. doi: 10.1177/21677026211055170. PMID: 36090949.
- [8] Andre F Marquand, Iead Rezek, Jan Buitelaar, and Christian F Beckmann. Understanding Heterogeneity in Clinical Cohorts Using Normative Models: Beyond Case-Control Studies. *Biol Psychiatry*, 80(7):552–61, 2016. doi: 10.1016/j.biopsych.2015.12.023. PMID: 26927419.
- [9] Thomas Wolfers, Nhat Trung Doan, Tobias Kaufmann, Dag Alnæs s, Torgeir Moberget, Ingrid Agartz, Jan K Buitelaar, Torill Ueland, Ingrid Melle, Barbara Franke, Ole A Andreassen, Christian F Beckmann, Lars T Westlye, and Andre F Marquand. Mapping the Heterogeneous Phenotype of Schizophrenia and Bipolar Disorder Using Normative Models. *JAMA Psychiatry*, 75(11):1146–1155, 2018. doi: 10.1001/jamapsychiatry.2018.2467. PMID: 30304337.
- [10] Ashlea Segal, Linden Parkes, Kevin Aquino, Seyed Mostafa Kia, Thomas Wolfers, Barbara Franke, Martine Hoogman, Christian F Beckmann, Lars T Westlye, Ole A Andreassen, Andrew Zalesky, Ben J Harrison, Christopher G Davey, Carles Soriano-Mas, Narcís Cardoner, Jeggan Tiego, Murat Yücel, Leah Braganza, Chao Suo, Michael Berk, Sue Cotton, Mark A Bellgrove, Andre F Marquand, and Alex Fornito. Regional, circuit and network heterogeneity of brain abnormalities in psychiatric disorders. *Nat Neurosci*, 26(9):1613–1629, 2023. doi: 10.1038/s41593-023-01404-6. PMID: 37580620.
- [11] Brenda W J H Penninx, Femke Lamers, Rick Jansen, Michael Berk, Golam M Khandaker, Livia De Picker, and Yuri Milaneschi. Immuno-metabolic depression: from concept to implementation. *Lancet Reg Health Eur*, 48:101166, 2024. doi: 10.1016/j.lanepe.2024.101166. PMID: 39801616.
- [12] Yuri Milaneschi, Femke Lamers, Michael Berk, and Brenda W J H Penninx. Depression Heterogeneity and Its Biological Underpinnings: Toward Immunometabolic Depression. *Biol Psychiatry*, 88(5):369–380, 2020. doi: 10.1016/j.biopsych.2020.01.014. PMID: 32247527.

- [13] Femke Lamers, Yuri Milaneschi, Christiaan H Vinkers, Robert A Schoevers, Erik J Giltay, and Brenda W J H Penninx. Depression profilers and immuno-metabolic dysregulation: Longitudinal results from the NESDA study. *Brain Behav Immun*, 88:174–183, 2020. doi: 10.1016/j.bbi.2020.04.002. PMID: 32272220.
- [14] Davy Vancampfort, Brendon Stubbs, Alex J Mitchell, Marc De Hert, Martien Wampers, Philip B Ward, Simon Rosenbaum, and Christoph U Correll. Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. *World Psychiatry*, 14(3):339–47, 2015. doi: 10.1002/wps.20252. PMID: 26407790.
- [15] Toby Pillinger, Katherine Beck, Cristian Gobjila, Jacek G Donocik, Sameer Jauhar, and Oliver D Howes. Impaired Glucose Homeostasis in First-Episode Schizophrenia: A Systematic Review and Meta-analysis. *JAMA Psychiatry*, 74(3):261–269, 2017. doi: 10.1001/jamapsychiatry.2016.3803. PMID: 28097367.
- [16] Benjamin Ian Perry, Rachel Upthegrove, Andrew Thompson, Steven Marwaha, Stanley Zammit, Swaran Preet Singh, and Golam Khandaker. Dysglycaemia, Inflammation and Psychosis: Findings From the UK ALSPAC Birth Cohort. *Schizophr Bull*, 45(2):330–338, 2019. doi: 10.1093/schbul/sby040. PMID: 29635418.
- [17] Brett A Clementz, John A Sweeney, Jordan P Hamm, Elena I Ivleva, Lauren E Ethridge, Godfrey D Pearlson, Matcheri S Keshavan, and Carol A Tamminga. Identification of Distinct Psychosis Biotypes Using Brain-Based Biomarkers. *Am J Psychiatry*, 173(4):373–84, 2016. doi: 10.1176/appi.ajp.2015.14091200. PMID: 26651391.
- [18] Brett A Clementz, Rebekah L Trotti, Godfrey D Pearlson, Matcheri S Keshavan, Elliot S Gershon, Sarah K Keedy, Elena I Ivleva, Jennifer E McDowell, and Carol A Tamminga. Testing Psychosis Phenotypes From Bipolar-Schizophrenia Network for Intermediate Phenotypes for Clinical Application: Biotype Characteristics and Targets. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 5(8):808–818, 2020. doi: 10.1016/j.bpsc.2020.03.011. PMID: 32600898.
- [19] Brett A Clementz, Michal Assaf, John A Sweeney, Elliot S Gershon, Sarah K Keedy, S Kristian Hill, Elena I Ivleva, Carol A Tamminga, Jennifer E McDowell, Matcheri S Keshavan, Robert D Gibbons, William T Carpenter, and Godfrey D Pearlson. Categorical and Dimensional Approaches for Psychiatric Classification and Treatment Targeting: Considerations from Psychosis Biotypes. *Adv Neurobiol*, 40:685–723, 2024. doi: 10.1007/978-3-031-69491-2_23. PMID: 39562461.
- [20] Steven P Reise. Invited Paper: The Rediscovery of Bifactor Measurement Models. *Multivariate Behav Res*, 47(5):667–696, 2012. doi: 10.1080/00273171.2012.715555. PMID: 24049214.
- [21] Tihomir Asparouhov and Bengt Muthén. Exploratory structural equation modeling. *Structural Equation Modeling*, 16(3):397–438, 2009. doi: 10.1080/10705510903008204.
- [22] Bengt Muthén and Tihomir Asparouhov. Bayesian structural equation modeling: A more flexible representation of substantive theory. *Psychological Methods*, 17(3):313–335, 2012. doi: 10.1037/a0026802.
- [23] Wes Bonifay and Li Cai. On the complexity of item response theory models. *Multivariate Behav Res*, 52(4):465–484, 2017. doi: 10.1080/00273171.2017.1309262.

- [24] Ashley L Greene, Nicholas R Eaton, Kaiqiao Li, Miriam K Forbes, Robert F Krueger, Kristian E Markon, Irwin D Waldman, David C Cicero, Christopher C Conway, Anna R Docherty, Eiko I Fried, Masha Y Ivanova, Katherine G Jonas, Robert D Latzman, Christopher J Patrick, Ulrich Reininghaus, Jennifer L Tackett, Aidan G C Wright, and Roman Kotov. Are fit indices used to test psychopathology structure biased? a simulation study. *J Abnorm Psychol*, 128(7):740–764, 2019. doi: 10.1037/abn0000434.
- [25] David A Parker, Rebekah L Trotti, Jennifer E McDowell, Matcheri S Keshavan, Godfrey D Pearlson, Elliot S Gershon, Sarah K Keedy, Elena I Ivleva, S Kristian Hill, John A Sweeney, Brett A Clementz, and Carol A Tamminga. Differentiating biomarker features and familial characteristics of B-SNIP psychosis biotypes. *Transl Psychiatry*, 15:234, 2025. doi: 10.1038/s41398-025-03501-5.
- [26] Nick Haslam, Elise Holland, and Peter Kuppens. Categories versus dimensions in personality and psychopathology: a quantitative review of taxometric research. *Psychol Med*, 42(5): 903–920, 2012. doi: 10.1017/S0033291711001966.
- [27] Nanfang Pan, Yajing Long, Kun Qin, Isaac Z Pope, Qiuxing Chen, Ziyu Zhu, Ying Cao, Lei Li, Manpreet K Singh, Robert K McNamara, Melissa P DelBello, Ying Chen, Alex Fornito, and Qiyong Gong. Mapping ADHD Heterogeneity and Biotypes by Topological Deviations in Morphometric Similarity Networks. *JAMA Psychiatry*, 83(5):478–490, 2026. doi: 10.1001/jamapsychiatry.2026.0001. PMID: 41739459.
- [28] Nils Kappelmann, Janine Arloth, Marios K Georgakis, Darina Czamara, Nicolas Rost, Symen Ligthart, Golam M Khandaker, and Elisabeth B Binder. Dissecting the Association Between Inflammation, Metabolic Dysregulation, and Specific Depressive Symptoms: A Genetic Correlation and 2-Sample Mendelian Randomization Study. *JAMA Psychiatry*, 78(2):161–170, 2021. doi: 10.1001/jamapsychiatry.2020.3436. PMID: 33079133.
- [29] Sarah A Norton, Aaron J Gorelik, Sarah E Paul, Emma C Johnson, and Ryan Bogdan. A phenome-wide association study (PheWAS) of genetic risk for C-reactive protein in children of european ancestry: Results from the ABCD study. *Brain Behav Immun*, 127:451–462, 2025. doi: 10.1016/j.bbi.2025.04.012.
- [30] Zheng Ye, Nils Kappelmann, Sylvain Moser, George Davey Smith, Stephen Burgess, Peter B Jones, and Golam M Khandaker. Role of inflammation in depression and anxiety: tests for disorder specificity, linearity and potential causality of association in the UK Biobank. *EClinicalMedicine*, 38:100992, 2021. doi: 10.1016/j.eclinm.2021.100992.
- [31] Yuri Milaneschi, Nils Kappelmann, Zheng Ye, Femke Lamers, Sylvain Moser, Peter B Jones, Stephen Burgess, Brenda W J H Penninx, and Golam M Khandaker. Association of inflammation with depression and anxiety: evidence for symptom-specificity and potential causality from UK Biobank and NESDA cohorts. *Mol Psychiatry*, 26(12):7393–7402, 2021. doi: 10.1038/s41380-021-01188-w.
- [32] Davy Vancampfort, Christoph U Correll, Britta Galling, Michel Probst, Marc De Hert, Philip B Ward, Simon Rosenbaum, Fiona Gaughran, John Lally, and Brendon Stubbs. Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. *World Psychiatry*, 15(2):166–74, 2016. doi: 10.1002/wps.20309. PMID: 27265707.
- [33] Nathalia Garrido-Torres, Miguel Ruiz-Veguilla, Luis Alameda, Manuel Canal-Rivero, María Juncal Ruiz, Marcos Gómez-Revuelta, Rosa Ayesa-Arriola, Ana Rubio-García, Benedicto Crespo-Facorro, and Javier Vázquez-Bourgon. Prevalence of metabolic syndrome and related factors in a large sample of antipsychotic naïve patients with first-episode

- psychosis: Baseline results from the PAFIP cohort. *Schizophr Res*, 246:277–285, 2022. doi: 10.1016/j.schres.2022.07.007. PMID: 35878542.
- [34] Emanuele Felice Osimo, Luke James Baxter, Glyn Lewis, Peter B Jones, and Golam M Khandaker. Prevalence of low-grade inflammation in depression: a systematic review and meta-analysis of CRP levels. *Psychol Med*, 49(12):1958–1970, 2019. doi: 10.1017/S0033291719001454. PMID: 31258105.
 - [35] Brisa S Fernandes, Johann Steiner, Marc L Molendijk, Seetal Dodd, Patricia Nardin, Carlos-Alberto Gonçalves, Felice Jacka, Cristiano A Köhler, Chandan Karmakar, André F Carvalho, and Michael Berk. C-reactive protein concentrations across the mood spectrum in bipolar disorder: a systematic review and meta-analysis. *Lancet Psychiatry*, 3(12):1147–1156, 2016. doi: 10.1016/S2215-0366(16)30370-4. PMID: 27838212.
 - [36] B S Fernandes, J Steiner, H-G Bernstein, S Dodd, J A Pasco, O M Dean, P Nardin, C-A Gonçalves, and M Berk. C-reactive protein is increased in schizophrenia but is not altered by antipsychotics: meta-analysis and implications. *Mol Psychiatry*, 21(4):554–64, 2015. doi: 10.1038/mp.2015.87. PMID: 26169974.
 - [37] F Lamers, N Vogelzangs, K R Merikangas, P de Jonge, A T F Beekman, and B W J H Penninx. Evidence for a differential role of HPA-axis function, inflammation and metabolic syndrome in melancholic versus atypical depression. *Mol Psychiatry*, 18(6):692–9, 2012. doi: 10.1038/mp.2012.144. PMID: 23089630.
 - [38] Yuri Milaneschi, Femke Lamers, Wouter J Peyrot, Bernhard T Baune, Gerome Breen, Abbas Dehghan, Andreas J Forstner, Hans J Grabe, Georg Homuth, Carol Kan, Cathryn Lewis, Niamh Mullins, Matthias Nauck, Giorgio Pistis, Martin Preisig, Margarita Rivera, Marcella Rietschel, Fabian Streit, Jana Strohmaier, Alexander Teumer, Sandra Van der Auwera, Naomi R Wray, Dorret I Boomsma, and Brenda W J H Penninx. Genetic Association of Major Depression With Atypical Features and Obesity-Related Immunometabolic Dysregulations. *JAMA Psychiatry*, 74(12):1214–1225, 2017. doi: 10.1001/jamapsychiatry.2017.3016. PMID: 29049554.
 - [39] M Bryant Howren, Donald M Lamkin, and Jerry Suls. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med*, 71(2):171–86, 2009. doi: 10.1097/PSY.0b013e3181907c1b. PMID: 19188531.
 - [40] Emanuele F Osimo, Toby Pillinger, Irene Mateos Rodriguez, Golam M Khandaker, Carmine M Pariante, and Oliver D Howes. Inflammatory markers in depression: A meta-analysis of mean differences and variability in 5,166 patients and 5,083 controls. *Brain Behav Immun*, 87:901–909, 2020. doi: 10.1016/j.bbi.2020.02.010. PMID: 32113908.
 - [41] Rita Haapakoski, Julia Mathieu, Klaus P Ebmeier, Harri Alenius, and Mika Kivimäki. Cumulative meta-analysis of interleukins 6 and 1 β , tumour necrosis factor α and C-reactive protein in patients with major depressive disorder. *Brain Behav Immun*, 49:206–15, 2015. doi: 10.1016/j.bbi.2015.06.001. PMID: 26065825.
 - [42] Brenda W J H Penninx and Sjors M M Lange. Metabolic syndrome in psychiatric patients: overview, mechanisms, and implications. *Dialogues Clin Neurosci*, 20(1):63–73, 2018. doi: 10.31887/DCNS.2018.20.1/bpenninx. PMID: 29946213.
 - [43] Toby Pillinger, Robert A McCutcheon, and Oliver D Howes. Variability of glucose, insulin, and lipid disturbances in first-episode psychosis: a meta-analysis. *Psychol Med*, 53(7):3150–3156, 2022. doi: 10.1017/S0033291721005213. PMID: 37449481.

- [44] Kateryna Maksyutynska, Nicolette Stogios, Femin Prasad, Jashan Gill, Zaineb Hamza, Riddhita De, Emily Smith, Angelina Horta, Benjamin I Goldstein, Daphne Korczak, Ariel Graff-Guerrero, Margaret K Hahn, and Sri Mahavir Agarwal. Neurocognitive correlates of metabolic dysregulation in individuals with mood disorders: a systematic review and meta-analysis. *Psychol Med*, 54(7):1245–1271, 2024. doi: 10.1017/S0033291724000345. PMID: 38450447.
- [45] Andrea Zagaria, Valeria Fiori, Mariacarla Vacca, Caterina Lombardo, Carmine M Pariante, and Andrea Ballesio. Inflammation as a mediator between adverse childhood experiences and adult depression: A meta-analytic structural equation model. *J Affect Disord*, 357:85–96, 2024. doi: 10.1016/j.jad.2024.04.072. PMID: 38677656.
- [46] Holly F Levin-Aspenson, David Watson, Lee Anna Clark, and Mark Zimmerman. What Is the General Factor of Psychopathology? Consistency of the p Factor Across Samples. *Assessment*, 28(4):1035–1049, 2020. doi: 10.1177/1073191120954921. PMID: 32909442.
- [47] Carlos M Carvalho, Nicholas G Polson, and James G Scott. Handling sparsity via the horseshoe. In *Proceedings of the Twelfth International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 5, pages 73–80, 2009.
- [48] Juho Piironen and Aki Vehtari. Sparsity information and regularization in the horseshoe and other shrinkage priors. *Electronic Journal of Statistics*, 11(2):5018–5051, 2017. doi: 10.1214/17-EJS1337SI.
- [49] Matthew D Hoffman and Andrew Gelman. The no-u-turn sampler: Adaptively setting path lengths in hamiltonian monte carlo. *Journal of Machine Learning Research*, 15:1593–1623, 2014.
- [50] Oriol Abril-Pla, Virgile Andreani, Colin Carroll, Larry Dong, Christopher J Fonnesbeck, Maxim Kochurov, Ravin Kumar, Junpeng Lao, Christian C Luhmann, Osvaldo A Martin, Michael Osthege, Ricardo Vieira, Thomas Wiecki, and Robert Zinkov. PyMC: a modern, and comprehensive probabilistic programming framework in Python. *PeerJ Computer Science*, 9:e1516, 2023. doi: 10.7717/peerj-cs.1516.
- [51] Du Phan, Neeraj Pradhan, and Martin Jankowiak. Composable effects for flexible and accelerated probabilistic programming in NumPyro. *arXiv preprint arXiv:1912.11554*, 2019. doi: 10.48550/arXiv.1912.11554.
- [52] Wes Bonifay, Sean P Lane, and Steven P Reise. Three concerns with applying a bifactor model as a structure of psychopathology. *Clin Psychol Sci*, 5(1):184–186, 2017. doi: 10.1177/2167702616657069.
- [53] Sumio Watanabe. Asymptotic equivalence of bayes cross validation and widely applicable information criterion in singular learning theory. *Journal of Machine Learning Research*, 11:3571–3594, 2010.
- [54] Aki Vehtari, Andrew Gelman, and Jonah Gabry. Practical bayesian model evaluation using leave-one-out cross-validation and waic. *Statistics and Computing*, 27(5):1413–1432, 2017. doi: 10.1007/s11222-016-9696-4.
- [55] Francis K C Hui, David I Warton, John T Ormerod, Viivi Haapaniemi, and Sara Taskinen. Variational approximations for generalized linear latent variable models. *J Comput Graph Stat*, 26(1):35–43, 2017. doi: 10.1080/10618600.2016.1164708.
- [56] David M Blei, Alp Kucukelbir, and Jon D McAuliffe. Variational inference: A review for statisticians. *J Am Stat Assoc*, 112(518):859–877, 2017. doi: 10.1080/01621459.2017.1285773.

- [57] Matthew D Hoffman, David M Blei, Chong Wang, and John Paisley. Stochastic variational inference. *J Mach Learn Res*, 14:1303–1347, 2013.
- [58] Adam Paszke, Sam Gross, Francisco Massa, Adam Lerer, et al. PyTorch: An imperative style, high-performance deep learning library. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 32, pages 8024–8035, 2019.
- [59] Diederik P Kingma and Max Welling. Auto-encoding variational Bayes. In *International Conference on Learning Representations (ICLR)*, 2014. doi: 10.48550/arXiv.1312.6114.
- [60] Adele Cutler and Leo Breiman. Archetypal analysis. *Technometrics*, 36(4):338–347, 1994. doi: 10.1080/00401706.1994.10485840.
- [61] Jo Bovy, David W Hogg, and Sam T Roweis. Extreme deconvolution: Inferring complete distribution functions from noisy, heterogeneous and incomplete observations. *Annals of Applied Statistics*, 5(2B):1657–1677, 2011. doi: 10.1214/10-AOAS439.
- [62] Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983. doi: 10.1093/biomet/70.1.41.
- [63] Peter C Austin. An Introduction to Propensity Score Methods for Reducing the Effects of Confounding in Observational Studies. *Multivariate Behav Res*, 46(3):399–424, 2011. doi: 10.1080/00273171.2011.568786. PMID: 21818162.
- [64] Tyler J VanderWeele and Peng Ding. Sensitivity Analysis in Observational Research: Introducing the E-Value. *Ann Intern Med*, 167(4):268–274, 2017. doi: 10.7326/M16-2607. PMID: 28693043.
- [65] Daniel Lewandowski, Dorota Kurowicka, and Harry Joe. Generating random correlation matrices based on vines and extended onion method. *Journal of Multivariate Analysis*, 100(9):1989–2001, 2009. doi: 10.1016/j.jmva.2009.04.008.
- [66] J A Hartigan and P M Hartigan. The dip test of unimodality. *Annals of Statistics*, 13(1): 70–84, 1985. doi: 10.1214/aos/1176346577.
- [67] Robert Tibshirani, Guenther Walther, and Trevor Hastie. Estimating the number of clusters in a data set via the gap statistic. *Journal of the Royal Statistical Society: Series B*, 63(2): 411–423, 2001. doi: 10.1111/1467-9868.00293.
- [68] Ricardo J G B Campello, Davoud Moulavi, and Jörg Sander. Density-based clustering based on hierarchical density estimates. In *Advances in Knowledge Discovery and Data Mining (PAKDD)*, volume 7819 of *Lecture Notes in Computer Science*, pages 160–172, 2013. doi: 10.1007/978-3-642-37456-2_14.
- [69] Gurjeet Singh, Facundo Mémoli, and Gunnar Carlsson. Topological methods for the analysis of high dimensional data sets and 3D object recognition. In *Eurographics Symposium on Point-Based Graphics*, pages 91–100, 2007. doi: 10.2312/SPBG/SPBG07/091-100.
- [70] Leland McInnes, John Healy, and James Melville. UMAP: Uniform manifold approximation and projection for dimension reduction. *arXiv preprint arXiv:1802.03426*, 2018. doi: 10.48550/arXiv.1802.03426.