

Given  $f, g \in L^p(\mu)$ , and we investigate

$$\phi(t) = \int_X |f + tg|^p d\mu.$$

One can expect that  $\Phi(t)$  is differentiable because the integrand is continuous. First, let me give an example

$$\phi(t) = \int_0^{2\pi} |\sin(x) + t \cos(x)|^2 d\mu,$$

which is illustrated in the following figure.




Figure 1:

To prove this theoretically, we take  $t = 0$  as example

$$\begin{aligned} \phi'(0) &= \lim_{t \rightarrow 0} \frac{\int_X |f + tg|^p - |f|^p}{t} \\ &= \int_X \lim_{t \rightarrow 0} \frac{|f + tg|^p - |f|^p}{t} d\mu. \end{aligned}$$

Notice that we need some convergence theorem to change the order of  $\lim_{t \rightarrow 0}$  and  $\int_X$ .

It's the convexity of  $|a + bt|^p, p \geq 1$  that make it available. For  $t \in (-0.5, 0.5)$ , we have

$$\frac{|f + tg|^p - |f|^p}{t} \leq \frac{|f + g|^p - |f|^p}{1 - 0} = |f + g|^p - |f|^p.$$

and it's easy to see that  $|f + g|^p - |f|^p \in L^1(\mu)$  by the simple inequality  $|a + b|^p \leq 2^{p-1}(|a|^p + |b|^p)$ .

Hence we apply L Hospital's rule to get

$$\phi'(0) = p \int_X |f|^{p-2} f g \, d\mu.$$

**Convex function.** Here I wanna write something about convex function.

- The simplest convex function which is not differentiable is  $|x|$ . More generally, a convex function can consist of connected line segments, also known as a piecewise linear function. Interestingly, we can obtain such a function by connecting the tangent lines of a convex function or connecting the points  $(x_i, f(x_i))$  where  $f$  is convex.