

**Physics 110A, Spring 2021**  
**Solution to Homework 4**  
**GSI: Yi-Chuan Lu**

1. (a) Due to spherical symmetry, without loss of generality, we may assume the observer (field point) is on the  $z$  axis at a distance  $r$  away from the center. Then

$$V(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_0^\pi \int_0^{2\pi} \frac{\sigma R^2 \sin\theta d\phi d\theta}{\sqrt{r^2 + R^2 - 2rR\cos\theta}} = \frac{\sigma}{2\epsilon_0} \frac{R}{r} \sqrt{r^2 + R^2 - 2rR\cos\theta} \Big|_0^\pi$$

$$= \frac{\sigma}{2\epsilon_0} \frac{R}{r} [(r+R) - |r-R|] = \begin{cases} \frac{\sigma}{\epsilon_0} \frac{R^2}{r}, & r > R, \\ -R, & r < R. \end{cases}$$

(b) The electric field is  $\mathbf{E}(\mathbf{r}) = -\nabla V(\mathbf{r}) = -\frac{\partial V}{\partial r} \hat{\mathbf{r}} = \begin{cases} \frac{\sigma}{\epsilon_0} \frac{R^2}{r^2} \hat{\mathbf{r}}, & r > R, \\ \mathbf{0}, & r < R. \end{cases}$

2. Note that the electric field inside the shell is zero, so the integral over the entire space reduces to that over the region just outside the spherical shell:

$$W = \frac{\epsilon_0}{2} \int_{\text{all space}} E^2 d\tau = \frac{\epsilon_0}{2} \int_{\text{outside}} E^2 d\tau$$

$$= \frac{\epsilon_0}{2} \int_R^\infty \int_0^\pi \int_0^{2\pi} \left( \frac{\sigma}{\epsilon_0} \frac{R^2}{r^2} \right)^2 r^2 \sin\theta d\phi d\theta dr = \frac{2\pi\sigma^2 R^3}{\epsilon_0}.$$

3. (a) We can either assume  $\sigma$  is constant or assume the total charge  $Q \equiv 4\pi R^2 \sigma$  is constant when the shell expands. The former is more complicated because we need to connect the shell to, e.g., a battery to provide charge, and therefore we will have to take into account the electric work done by the battery<sup>1</sup>. So let's assume  $Q = 4\pi R^2 \sigma$  is constant. In terms of  $Q$ , the energy  $W$  is expressed as

$$W = \frac{2\pi R^3}{\epsilon_0} \left( \frac{Q}{4\pi R^2} \right)^2 = \frac{Q^2}{8\pi\epsilon_0 R},$$

so the change in  $W$  at constant  $Q$  is

$$\delta W = \frac{dW}{dR} \delta R = -\frac{Q^2}{8\pi\epsilon_0 R^2} \delta R = \frac{2\pi R^2 \sigma^2}{\epsilon_0} \delta R.$$

- (b) By letting  $\delta W = -f(4\pi R^2) \delta R$ , we obtain

$$f = -\frac{1}{4\pi R^2} \frac{\delta W}{\delta R} = \frac{\sigma^2}{2\epsilon_0}.$$

Note that this is consistent with what we would obtain by using Griffiths equation (2.50):

$$\mathbf{f} = \sigma \frac{\mathbf{E}_{\text{above}} + \mathbf{E}_{\text{below}}}{2} = \sigma \left( \frac{\sigma}{2\epsilon_0} \hat{\mathbf{n}} \right).$$

---

<sup>1</sup>There will be a discussion on this matter in Griffiths section 4.4.4.