PHYS 3031 Course Notes

Mathematical Methods in Physics II

LIN, Xuanyu

MATH METHODS IN PHYSICS

PHYS 3031 Mathematical Methods in Physics II



September 6, 2023



1 Series

1.1 Convergence Condition for Positive Series $\sum\limits_{n=1}^{\infty}a_n$

Necessary condition: $\lim_{N\to\infty} a_N = 0$ Hierarchy: $N! > a^N > N^b > \ln N$

Sterling Formula $\ln N! \approx N \ln N - N \approx N \ln N$

Comparison Test 1 $\sum a_n < \sum b_n,\, b \text{ converges} \to a \text{ converges}$

Comparison Test 2 (Integral Test) $\sum a_n \& \int a(n) dn$ share the same fate

Ratio Test $\lim_{n\to\infty}\frac{a_{n+1}}{a_n}=\rho,\, \rho>1\to \text{Diverges},\, \rho<1\to \text{Converges}$

Extended (Special) Comparison Test $\lim_{n\to\infty}\frac{a_n}{b_n}=1$, then $\sum a_n \& \sum b_n \mathrm{d}n$ share the same fate



Homework 1

Problem 1

Use the comparison test to prove the convergence of the following series:

(a)
$$\sum_{n=1}^{\infty} \frac{1}{2^n + 3^n}$$

(b)
$$\sum_{n=1}^{\infty} \frac{1}{n2^n}$$

Solution.

(a) By considering the n-th term, we have

$$\frac{1}{2^n + 3^n} < \frac{1}{2^n + 2^n} = \frac{1}{2^{n+1}}$$

Since the series is positive and that

$$\sum_{n=1}^{\infty} \frac{1}{2^{n+1}} = \frac{1}{2} < \infty,$$

the original series converges.

(b) Consider the n-th term, we have

$$\frac{1}{n2^n} < \frac{1}{2^n}$$

for $n \geq 2$.

Since the series is positive and that

$$\sum_{n=1}^{\infty} \frac{1}{n2^n} < \frac{1}{2} + \sum_{n=2}^{\infty} \frac{1}{2^n} = 1 < \infty,$$

the original sequence converges.

Test the following series for convergence using the comparison test:

(a)
$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$$

(b)
$$\sum_{n=2}^{\infty} \frac{1}{\ln n}$$

Solution.

(a) Consider the n-th term, we have

$$\frac{1}{\sqrt{n}} > \frac{1}{n}$$

for $n \geq 2$.

Since

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} > 1 + \sum_{n=2}^{\infty} \frac{1}{n} \to \infty,$$

the series diverges by the comparison test.

(b) Since $\ln n < n$ for $\forall n \in \mathbb{N}$,

$$\sum_{n=2}^{\infty} \frac{1}{\ln n} > \sum_{n=2}^{\infty} \frac{1}{n} \to \infty$$

Thus, the series diverges by the comparison test.



Use the integral test to find whether the following series converge or diverge:

(a)
$$\sum_{n=2}^{\infty} \frac{1}{n \ln n}$$

(b)
$$\sum_{n=1}^{\infty} \frac{e^n}{e^{2n} + 9}$$

Solution.

(a) Consider the integral

$$\int_{2}^{\infty} \frac{1}{x \ln x} dx = \int_{\ln 2}^{\infty} \frac{d \ln x}{\ln x} = \ln \ln x \Big|_{2}^{\infty} \to \infty$$

Thus, the original series diverges.

(b) Consider the integral

$$\int_{2}^{\infty} \frac{e^{x}}{e^{2x}+9} \mathrm{d}x = \int_{2}^{\infty} \frac{1}{e^{x}+\frac{9}{e^{x}}} \mathrm{d}x \geq \int_{2}^{\infty} \frac{1}{2e^{x}} = -\frac{1}{2} e^{-x} \big|_{2}^{\infty} = \frac{e^{-2}}{2} < \infty$$

Thus, the original series converges.



Use the ratio test to find whether the following series converge or diverge: $\sum_{n=0}^{\infty} \frac{(n!)^3 e^{3n}}{(3n)!}$

Solution.

The n-th term si given by

$$a_n = \frac{(n!)^3 e^{3n}}{(3n)!}$$

Thus,

$$\frac{a_{n+1}}{a_n} = \frac{[(n+1)!]^3 e^{3(n+1)}}{[3(n+1)]!} \frac{(3n)!}{(n!)^3 e^{3n}} = \frac{(n+1)^3 \cdot e^3}{(3n+1)(3n+2)(3n+3)} = e^3 \cdot \frac{n^3 + 3n^2 + 3n + 1}{27n^3 + 54n^2 + 33n + 6}$$

$$\lim_{n \to \infty} \frac{a_{n+1}}{a_n} = \lim_{n \to \infty} e^3 \cdot \frac{n^3 + 3n^2 + 3n + 1}{27n^3 + 54n^2 + 33n + 6} = \lim_{n \to \infty} e^3 \cdot \frac{1 + \frac{3}{n} + \frac{3}{n^2} + \frac{1}{n^3}}{27 + \frac{54}{n} + \frac{33}{n^2} + \frac{6}{n^3}} = \frac{e^3}{27} < 1$$

Thus, the series converges by the ratio test.

Use the special comparison test to find whether the following series converge or diverge:

(a)
$$\sum_{n=9}^{\infty} \frac{(2n+1)(3n-5)}{\sqrt{n^2-73}}$$

(b)
$$\sum_{n=3}^{\infty} \frac{(n-\ln n)^2}{5n^4-3n^2+1}$$

(c)
$$\sum_{n=1}^{\infty} \frac{\sqrt{n^3 + 5n - 1}}{n^2 - \sin n^3}$$

Solution.

(a)

$$a_n = \frac{(2n+1)(3n-5)}{\sqrt{n^2-73}}$$

Consider another sequence with $b_n=6n.$ Obviously, $\sum b_n$ diverges.

$$\lim_{n \to \infty} \frac{a_n}{b_n} = \lim_{n \to \infty} \frac{(2n+1)(3n-5)}{6n\sqrt{n^2-73}} = \lim_{n \to \infty} \frac{6-\frac{7}{n}-\frac{5}{n^2}}{6\sqrt{1-\frac{73}{n^2}}} = 1$$

Thus, $\sum a_n$ also diverges.

(b)

$$a_n = \frac{(n - \ln n)^2}{5n^4 - 3n^2 + 1}$$

Consider another sequence with $b_n = \frac{1}{5n^2}$. Obviously, $\sum b_n$ converges.

$$\lim_{n \to \infty} \frac{a_n}{b_n} = \lim_{n \to \infty} \frac{5n^2(n - \ln n)^2}{5n^4 - 3n^2 + 1} = \lim_{n \to \infty} \frac{(1 - \frac{\ln n}{n})^2}{1 - \frac{3}{5n^2} + \frac{1}{5n^4}} = 1$$

Thus, $\sum a_n$ also converges.

(c)

$$a_n = \frac{\sqrt{n^3 + 5n - 1}}{n^2 - \sin n^3}$$

Consider another sequence with $b_n = \frac{1}{n^{\frac{1}{2}}}$. Then $\sum b_n$ diverges from the previous problem.

$$\lim_{n \to \infty} \frac{a_n}{b_n} = \lim_{n \to \infty} \frac{\sqrt{n^4 + 5n^2 - n}}{n^2 - \sin n^3} = \lim_{n \to \infty} \frac{\sqrt{1 + \frac{5}{n^2} - \frac{1}{n^3}}}{1 - \frac{\sin n^3}{n^2}} = 1$$

as

$$0=\lim_{n\to\infty}-\frac{1}{n^2}\leq\lim_{n\to\infty}\frac{\sin n^3}{n^2}\leq\lim_{n\to\infty}\frac{1}{n^2}=0$$

Thus, $\sum a_n$ also diverges.

Test the following series for convergence:

(a)
$$\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$$

(b)
$$\sum_{n=1}^{\infty} \frac{(-2)^n}{n^2}$$

Solution.

(a) Since

$$\frac{|a_{n+1}|}{|a_n|}=\frac{\sqrt{n}}{\sqrt{n+1}}<1$$

and

$$\lim_{n\to\infty}a_n=\lim_{n\to\infty}\frac{(-1)^n}{\sqrt{n}}=0$$

Thus, the original series converges.

(b)

$$\lim_{n \to \infty} \frac{a_{n+1}}{a_n} = \lim_{n \to \infty} -\frac{2n^2}{(n+1)^2} = \lim_{n \to \infty} -\frac{2}{(\frac{n+1}{n})^2} = -2$$

Thus, the original series diverges.



Homework 2

Problem 7

Find the interval of convergence of the following power series; be sure to investigate the endpoints of the interval: $\sum_{n=1}^{\infty} \frac{(-1)^n (x+1)^n}{n}$



The following series are not power series, but you can transform each one into a power series by a change of variable and so find out where it converges:

- (a) $\sum_{n=2}^{\infty} \frac{(-1)^n x^{n/2}}{n \ln n}$
- (b) $\sum_{n=0}^{\infty} (\sqrt{x^2+1})^n \frac{2^n}{3^n+n^3}$
- (c) $\sum_{n=0}^{\infty} (\sin x)^n (-1)^n 2^n$

- (a)
- (b)
- (c)



Find the first few terms of the Maclaurin series for the following functions and check your results by computer: $\frac{e^x}{1-x}$



Show that $\ln(1-x) = -x$ with an error less than 0.0056 for |x| < 0.1.



Show that $2/\sqrt{4-x} = 1 + \frac{1}{8}x$ with an error less than $\frac{1}{32}$ for 0 < x < 1.



Use power series to evaluate the function at the given point. Compare with computer results, using the computer to find the series, and also to do the problem without series. Resolve any disagreement in results: $e^{\arcsin x} + \ln(\frac{1-x}{e})$ at x = 0.0003



Use Maclaurin series to evaluate each of the following. Although you could do them by computer, you can probably do them in your head faster than you can type them into the computer. So use these to practice quick and skillful use of basic series to make simple calculations: $\frac{d^3}{dx^3}(\frac{x^2e^x}{1-x})$ at x=0



Find the sum of the following series by recognizing it as the Maclaurin series for a function evaluated at a point: $\sum_{n=1}^{\infty} \frac{1}{n2^n}$



Evaluate the following indeterminate forms by using L' Hopital's rule and check your results by computer. (Note that Maclaurin series would not be useful here because x does not tend to zero, or because a function (ln x, for example) is not expandable in a Maclaurin series.): $\lim_{x\to\pi} \frac{x\sin x}{x-\pi}$



Homework 3

Problem 16