

The Breakthrough Listen Search for Intelligent Life in the Universe: Searching for Anomalous Variability in Modern Time-Domain Surveys

Anastasios Tzanidakis^{*1} and James R. A. Davenport¹

¹Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA

1 Introduction

One of the pressing questions in contemporary astronomy is the Search for Extraterrestrial Intelligence (SETI). Highlighted by the Astro 2020 Decadal Survey, SETI

However, SETI has not utilized a new generation of optical wide, fast, and deep time-domain surveys. With the advent of modern ground-based large time-domain surveys, we propose the expansion of frameworks and searches for anomalous variability over a large data volume in real time.

Modern time-domain surveys give us an unprecedented volume of data and parameter space to search for technosignatures in temporal datasets for the Search of Extraterrestrial Intelligence (SETI). For example, large sky surveys have been argued to be the ideal springboard advancing the search for technosignatures (Djorgovski, 2000; Davenport, 2019).

Another strange variability such as ASASSN-21qj, a Sun-like star, that exhibits anomalous long-term variability trends.

For example, (Wright et al., 2016) ...

- Some basic SETI technosignature search assumption - and why time series? (need literature i.e Kipping and Teachy 2016, Socas-Navarro et al. 2022)
- Why OSETI in Modern Time-Domain Surveys? For example, large sky surveys have been argued to be the ideal springboard advancing the search for technosignatures (Djorgovski 2000, Davenport 2019)
- What are the challenges, and things that need to be accomplished ahead?

Based on our own evaluation of the 9-axis of merit developed by Sheikh (2020), searches for technosignatures through variable sources are relevant now.

Until most recently, a systematic search for anomalous stellar variability has not been undertaken as rigorously on wide field-of-view modern time-domain surveys as we proposed in this program. For example, Malanchev et al. (2021) performed a search for anomalous events in the ZTF DR3 that contained 2.5 million objects over the span of nearly two years of ZTF observations. Using a series of machine-learning filtering algorithms, their work demonstrates the ability to discover rare variables and transients. Since the work of Malanchev et al. (2021) the current onset of ZTF DR19 contains over 4.8 billion cataloged sources, and more than 5 years of observations begs the opportunity for the thorough investigation of technosignatures. Other hyper-catalogs exist, such as ZTF Ubercalibration¹, that enables even higher photometric precision for the ZTF photometry pipeline that can be utilized in this work.

We present methods here expanding on this work, enabling more robust anomaly detection, while also implementing state-of-the-art algorithms for catalogs of greater volume.

By now, the canonical search for anomalous stellar variability is often associated with KIC 1223471

By now, the canonical search for "anomalous variability" is more or less described. Even though converging literature suggests the presence of complex dust behavior. KIC 1233 however remains a mystery, given that no other star exhibits. However, other patterns of odd variability could exist such as asymmetric dimming events, prolonged dimming events, complex dimming event profiles, and the overall color-evolution of the variable. **Our program will be the first to place meaningful upper limits to possible detection rates of technosignatures. We will develop a catalog of possible signals in the theme of duration, depth, color, and shape.**

^{*}Corresponding Author: atzanida@uw.edu

¹<http://nesssi.cacr.caltech.edu/ZTF/Web/Zuber.html>

This document is part of the Optical SETI Roadmap document.

In the following sections, we discuss the overall themes and work plan as part of our anomalous variability search using time-domain surveys. The proposed sections will be part of the lead author’s Ph.D thesis.

- Build upon the methods for systematically detecting outliers (Giles and Walkowicz, 2019), and using new methods for comparing variability across the color-magnitude diagram, carry out the most comprehensive search for rare or unique stars to date in the ZTF public survey
- Incorporate established methods to search for technosignature candidates, including searching the “SETI Ellipsoid”, the “Earth Transit Zone”, and for disappearing stars
- Prototype and implement a real-time technosignature alert-based anomaly detection and follow-up strategy for the forthcoming Rubin observatory alerts
- Create an extensible, general-purpose technosignature and outlier software framework, which can be built upon to search for new or novel signal types in future surveys

1.1 Pilot Study and Early Preparation

In preparation for our program, we suggest a proprietary period of a few months including the collection of open-source tools, data, additional sources of funding, follow-up proposals, and any other resources that will be utilized for this project. In each section, we briefly summarize our proprietary focus.

1.1.1 Access to Modern Time-Domain Surveys

Our home institution provides strong expertise in both in-house algorithms and tools for accessing and processing large volumes of time-domain data from large surveys such as Astronomy eXtensions for Spark (AXS) (Zečević et al., 2019). For example, this can be enabled immediately with the LSST Interdisciplinary Network For Collaboration and Computing (LINCC) with the use of the Large Survey Database v2 (LSDB)² a new tool that will enable rapid access to synoptic data. Discussions with engineers will include the ingestion of large astronomical catalogs that will be readily available for our project.

1.2 Open-Source Tools

Our program will require a strong suite of tools to provide a rapid assessment of stellar classification, including the identification of fundamental stellar parameters for further assessment. Here is a short but not fully complete list of tools that could be used, while we examine other options as part of our tool-box: SNAD (Malanchev et al., 2023), AstroARIADNE (Vines and Jenkins, 2022), VOSA (Bayo et al., 2008), MESA (Paxton et al., 2011), STARHorse (Queiroz et al., 2018), SNID (Liu and Modjaz, 2014), CLOUDY (Chatzikos et al., 2023), batman (Kreidberg, 2015). Prior to our technosignature search, we will ensure that all tools are installed, and streamlined to our regular pipelines of investigating astrophysical sources.

1.2.1 Observing Programs and Capabilities

Upon the successful identification of possible **Anomalous Targets of Opportunity** (ATO), follow-up capabilities will enable reliable identification, classification, and further examination of identified variable sources. In particular, optical and near-IR medium-resolution spectroscopic follow-up capabilities can be used as a diagnostic tool to differentiate anomalous candidates from other stochastic variables such as Young Stellar Objects (YSO), Cataclysmic Variables (CVS), and R Coronae Borealis (RcB) stars. As part of our recommended program, both authors will have regular observing programs allocated on the Apache Point Observatory (APO) 3.5m telescope for regular follow-up capabilities with medium-resolution spectrographs and deep multi-band imaging. As a result of our complementary spectroscopic campaign, upon the approval of the Breakthrough Listen project, all spectra can be released publicly either via the Transient Naming Server (TNS), The Astronomer’s Telegram (ATel), or other related services that can enable future studies of our targets.

²<https://github.com/astronomy-commons/lsc2>

83 1.2.2 F-dwarf Search and Limits with ZTF

84 To date, no other documented study has found a star that exhibits similar variability patterns to that of KIC 8462852,
85 although converging literature suggests the complex presence of dust driving such variability (Wright and Sigurdsson,
86 2016). Yet it remains upon debate why such other variables have not been discovered. Tzanidakis et al. (in-prep) are
87 currently undertaking a search for KIC 8462852 F-dwarf analogs using the Zwicky Transient Facility (ZTF) survey
88 Data Release 18 for anomalous variability of F0-9 main sequence stars. Upon the completion of this brief paper, will
89 conclude with one of the largest volume searched sampled of F-dwarfs conducted to date. The preliminary sample
90 contains over 5 million light curves from ZTF that will require extensive investigation including the prototyping of
91 robust algorithms that will identify anomalous variable stars.

92 1.3 Algorithms and Limits

93 In this part of our proposed work, we aim to establish a strong understanding of the limit of our algorithms, including
94 simulated data, and the sensitivity of possible technosignature outlier signals we might be capable of finding. The work
95 presenter here will provide a comprehensive examination and detection limit for the most rich time-domain catalogs
96 such as ZTF, LSST,...

97 Preliminary

- 98 • Injection and recovery analysis for sensitivity estimations using emulated technosignature searches via time-domain
99 schedulers such as LSST OPSIMS, ZTF...
- 100 • Period synchronization search in the era of LSST, ZTF, TESS (for example; Nillpour + 2023; *Signal Synchronization*
101 *Strategies and Time Domain SETI with Gaia DR3*)- via Lomb-Scargle Periodogram - we have the sensitivity
102 and algorithms to test this and report limits. For example, Tzanidakis et al. (dmtn-221) Signal Synchronization
103 Strategies and Time Domain SETI with Gaia DR3 have shown the ability to perform an LSST-scalable approach
104 for recovering significant periods for a large volume of data.
105 For example, the lead author has already shown in dmtn-221 has shown the ability to do such work.
- 106 • Volume sensitivity function enabled through the big surveys.
- 107 • we can generate a grid of interesting light curve with (SHADOW IMAGING OF TRANSITING OBJECTS -
108 Sandford and Kipping, Arnord 2015)

109 1.4 ASTRA-VAR: A Census for Anomalous Temporal Stellar Variability using Time-Domain 110 Surveys

111 A complete and robust search for anomalous stellar variability will be performed with our newly developed anomaly
112 detection outlier time series algorithm. There are several surveys we can explore the use of our algorithms: ZTF DR19,
113 Gaia DR4, and unWISE. All work with prepare for the next generation of the truly exceptional volume of data from
114 the Vera C. Rubin Observatory Legacy Survey of Space and Time (LSST), and Gaia DR4³ that includes the full
115 astrometric, photometric, and radial-velocity catalogs and all epochs and transit data for all sources

116 The search fast-slow variability that does not fall within the normal boundaries of stellar variability...

117 2 Outreach and Community Impact

118 2.1 Undergraduate Research

119 Undergraduate research opportunities are the pillars for inspiring students to pursue careers in STEM. Our proposal
120 has a plethora of SETI-related research projects aimed at all levels of undergraduate career and experience. The skills
121 acquired from our research projects can drastically vary depending on the interest of our students and their level of
122 involvement. For example, projects with a concentration on algorithms and big-data, observational astronomy, and
123 data reduction, and computational work. We strive to expand the diversity of our projects such that it directly matches
124 the interests of our students. Most importantly, our program is committed to prioritizing students from marginalized
125 communities and first-generation college students. This will be achieved directly in partnership with faculty and staff
126 with a specific focus on Diversity, Equity, Inclusion, and Accessibility (DEIA), and our in-house outreach and DEIA
127 committee.

³Not before the end of 2025

2.2 Public Series *Astrobiology and SETI: Are We Alone in the Universe?*

Both authors lead substantial outreach efforts at the University of Washington in the Department of Astronomy. The primary author, A. Tzanidakis, serves as the director and program coordinator for the University of Washington's Astronomy Planetarium and is a co-leading member of the UW Astronomy outreach initiative. **As part of our Breakthrough Listen program, we aim to complement our work with a series of free public talks, planetarium shows, and public observing sessions with a focus on time-domain astronomy and SETI to the greater Washington community.**

Our home institution currently has all the tools and programs in place for such specialized programs. For example, In 2020, our program was successfully awarded over \$70,000 grant from the University of Washington Student Technology Fund (STF), updated with the state-of-the-art 4k projectors system⁴. Our planetarium currently operates on the World Wide Telescope (WWT) (Goodman et al., 2012) platform that enables data-driven astronomical 3D visualizations to create stunning planetarium programs. Other data-driven outreach media include the sonification time-series data (Zanella et al., 2022) by expanding the accessibility of media during our shows. Our suggest program can also be expanded to reach beyond the local Seattle vicinity with the UW Mobile Planetarium⁵.

Similarly, A. Tzanidakis was a co-investigator in collaboration with the undergraduate Astronomy club (League of Astronomers⁶; LOA) for the successful application of another STF proposal over \$18,000 for an H- α solar telescope that can be used as part of this program for daytime solar observations with the public. This will also be reinforced with our in-house telescope gear in partnership with the on-campus Jacobsen Observatory⁷.

2.3 Enabling Broad Interdisciplinary Collaborations

The Search for Extraterrestrial Intelligence in the context of our time-domain program is inherently interdisciplinary. The temporal evolution of astrophysical sources is complex that requires a suite of tools and understanding spanning from data science, statistics, and overall expertise in many domains of astrophysics such as transients, AGN, and various other sub-fields of TDA. The authors propose a journal-club equivalent meeting that will focus on the search for technosignatures. The focus of the meetings will successfully allow all academically staged students and faculty to engage with the SETI literature, but most importantly facilitate collaboration with others. The meetings will be advertised beyond the Astronomy department such as the Astrobiology (AB) division, Earth and Atmospheric Sciences (EAS), and statistics.

3 Summary Timeline

The corresponding lead author, A. Tzanidakis, will be entering his 3rd year of graduate studies at the University of Washington (UW). Here we summarize a rough timeline of the proposed research:

References

- A. Bayo, C. Rodrigo, D. Barrado Y Navascués, E. Solano, R. Gutiérrez, M. Morales-Calderón, and F. Allard. VOSA: virtual observatory SED analyzer. An application to the Collinder 69 open cluster. , 492(1):277–287, December 2008. doi: 10.1051/0004-6361:200810395.
- Marios Chatzikos, Stefano Bianchi, Francesco Camilloni, Priyanka Chakraborty, Chamani M. Gunasekera, Francisco Guzmán, Jonathan S. Milby, Arnab Sarkar, Gargi Shaw, Peter A. M. van Hoof, and Gary J. Ferland. The 2023 release of Cloudy. *arXiv e-prints*, art. arXiv:2308.06396, August 2023. doi: 10.48550/arXiv.2308.06396.
- James. R. A. Davenport. SETI in the Spatio-Temporal Survey Domain. *arXiv e-prints*, art. arXiv:1907.04443, July 2019. doi: 10.48550/arXiv.1907.04443.
- Stanislav Djorgovski. Generalized SETI in a Virtual Observatory. In Guillermo Lemarchand and Karen Meech, editors, *Bioastronomy 99*, volume 213 of *Astronomical Society of the Pacific Conference Series*, page 519, January 2000.
- Daniel Giles and Lucianne Walkowicz. Systematic serendipity: a test of unsupervised machine learning as a method for anomaly detection. , 484(1):834–849, March 2019. doi: 10.1093/mnras/sty3461.

⁴<https://astro.washington.edu/uw-planetarium>

⁵<https://astro.washington.edu/uw-mobile-planetarium>

⁶<https://huskylink.washington.edu/organization/leaguoia/>

⁷<https://astro.washington.edu/jacobsen-observatory>

- 171 A. Goodman, J. Fay, A. Muench, A. Pepe, P. Udomprasert, and C. Wong. WorldWide Telescope in Research and
172 Education. In P. Ballester, D. Egret, and N. P. F. Lorente, editors, *Astronomical Data Analysis Software and*
173 *Systems XXI*, volume 461 of *Astronomical Society of the Pacific Conference Series*, page 267, September 2012. doi:
174 10.48550/arXiv.1201.1285.
- 175 Laura Kreidberg. batman: BASic Transit Model cAlculation in Python. , 127(957):1161, November 2015. doi:
176 10.1086/683602.
- 177 Yuqian Liu and Maryam Modjaz. SuperNova IDentification spectral templates of 70 stripped-envelope core-collapse
178 supernovae. *arXiv e-prints*, art. arXiv:1405.1437, May 2014. doi: 10.48550/arXiv.1405.1437.
- 179 K. L. Malanchev, M. V. Pruzhinskaya, V. S. Korolev, P. D. Aleo, M. V. Kornilov, E. E. O. Ishida, V. V. Krushinsky,
180 F. Mondon, S. Sreejith, A. A. Volnova, A. A. Belinski, A. V. Dodin, A. M. Tatarnikov, S. G. Zheltoukhov, and
181 (The SNAD Team). Anomaly detection in the Zwicky Transient Facility DR3. , 502(4):5147–5175, April 2021. doi:
182 10.1093/mnras/stab316.
- 183 Konstantin Malanchev, Matwey V. Kornilov, Maria V. Pruzhinskaya, Emille E. O. Ishida, Patrick D. Aleo, Vladimir S.
184 Korolev, Anastasia Lavrukhina, Etienne Russeil, Sreevarsha Sreejith, Alina A. Volnova, Anastasiya Voloshina, and
185 Alberto Krone-Martins. The SNAD Viewer: Everything You Want to Know about Your Favorite ZTF Object. , 135
186 (1044):024503, February 2023. doi: 10.1088/1538-3873/acb292.
- 187 Bill Paxton, Lars Bildsten, Aaron Dotter, Falk Herwig, Pierre Lesaffre, and Frank Timmes. Modules for Experiments
188 in Stellar Astrophysics (MESA). , 192(1):3, January 2011. doi: 10.1088/0067-0049/192/1/3.
- 189 A. B. A. Queiroz, F. Anders, B. X. Santiago, C. Chiappini, M. Steinmetz, M. Dal Ponte, K. G. Stassun, L. N. da Costa,
190 M. A. G. Maia, J. Crestani, T. C. Beers, J. G. Fernández-Trincado, D. A. García-Hernández, A. Roman-Lopes, and
191 O. Zamora. StarHorse: a Bayesian tool for determining stellar masses, ages, distances, and extinctions for field stars.
192 , 476(2):2556–2583, May 2018. doi: 10.1093/mnras/sty330.
- 193 Sofia Z. Sheikh. The Nine Axes of Merit for Technosignature Searches. *International Journal of Astrobiology*, 19(3):
194 237–243, June 2020. doi: 10.1017/S1473550419000284.
- 195 Jose I. Vines and James S. Jenkins. ARIADNE: measuring accurate and precise stellar parameters through SED fitting.
196 , 513(2):2719–2731, June 2022. doi: 10.1093/mnras/stac956.
- 197 Jason T. Wright and Steinn Sigurdsson. Families of Plausible Solutions to the Puzzle of Boyajian’s Star. , 829(1):L3,
198 September 2016. doi: 10.3847/2041-8205/829/1/L3.
- 199 Jason T. Wright, Kimberly M. S. Cartier, Ming Zhao, Daniel Jontof-Hutter, and Eric B. Ford. The Search for
200 Extraterrestrial Civilizations with Large Energy Supplies. IV. The Signatures and Information Content of Transiting
201 Megastructures. , 816(1):17, January 2016. doi: 10.3847/0004-637X/816/1/17.
- 202 A. Zanella, C. M. Harrison, S. Lenzi, J. Cooke, P. Damsma, and S. W. Fleming. Sonification and sound design
203 for astronomy research, education and public engagement. *Nature Astronomy*, 6:1241–1248, November 2022. doi:
204 10.1038/s41550-022-01721-z.
- 205 P. Zečević, C. T. Slater, M. Jurić, A. J. Connolly, S. Lončarić, E. C. Bellm, V. Z. Golkhou, and K. Suberlak. AXS: A
206 Framework for Fast Astronomical Data Processing Based on Apache Spark. , 158:37, July 2019.

Phase	Timeline	Goals and Milestones
Pilot Study	Year 1	<ul style="list-style-type: none"> • Assemble tools and data products used for the program • ApJL paper submission for <i>Search for F-Type Main Sequence KIC 8462852 Analogs with the Zwicky Transient Facility and Gaia DR3</i> (Tzanidakis et al. in-prep) • Target of Opportunity (ToO) proposals with APO/Gemini S (for later), ATA radio observatory follow-up ... • Compile a census of anomalous stellar variability patterns: <i>duration, depth, light curve shape, color</i> with ground based data • Applying to NSF for the Search Anomalous Stellar Variability in the Rubin Era • Can we establish an upper limit with ZTF a boundary condition/limits we can use the alert framework to find anomalous events in <i>gri</i>. Needs to be deployed on the alert brokers in ~ 2 years.
Algorithms & Limits	Year 2	<ul style="list-style-type: none"> • Collaboration with Rubin/LSST OpSims simulations to place meaningful upper limits on the search for optical technosignatures through stellar variability • Estimate upper limit of detectability of megastructures induced variability with most modern surveys: ZTF, SDSS, LSST, and Gaia • How would our detection rates change if we wanted to work with alerts? Can we enable the first real-time search for technosignatures through alert-based classifiers? • Signal synchronization upper limits with LSST and other surveys. <i>Spatio-Temporal Signal Synchronization for the Search of Technosignatures using Time-Domain Surveys. Deploying the ellipsoid at scale for Rubin. ElaSTICC technosignature for Andy Nilipour et al. (including Davenport, Tzanidakis, Croft)</i> • Development of open-source tool Astra-Var extensive time series tools for enabling robust anomaly searches with sparse multi-band data • Student project ideas for undergraduate researchers & post-bacc., explore some of the exotic ideas for spatiotemporal ideas, look for VASCO's...
Systematic Search	Year 3-4	<ul style="list-style-type: none"> • The big search. Using our pilot program, limits, and newly developed algorithms let's search for technosignatures: ZTF DR19, SDSS, Gaia DR4, and LSST • Enable auxiliary stellar variability estimates. After the completion of this project, our study will have the largest temporal characterization of stellar variability with a sampling cadence of 3-7 days. Could be also used for future studies to enable robust classification and for more anomaly searching in alerts.